

Name of journal: *World Journal of Critical Care Medicine*

ESPS Manuscript NO: 11891

Columns: Editorial

Intensive care performance: How should we monitor performance in the future?

Timmers TK *et al.* Intensive care performance

Tim K Timmers, Michiel HJ Verhofstad, Karl GM Moons, Luke PH Leenen

Tim K Timmers, Luke PH Leenen, Department of Surgery, University Medical Center Utrecht, 3508 GA Utrecht, The Netherlands

Michiel HJ Verhofstad, Department of Surgery, Erasmus Medical Center Rotterdam, 3000 CA Rotterdam, The Netherlands

Karl GM Moons, Department of Clinical Epidemiology, Julius Centre for Health Sciences and Primary Care, 3508 GA Utrecht, The Netherlands

Author contributions: All authors contributed to this paper.

Correspondence to: Tim K Timmers, MD, PhD, Department of Surgery, University Medical Center Utrecht, PO Box 85500, 3508 GA Utrecht, The Netherlands.
tk.timmers@gmail.com

Telephone: +31887559882

Fax: +31887555555

ACKNOWLEDGMENTS AND FUNDING

We received no source of funding for our study.

The authors declared that they have no competing interests.

Each author certifies that he has participated sufficiently in the intellectual content, the analysis of data, if applicable, and the writing of the Work to take public responsibility for it. Each has reviewed the final version of the Work, believes it represents valid work, and approves it for publication. We herewith state that all authors provided an equal contribution to this study and act in concordance to the 'Uniform Requirements'.

Abstract

Intensive care faces economic challenges. Therefore evidence proving both effectiveness and efficiency, i.e. cost-effectiveness, of delivered care is needed. Today, the quality of care is an important issue in the health care debate. How do we measure quality of care, and how accurate and representative is this measurement? In the following report several topics, which are used for the evaluation of ICU performance, are discussed: (1) The use of general outcome prediction models to assess/ quantify the risk of patients who are admitted to our ICUs in an increasing variety in case mix for all the different intensive care units. Together with three major limitation; (2) As critical care outcomes research becomes a more established entity mortality is now only one of many endpoints that are of relevance. Mortality is a limited outcome when assessing critical care performance, while patient interest in QoL outcomes is of relevance; and (3) The Quality Indicators Committee of the Society of Critical Care Medicine recommended that short-term readmission is a leading performance indicator of the quality of intensive care medicine.

Key words: Intensive care performance; Quality of Care; Critical Care; Intensive Care Medicine

Core tip: Variations in case-mix, ICU demographics, clinical and non-clinical factors not addressed by the present severity of illness scores must be quantified to improve the accuracy of future prediction models. A completely different benefit, in the use of HrQoL as a performance-benchmark, could be the possibility of follow-up evaluation of the patients' health status after ICU or hospital discharge. The moment when outcome research can predict the short-term (ICU discharge) QoL of a critically ill patient during the first 24 h of ICU admission; It will give physicians and health care policy makers an up-to-date and reliable evaluation of quality of care in the ICU during the upcoming years.

INTRODUCTION

The intensive care unit (ICU) is a hospital unit providing continuous surveillance and highly specialized care to critically ill patients, either medical or surgical. Patient's conditions are life-threatening and require comprehensive care. Established approximately five decades ago, the ICU is now an essential part of hospital care. It presents itself as the discipline that aims to help patients with extended needs of care and organ support.

Intensive care faces economic challenges. Therefore, evidence proving both effectiveness and efficiency, i.e. cost-effectiveness, of delivered care is needed. ICUs consume a significant proportion of health care resources, accounting for up to 20% of a hospital's cost [1-6]. By 2005, critical care medicine (CCM) costs in the United States were estimated to be \$81.7 billion accounting for 4.1% of the national health expenditures, 0,66% of the gross domestic product. The US spends 15% of the gross domestic product on healthcare (9-11% in Germany, France and Canada; 7-8% in Spain and the United Kingdom). Intensive care costs are predicted to grow all over the developed world [2,5,7-14].

Today, quality of care is an important issue in the health care debate. All countries struggle to optimize quality of care while minimizing costs. Evaluation of clinical performance is a prerequisite for the assessment of both the effectiveness and efficiency of care [15] and, therefore, several questions arise: How do we measure quality of care, and how accurate and representative is this measurement?

The goal of intensive care medicine is to achieve the best outcome for critically ill patients, and usually this is accompanied by the use of very complex care [16]. All patients carry both an intrinsic risk (disease-related) and an extrinsic risk (care-related) at the same time [16,17]. There is an ever-increasing recognition of the wide variation in the quality of care across ICUs and its effect on outcome. Indicators to evaluate the quality of care are increasingly being used and focus on patient outcome

[18]. Finding a reliable method to quantify the performance of single ICUs has been a difficult quest in the last 30 years.

Outcome prediction models: shall we continue in the same way?

Each new development in critical care treatment over the past 30 years has been implemented to improve the quality of care. Therefore, the extrinsic risks that patients carry should be as low as possible. Ideally, quality of care performance research should give more information about the extrinsic risks rather than the intrinsic ones. At this moment, ICU performance evaluation is becoming increasingly difficult because of the presence of an increasing variety in patient case mix for all the different intensive care units. Since the development of prediction mortality models in the early 1980s, physicians have tried to normalize certain different ICU populations through the use of severity of illness measurements. At the time that a general outcome prediction model (GOPM) had been developed the intrinsic risk had been adjusted in such a way that performance mainly illuminates the extrinsic risk factors. Most published approaches, concerning the evaluation of ICU performance, adopt more or less the same procedures: the development of a GOPM and its calibration in a suitable database. Such models are then applied to different cohorts of ICU patients, and the comparison of the predicted number of deaths with the actual number is used as a reference for the clinical performance of the unit [15]. For more than 30 years, outcome research in critical care relied heavily on these risk adjustment methods (GOPM) to assess and quantify the risk of patients who are admitted to our ICUs. Using several GOPMs this methodology has become the 'gold standard' to compare ICUs across different geographical areas, or within a specific individual nation, or other specific subgroups. Several risk adjustment systems have been developed or have undergone an update and are used in daily practice. In the use of general outcome prediction models, several limitations should be considered.

1. Most systems give a single estimate, known as the standardized mortality ratio (SMR). A single estimate considers that the performance of an ICU is constant over the whole spectrum of the severity of illness [19]. In other words, an ICU with a 'good' performance (low SMR) is assumed to be uniformly good for both low-risk and high-risk patients; likewise, an ICU with a 'bad' performance (high SMR) is assumed to be uniformly bad. However, since performance can vary not only between ICUs but also within the same unit across patients and doctors, this assumption is likely not true [15]. Several studies have provided conclusive evidence that the clinical performance of ICUs may vary over the spectrum of severity of illness [15,16,20-23].

2. It is unknown whether variations in SMR reflect quality of care or case mix differences. Debate continues whether higher than predicted mortality (high SMR) is a warning about the quality of care or rather reflects a difference of case mix between hospitals [24]. In the past, GOPMs have been revised or even updated to newer versions to predict expected death more accurately. However, before a new GOPM version is used, many years have elapsed. Although, the newer third and fourth versions of the APACHE prognostic model were developed many years ago [25-27], the APACHE II score is still one of the prediction models most widely used [12,28].

3. There is no consensus which GOPM must be used for which type of ICU (general mixed unit, specialized unit, or even in different sub-populations). For critical care physicians there are three overall GOPMs for predicting overall mortality and used for performance evaluations: the APACHE model [25-27,29], the MPM system [30-32] and the SAPS model [22,23,33-35]. These scoring systems differ in the choice and the relative weight given to patient characteristics and physiological parameters [18]. Quality of care performance evaluation should be done with the same - and ideally most reliable - outcome prediction model for each intensive care unit. Because there is no consensus which GOPM should be used, they seem to be used randomly. Within the Netherlands, this last item has been addressed: Since the year 2008, all 61

participating ICUs in the NICE registry started using the APACHE IV prognostic model.

The quality of intensive care performance.

Until today, one of the most used ICU performance measurements is the standardized mortality ratio (SMR). The SMR has been developed in a period that the evaluation of quality of care was done exclusively through primary patient outcome (short-term mortality). Some authors have evaluated the use of SMR as an indicator of ICU quality of care and debated its specific relevance [17,36,37]. The SMR value gives insight in the observed mortality compared with the associated predicted mortality, but it does not give insight in the health status of these patients.

As critical care outcomes research becomes a more established entity mortality is now only one of many endpoints that is of relevance, and mortality is a limited outcome evaluation method when assessing critical care performance. The health-related quality of life (HrQoL), defined as the degree to which a patient's health status affects the subjective evaluation of his or her satisfaction with life, seems to be a better indicator, especially from the patient-centred view [38]. ICU and hospital survival will always have an important role in the evaluation of performance at the moment different units or hospitals are being benchmarked. Consequently, in the last decennia the quality of life (QoL) has gained great interest when not only physicians but also the patients' relatives mention patient outcome. Therefore, QoL clearly challenges survival whenever we address secondary (long-term) patient outcome.

Difficulties are being foreseen when using health status as a performance-benchmark [39], because of the great diversity in intrinsic risk that patients carry in all the different ICUs (i.a. specialized units, general mixed units). How should we use health status as performance-benchmark? Should we cross-section the mean health status of a given cohort against the general population norm, or must we compare individual outcome with individual pre-admission values? The latter will invariably

provide more patient oriented and thus clinically relevant outcome values, but also result in an administrative burden. A third possibility is to compare such an individual QoL value with a predicted individual health status.

The ability to predict a patient's QoL after ICU admission could be useful in many ways. Firstly, it could help patients and their relatives to make decisions. Secondly, it could help families to prepare themselves to care for the patient after hospital discharge. Thirdly, it could help critical care physicians to give useful information, avoid unrealistic expectations, and - possibly - could help in making treatment decisions. Fourthly, it could help society to realize in which ICUs patients have a good chance of recovery and give health policy makers and insurance companies the insight in the needs of ICUs [39,40].

A completely different benefit, in the use of HrQoL as a performance-benchmark, could be the possibility of follow-up evaluation of patients' health status after ICU or hospital discharge. Post-ICU patients are known to express a reduced HrQoL compared to the general population. It is still not clear to what extend and how long this reduced HrQoL persists, although this effect may be long-lasting [41]. Therefore, a continuous survey as part of regular after care for each individual patient, would be the ideal way to investigate this, and gives the possibility to better manage patients in which HrQoL does not increase as expected.

Readmission to the ICU: Can we predict patients at risk for readmission?

The Quality Indicators Committee of the Society of Critical Care Medicine recommended that readmission within 48 hours is a leading performance indicator of the quality of intensive care medicine [42]. Readmitted patients are most often the sickest in the ICU; therefore, it is an unexpected and unfavourable event for the patient, and is associated with a more severe outcome [43-52]. Moreover, a strategy to reduce premature discharges in patients at high risk of in-hospital death could result in a reduction of post-ICU mortality (Daly *et al.*: 39% reduction in mortality) [52-54]. In times of great pressure on the ICU capacity, should we not be more careful in

deciding which patient may be discharged and who has a greater risk of readmission? Ideally, such decisions are made on sound criteria, rather than subjective parameters. In the last 10 years several authors have proven that it is difficult to analyze and predict readmittance-risk for ICU patients in general [44-47]. Various authors concluded that patients readmitted to the ICU had a higher severity of illness score at the time of initial ICU discharge compared to single ICU admission patients [45,46,55]. Ideally perhaps, severity of illness is scored on a daily base and discharge is initiated from these values. Unfortunately, these severity of illness scores have not been validated after the first 24 hours of ICU admission. The Sequential (Sepsis-related) Organ Failure Assessment score (SOFA score), is used to track a patient's status during the admission in the ICU (also validated to be used after 24 hours). The SOFA score is a scoring system to determine the extent of a person's organ function or rate of failure [56-59]. This particular score has been validated to predict ICU mortality [60]. Nevertheless, the possible association with readmission has not been evaluated as yet. Currently, there are hardly any systematic studies of how daily severity of illness score changes from admission to initial discharge predict ICU readmission [28,47]. Besides the severity of illness score there is also an association between nursing workload and post-ICU mortality [61,62]. The Therapeutic Intervention Scoring System (TISS) has been widely applied to assess workload and resource allocation in intensive care, measuring treatment intensity [63]. Consequently, attempts have been made to use TISS scores to categorize the level of care that patients require, even to evaluate the care required after ICU discharge [64,65]. Several authors have shown an association of the TISS value of the last ICU day with post-ICU mortality [61,62,66,67], and therefore, indirectly, the association with ICU-readmission. Smith *et al.* concluded in their research that the mean TISS scores in patients readmitted to the ICU were significantly higher than in patients who did not require readmission [62].

Since a couple of years Spanish physicians have shown great interest in this topic and developed the Sabadell score system, a modification of the McCabe score [68-70]. They

have validated the applicability of the Sabadell score as a system for classifying patients' ward survival at discharge from the ICU [71]. They even found an association of the Sabadell score with ICU readmission. Unfortunately, the lack of reliable predictors of ICU readmission precludes the clinical usefulness of this variable. However, this information may improve the ability to predict the readiness for discharge for individual patients and improve the efficiency of intensive care units. Would critical care physicians have more information about patients' disease-status when they use a combination of several systems (TISS, severity of illness score and Sabadell score) as prediction measurement for ICU discharge readiness? This value could also give an indication if the patient could be discharged to the normal ward or that he should first be admitted to a step-down unit (high dependency unit).

Hospital death rates would be particularly useful if patients and physicians could use the statistics for a given diagnosis to select a hospital that offers the best prospect of survival. If the data are only partially corrected for differences in the health status of patients, then they must be used with caution [19]. Variations in case-mix, ICU demographics, clinical and non-clinical factors not addressed by the present severity of illness scores must be quantified to improve the accuracy of future prediction models. If the variation between ICUs is important, it will impair the stability of the equations used to compute predicted mortality and preclude the use of indirect standardization in the evaluation of differences between ICUs. These GOPMs consider the relation between performance and severity of illness as constant, although performance can vary within ICUs according to the degree of severity of illness in patients. Hypothetically, performance should be evaluated through the combination of survival (SMR) and the health status (QoL) at the time of discharge. As yet, this combination of both outcome measurements has not been used in a single benchmark value. Therefore, future research should focus on predicting quality of life together with an accuracy study of the TISS, severity of illness and the Sabadell scores to identify and weigh the specific variables for readmission. The

moment when outcome research can predict the short-term (ICU discharge) QoL of a critically ill patient during the first 24 hours of ICU admission; It will give physicians and health care policy makers an up-to-date and reliable evaluation of quality of care in the ICU during the upcoming years.

REFERENCES

1. Pronovost PJ, Angus DC, Dorman T, Robinson KA, Dremsizov TT, Young TL. Physician staffing patterns and clinical outcomes in critically ill patients: A systematic review. *J Am Med Assoc* 2002; 288: 2151-2162. [PMID: 12413375, doi: unknown].
2. Giovanni M, Scatto P, Scatto P. Economics and outcome in the intensive care unit. *Curr Opin Anaesthesiol.* 2009; 22: 232-6. [PMID: 19390250, doi: 10.1097/ACO.0b013e328323f9a8].
3. Moerer O, Plock E, Mgbor U, Schmid A, Schneider H, Wischnowsky MB, Burchardi H.. A German national prevalence study on the cost of intensive care: an evaluation from 51 intensive care units. *Crit Care Med* 2007; 11: 69-79. [PMID: 17594475, doi: unknown].
4. Chalfin DB. Cost-effectiveness analysis in health care. *Hosp Cost Manag Account* 1995; 7: 1-8. [PMID: 10143321, doi: unknown].
5. Zilberberg M, Luippold R. Prolonged acute mechanical ventilation, hospital resource utilization, and mortality in the United States. *Crit Care Med* 2008; 36: 726-730. [PMID: 18209667, doi: 10.1097/CCM.0B013E31816536F7].
6. Halpern NA, Bettes L, Greenstein R. Federal and nationwide intensive care units and healthcare costs: 1986-1992. *Crit Care Med* 1994; 22: 2001-7. [PMID: 7988140, doi: unknown].
7. Halpern NA, Pastores SM, Greenstein RJ. Critical care medicine in the United States 1985-2000: an analysis of bed numbers, use, and costs. *Crit Care Med* 2004; 32: 1254-9. [PMID: 15187502, doi: unknown].
8. Halpern NA, Pastores SM. Critical care medicine in the United States 2000-2005: an analysis of bed numbers, occupancy rates, payer mix, and costs. *Crit Care Med* 2010; 38: 65-71. [PMID: 19730257, doi: 10.1097/CCM.0b013e3181b090d0].
9. Halpern NA. Can the costs of critical care be controlled? *Curr Opin Crit Care* 2009; 15: 591-6. [PMID: 19823083, doi: 10.1097/MCC.0b013e328332f54f].
10. Cohen NH. Critical care medicine in the United States: What we know, what we do not, and where we go from here. *Crit Care Med* 2010; 38: 304-306. [PMID: 20023473, doi: 10.1097/CCM.0b013e3181b4a2b6].

11. Kahn JM, Angus DC. Reducing the cost of critical care: new challenges, new solutions. *Am J Respir Crit Care Med* 2006; 174: 1167-1168. [PMID: 17110650, doi: unknown].
12. Vincent JL, Opal SM, Marshall JC. Ten reasons why we should NOT use severity scores as entry criteria for clinical trials or in our treatment decisions. *Crit Care Med* 2010 Jan; 38: 283-287. [PMID: 19730252, doi: 10.1097/CCM.0b013e3181b785a2].
13. Organisation for Economics Co-operation and Development: Health at a Glance: OECD indicators. 2005.
14. Wunsch H, Angus DC, Harrison DA, Collange O, Fowler R, Hoste EA, de Keizer NF, Kersten A, Linde-Zwirble WT, Sandiumenge A, Rowan KM. Variation in critical care services across North America and Western Europe. *Crit Care Med* 2008; 36: 2787-2793. [PMID: 18766102, doi: 10.1097/CCM.0b013e318186aec8].
15. Moreno RP, Hochrieser H, Metnitz B, Bauer P, Metnitz PGH. Characterizing the risk profiles of intensive care units. *Intensive Care Med* 2010; 36: 1207-1212. [PMID: 20306015, doi: 10.1007/s00134-010-1852-2].
16. Metnitz PG. From risk assessment to risk management. *Crit Care Med* 2010; 16: 475-476. [PMID: 20827074, doi: 10.1097/MCC.0b013e32833e81ed].
17. Teres D, Lemeshow S. Why severity models should be used with caution. *Crit Care Clin* 1994; 10: 93-115. [PMID: 8118735, doi: unknown].
18. De Lange DW, Dusseljee J, Brinkman S, van Berkel G, van Maanen R, Bosman RJ, Joore H, de Keizer NF, van der Voort PHJ, de Waal R, Wesselink RMJ, de Jonge E. Severity of illness and outcome in ICU patients in the Netherlands: results from the NICE registry 2006-2007. *Nether J Crit Care* 2009; 13: 16-22. [PMID: Unknown, doi: unknown].
19. Moreno R, Apolone G. Impact of different customization strategies in the performance of a general severity score. *Crit Care Med* 1997; 25: 2001-2008. [PMID: 9403750, doi: unknown].
20. Moreno RP, Bauer P, Metnitz PGH. Characterizing performance profiles of ICUs. *Crit Care Med* 2010; 16: 477-481. [PMID: 20613499, doi: 10.1097/MCC.0b013e32833cc925].
21. Civetta JM. Scoring systems: Do we need a different approach? *Crit Care Med* 1991; 19: 1460-1461. [PMID: 1959362, doi: unknown].
22. Metnitz PG, Moreno RP, Almeida E, Jordan B, Bauer P, Campos RA, Iapichino G, Edbrooke D, Capuzzo M, Le Gall JR; SAPS 3 Investigators. SAPS 3 – From evaluation of the

patient to evaluation of the intensive care unit. Part 1: Objectives, methods and cohort description. *Intensive Care Med* 2005, 31: 1336-1344. [PMID: 16132893, doi: unknown].

23. Moreno RP, Metnitz PGH, Almeida E, Jordan B, Bauer P, Campos RA, Iapichino G, Edbrooke D, Capuzzo M, Le Gall JR; SAPS 3 Investigators. SAPS 3 – From evaluation of the patient to evaluation of the intensive care unit. Part 2: Development of a prognostic model for hospital mortality at ICU admission. *Intensive Care Med* 2005, 31: 1345-1355. [PMID: 16132892, doi: unknown].

24. Green J, Passman LJ, Wintfeld N. Analyzing hospital mortality: The consequences of diversity in patient mix. *J Am Med Assoc* 1991; 265 (14): 1849-1853. [PMID: 1900894, doi: unknown].

25. Knaus WA, Draper EA, Wagner DP, Zimmerman JE. APACHE II: a severity of disease classification system. *Crit Care Med* 1985, 13: 818-829. [PMID: 3928249, doi: unknown].

26. Knaus WA, Wagner DP, Draper EA, Zimmerman JE, Bergner M, Bastos PG, Sirio CA, Murphy DJ, Lotring T, Damiano A. The APACHE III prognostic system. Risk prediction of hospital mortality for critically ill hospitalised adults. *Chest* 1991; 100: 1619-1636. [PMID: 1959406, doi: unknown].

27. Zimmerman JE, Kramer AA, McNair DS, Malila FM. Acute Physiology and Chronic Health Evaluation (APACHE) IV: Hospital mortality assessment for today's critically ill patients. *Crit Care Med* 2006; 34: 1297-1310. [PMID: 16540951, doi: unknown].

28. Cohen NH. The real reasons not to rely on severity scores. *Crit Care Med* 2010; 38: 334-335. [PMID: 20023491, doi: 10.1097/CCM.0b013e3181b9d3e4].

29. Knaus WA, Zimmerman JE, Wagner DP, Draper EA, Lawrence DE. APACHE — Acute Physiology and Chronic Health Evaluation: A physiologically based classification system. *Crit Care Med* 1981; 9: 591-597. [PMID: 7261642, doi: unknown].

30. Lemeshow S. Refining intensive care unit outcome prediction by using changing probabilities of mortality. *Crit Care Med* 1988; 16: 470-477. [PMID: 3359785, doi: unknown].

31. Lemeshow S, Teres D, Klar J, Avrunin JS, Gehlbach SH, Rapoport J. Mortality Probability Models (MPM II) based on an international cohort of intensive care unit patients. *J Am Med Assoc* 1993, 270: 2478-2486. [PMID: 8230626, doi: unknown].

1. Higgins TL, Teres D, Copes WS, Nathanson BR, Stark M, Kramer AA. Assessing contemporary intensive care unit outcome: An updated Mortality Probability Admission Model (MPM0-III). *Crit Care Med* 2007; 35: 827-835. [PMID:
2. Le Gall J-R, Loirat P, Alperovitch A, *et al.* A simplified acute physiology score for ICU patients. *Crit Care Med* 1984; 12: 975-977. [PMID:
3. Le Gall J-R, Lemeshow S, Saulnier F: A new Simplified Acute Physiology Score (SAPS II) based on a European/North American multi centre study. *J Am Med Assoc* 1993, 270: 2957-2963. [PMID:
4. Le Gall J-R, Neumann A, Hemery F, *et al.* Mortality prediction using SAPS II: an update for French intensive care units. *Crit Care Med* 2005, 9: 645-652. [PMID:
5. Sherck JP, Shatney CH. ICU Scoring Systems do not allow predictions of patient outcomes or comparison of ICU performance. *Crit Care Clin* 1996; 12: 515-523. [PMID:
6. Boyd O, Grounds RM. Can standardized mortality ratio be used to compare quality of intensive care unit performance? *Crit Care Med* 1994; 22: 1706-1709. [PMID:
7. Capuzzo M., Moreno RP, Jordan B, Bauer P, Alvisi R, Metnitz PGH. Predictors of early recovery of health status after intensive care. *Intensive Care Med* 2006; 32: 1832-8.
8. Valentin A. The importance of risk reduction in critically ill patients. *Crit Care Med* 2010; 16: 482-486. [PMID:
9. Angus DC. Caring for the critically ill patient: Challenges and opportunities. *J Am Med Assoc* 2007; 298: 456-458. [PMID:
10. Timmers TK, Verhofstad MHJ, Moons KGM, van Beeck EF, Leenen LPH. Long-term (>6 years) quality of life after surgical intensive care admission. *Arch Surg* 2011; 146: 412-418. [PMID:
11. Angus DC. Grappling with intensive care unit quality: does the readmission rate tell us anything? *Crit Care Med* 1998; 26: 1779-1780. [PMID:
12. Metnitz PG, Fieux F, Jordan B, Lang T, Moreno R, Gall JR. Critically ill patients readmitted to intensive care units – lessons to learn? *Intensive Care Med* 2003; 29: 241-248. [PMID:

13. Chen LM, Martin CM, Keenan SP, Sibbald WJ. Patients readmitted to the intensive care unit during the same hospitalization: clinical features and outcomes. *Crit Care Med* 1998; 26: 1834-1841. [PMID:
14. Alban RF, Nisim AA, Ho J, Nishi GK, Shabot MM. Readmission to surgical intensive care increases severity-adjusted patient mortality. *J Trauma* 2006; 60: 1027-1031. [PMID:
15. Rosenberg AL, Watts C. Patients readmitted to ICUs: A systematic review of risk factors and outcomes. *Chest* 2000; 118: 492-502. [PMID:
16. Rosenberg AL, Hofer TP, Hayward RA, Strachan C, Watts CM. Who bounces back? Physiologic and other predictors of intensive care unit readmission. *Crit Care Med* 2001; 29: 511-518. [PMID:
17. Nishi GK, Suh RH, Wilson MT, Cunneen SA, Margulies R, Shabot MM. Analysis of causes and prevention of early readmission to surgical intensive care. *Am Surg* 2003; 69: 913-917. [PMID:
18. Lee HF, Lin S-C, Lu C-L, Chen CF, Yen M. Revised Acute Physiology and Chronic Health Evaluation score as a predictor of neurosurgery intensive care unit readmission: A case-controlled study. *J Crit Care* 2010; 25: 294-299. [PMID:
19. Kaben A, Correa F, Reinhart K, Settmacher U, Gummert J, Kalff R, Sakr Y. Readmission to a surgical intensive care unit: incidence, outcome and risk factors. *Crit Care* 2008; 12: R123. [PMID:
20. Chan K-S, Tan C-K, Fang C-S *et al.* Readmission to the intensive care unit: an indicator that reflects the potential risks of morbidity and mortality of surgical patients in the intensive care unit. *Surg Today* 2009; 39: 295-299. [PMID:
21. Afessa B, Keegan MT, Hubmayr RD *et al.* Evaluating the performance of an institution using an intensive care unit benchmark. *Mayo Clin Proc* 2005; 80: 174-180. [PMID:
22. Daly K, Beale R, Chang RW. Reduction in mortality after inappropriate early discharge from intensive care unit: logistic regression triage model. *BMJ* 2001; 322: 1274-1276. [PMID:

23. Metnitz PG, Fieux F, Jordan B, et al. Critically ill patients readmitted to intensive care units – lessons to learn? *Intensive Care Med* 2003; 29: 241-248. [PMID:
24. Durbin CG, Kopel RF. A case-control study of patients readmitted to the intensive care unit. *Crit Care Med* 1993; 21: 1547-1553. [PMID:
25. Vincent JL, Moreno R, Takala J, et al. The SOFA (Sepsis-related Organ Failure Assessment) score to describe organ dysfunction/failure. On behalf of the Working Group on Sepsis-Related Problems of the European Society of Intensive Care Medicine. *Intensive Care Med* 1996; 22: 707-710. [PMID:
26. Vincent JL, de Mendonça A, Cantraine F, et al. Use of the SOFA score to assess the incidence of organ dysfunction/failure in intensive care units: results of a multicenter, prospective study. Working group on “sepsis-related problems” of the European Society of Intensive Care Medicine. *Crit Care Med* 1998; 26: 1793-1800. [PMID:
27. Moreno R, Vincent JL, Matos R, et al. The use of maximum SOFA score to quantify organ dysfunction/failure in intensive care. Results of a prospective, multicentre study. Working Group on Sepsis related Problems of the ESICM. *Intensive Care Med* 1999; 25: 686-696. [PMID:
28. de Mendonça A, Vincent JL, Suter PM, et al. Acute renal failure in the ICU: risk factors and outcome evaluated by the SOFA score. *Intensive Care Med* 2000; 26: 915-921. [PMID:
29. Ferreira FL, Bota DP, Bross A, Mélot C, Vincent JL. Serial evaluation of the SOFA score to predict outcome in critically ill patients. *J Am Med Assoc* 2001; 286: 1754-1758. [PMID:
30. Moreno R, Miranda DR, Matos R, Fevereiro T. Mortality after discharge from intensive care: the impact of organ system failure and nursing workload use at discharge. *Intensive Care Med* 2001; 27: 999-1004. [PMID:
31. Smith L, Orts CM, O’Neil I, et al. TISS and mortality after discharge from intensive care. *Intensive Care Med* 1999; 25: 1061-1065. [PMID:

32. Cullen DJ, Civetta, JM, Briggs BA, Ferrara LC. Therapeutic Intervention Scoring System: a method for quantitative comparison of patient care. *Crit Care Med* 1976; 2: 57-60. [PMID:
33. Keene AR, Cullen DJ. Therapeutic Intervention Scoring System: update 1983. *Crit Care Med* 1983; 11: 1-3. [PMID:
34. Malstam J, Lind L. Therapeutic Intervention Scoring System (TISS) – a method for measuring workload and calculating costs in the ICU *Acta Anaesthesiol Scand* 1992; 36: 758-763. [PMID:
35. Capuzzo M, Moreno RP, Alvisi R. Admission and discharge of critically ill patients. *Crit Care Med* 2010; 16: 499-504. [PMID:
36. Beck DH, McQuillan P, Smith GB. Waiting for the break of dawn? The effects of discharge time, discharge TISS scores and discharge facility on hospital mortality after intensive care. *Intensive Care Med* 2002; 28: 1287-1293. [PMID:
37. Fernandez R, Baigorri F, Navarro G, Artigas A. A modified McCabe score for stratification of patients after intensive care unit discharge: the Sabadell score. *Crit Care Med* 2006; 10: R179. [PMID:
38. <http://www.sabadellscore.es/>
39. McCabe WR, Jackson GG. Gram negative bacteremia: I. Etiology and ecology. *Arch Intern Med* 1962, 110: 845-847. [PMID:
40. Fernandez R, Serrano JM, Umaran I, Abizanda R, Carrillo A, Lopez-Pueyo MJ, Rascado P, Balerdi B, Suberviola B, Hernandez G. Ward mortality after ICU discharge: a multicenter validation of the Sabadell score. *Intensive Care Med* 2010; 36: 1196-1201. [PMID: