

# **The prevalence of cam and pincer morphology and its association with development of hip osteoarthritis**

P. van Klij<sup>1</sup>, J. Heerey<sup>2</sup>, J.H. Waarsing<sup>1</sup>, R. Agricola<sup>1</sup>

<sup>1</sup>. Department of Orthopaedic Surgery, Erasmus University Medical Centre, Rotterdam, The Netherlands

<sup>2</sup>. La Trobe Sports and Exercise Medicine Research Centre, School of Allied Health, La Trobe University, Bundoora, Victoria, Australia

The authors affirm that they have no financial affiliation (including research funding) or involvement with any commercial organization that has a direct financial interest in any matter included in this manuscript.

## **Correspondence:**

Pim van Klij

‘s-Gravendijkwal 230

3000CA Rotterdam

The Netherlands

[p.vanklij@erasmusmc.nl](mailto:p.vanklij@erasmusmc.nl)

## **Synopsis**

Our understanding of femoroacetabular impingement (FAI) syndrome is slowly improving. The number of studies on all aspects (aetiology, prevalence, pathophysiology, natural history, treatment, and preventative measures) of FAI syndrome has grown exponentially over the past few years. This commentary provides the latest updates on the prevalence of cam and pincer hip morphology and its relationship with development of hip osteoarthritis (OA). Cam and pincer morphology is highly prevalent in the general population and in this paper is presented for different subgroups based on: age, sex, ethnicity, and athletic activity. Methodological issues in determining prevalence of abnormal hip morphology are also discussed. Cam morphology has been associated with development of hip OA while the association between pincer morphology and hip OA is much less clear. Results from reviewed studies as well as remaining gaps in literature on this topic are critically discussed and put into perspective for the clinician.

**Key words:** aetiology, cam, FAI syndrome, hip, impingement, pincer, osteoarthritis

Femoroacetabular impingement (FAI) syndrome has recently been defined by authors of an international consensus statement as “a motion-related clinical disorder of the hip with a triad of symptoms, clinical signs, and imaging findings”.<sup>26</sup> They also described the most commonly seen symptoms and clinical signs. The primary symptom of FAI syndrome is motion-related or position-related pain in the hip or groin. Pain may also be felt in the back, buttock, or thigh. In addition to pain, patients may also describe clicking, catching, locking, stiffness, restricted range of motion, or giving way. Diagnosis of FAI syndrome does not depend on a single sign. The flexion, adduction, internal rotation (FADIR) test is most commonly used, and is sensitive but not specific. There is often limited hip motion, especially restricted internal rotation when in hip flexion.<sup>26</sup> Imaging findings, the focus of this manuscript, include the presence of cam

and/or pincer hip morphology. Cam hip morphology is characterized by a nonspherical femoral head while pincer morphology is defined as overcoverage of the acetabulum relative to the femoral head, which can be either global (bony overgrowth of the acetabulum or a deep socket) or focal (acetabular retroversion). This manuscript provides an overview of studies that report on the prevalence of cam and pincer morphology, as well as studies investigating the relationship between cam and pincer morphology and hip osteoarthritis. Future research direction for FAI syndrome will be discussed.

### **Prevalence of cam morphology**

A recent systematic review,<sup>16</sup> which included 30 studies, showed that the prevalence of cam morphology has yet to be defined in a truly overall population based cohort. The prevalence of cam morphology in this systematic review ranged from 5 to 75%. This high extent of prevalence variation among studies was based on population characteristics (age, sex, ethnicity, athletic activity, presence/absence of symptoms), the measures and concurrent threshold values used to quantify hip morphology, and the imaging technique.

#### *Cam morphology: age*

Cam morphology is less prevalent in adolescents than in adults and has been shown to gradually increase during skeletal growth.<sup>1,3,55,56,63-65</sup> Cam morphology can first be identified and starts to develop from the age of 12 years old<sup>1,55,63</sup>, with prevalence increasing with age until the completion of growth.<sup>3</sup> In addition, the extent of athletic activity during skeletal growth may increase the risk of cam morphology development.<sup>3,55,64</sup> Cam morphology is therefore an acquired phenomenon during the second growth spurt and highly influenced by exercise related loads applied to the hip during this phase.

### *Cam morphology: sex*

Cam morphology is probably more common in males. The prevalence of cam morphology in asymptomatic males ranges from 13.0 to 72.0% compared to 0 to 11.7% in asymptomatic women (**TABLE 1**).<sup>30,32,39,57</sup> Studies on symptomatic individuals are more inconsistent because of the selection bias related to symptomatic status. A study by Clohisy et al<sup>14</sup> showed an average prevalence of cam morphology of 47.6% in a symptomatic group of 1076 patients (55% women and 45% men) that underwent surgery for FAI syndrome. Symptomatology and functional limitations are pre-operatively significantly more severe in females compared with males.<sup>29,51</sup>

### *Cam morphology: ethnicity*

Mosler et al<sup>47</sup> identified a significantly lower prevalence of cam morphology amongst young East Asian (19%) professional soccer players, when compared to other ethnicities including Arabic, Black, Persian, and Whites in whom the prevalence ranged between 58% and 72%. Similarly, cam morphology prevalence has been shown to be lower in asymptomatic Chinese men and women compared to Caucasians in another article.<sup>68</sup> However, in contrast, another prevalence study of asymptomatic higher aged individuals report that East Asian populations have a high prevalence of cam morphology (45.3% of 1178 hips).<sup>44</sup>

### *Cam morphology: athletic activity*

In their systematic review and meta-analysis, Nepple et al<sup>52</sup> reported that professional athletes exhibit a higher prevalence of cam morphology relative to non-athletic individuals. The pooled prevalence of cam morphology in male athletes was 41% compared with 17% in male controls. In another systematic review<sup>20</sup>, the authors reported prevalence of cam morphology in up to

55% of male athletes compared with 23% in the general population. In their systematic review, Dickenson et al<sup>16</sup> reported prevalence of cam morphology in athletes ranging from 48 to 75%.

#### *Cam morphology: symptomatology*

It is currently unknown whether or not the presence of cam morphology by itself is associated with symptoms. Only one prospective study is available, which investigated 200 asymptomatic volunteers over a period of 4.4 years and showed that the presence of cam morphology resulted in a relative risk of 4.3 (95% confidence interval (CI) 2.3, 7.8) of developing hip pain.<sup>35</sup> Similarly a cross-sectional study found an association between an increased alpha angle (indicative of cam morphology) and prior or current athletic-related groin pain in 125 collegiate national football league prospects.<sup>37</sup> This is consistent with the results of another study which showed a relationship between cam morphology based on higher alpha angles and hip symptoms.<sup>8</sup> However, Gosvig et al<sup>24</sup>, studying a large non-athletic population consisting of 3202 individuals, showed no significant association between self-reported hip pain and cam morphology. Other studies also could not identify an association between symptoms and cam morphology.<sup>9,33,48</sup> When asymptomatic and symptomatic subgroups are compared, Mascarenhas et al<sup>41</sup> found a higher prevalence of cam morphology in symptomatic hips compared to asymptomatic hips. However, these studies consisted generally of less than 50 participants per subgroup.

#### **Prevalence of pincer morphology**

Pincer morphology is even more heterogeneously defined than cam morphology. However, similar to cam morphology, the prevalence of pincer morphology appears to vary across different subpopulations.

### *Pincer morphology: age*

Only a few studies have been published on how the prevalence of pincer morphology changes with age. A study on an asymptomatic paediatric and adolescent population of a mean age of 10.4 years, identified the presence of pincer morphology starting at 12 years of age.<sup>45</sup> In adolescents with an average age of 14.4 years old, Li et al<sup>40</sup> reported a prevalence of pincer morphology of 32.4%. Laborie et al<sup>36</sup>, in a study including 2081 young adults with an average age of 18.6 years, reported pincer morphology prevalence of 34.3% in men and 16.6% in women (**TABLE 2**).

### *Pincer morphology: sex*

Multiple studies have directly compared the incidence of pincer morphology between males and females, showing very little difference. Li et al<sup>40</sup> did not find a difference in prevalence of pincer morphology between asymptomatic males and females. Prevalences of 29.7% and 35.1% in males and females ( $P=.17$ ) were presented. Other studies showed conflicting results. A higher prevalence of pincer morphology in males was observed in the study of 2081 individuals by Laborie et al<sup>36</sup> who reported a prevalence of pincer morphology of 34% in males, compared to 17% in females ( $P<.001$ ). In contrast, coxa profunda was found to be significantly associated with female sex in 3 studies.<sup>15,17,28</sup> Two additional studies provided data on the prevalence of pincer morphology only in women, which ranged between 1 and 10%.<sup>33,39</sup> In comparison, the reported prevalence in males ranges between 3 and 66%.<sup>32,47</sup> There is also probably not a great difference in prevalence of pincer morphology between sexes in symptomatic individuals, based on a study by Nepple et al<sup>51</sup> who showed a prevalence of isolated pincer morphology in 56% of the males and 47% of females ( $P=.46$ ) undergoing FAI surgery.

### *Pincer morphology: ethnicity*

Less is known about the association between pincer morphology and ethnicity. The study of Mosler et al<sup>47</sup> compared the prevalence of pincer morphology (lateral center-edge angle (LCEA) >40°) between young soccer players with different ethnic backgrounds. No pincer morphology was found in white and East Asian soccer players. Arabic (3.6%), black (2.3%), and Persian soccer players (1.7%) showed also a low prevalence. Tannenbaum et al<sup>66</sup> did not find a difference in acetabular retroversion of pelvic specimens between African Americans and Caucasians. Several studies only investigated Asian persons, specifically Japanese, and found a prevalence of pincer morphology ranging from 7.4% to 37.4%.<sup>7,21,44,46</sup>

### *Pincer morphology: athletic activity*

The prevalence of pincer morphology in athletes is highly variable. Harris et al<sup>28</sup> investigated a group of elite ballet dancers and found a prevalence of 74%. In studies, which investigated soccer/football players, prevalence of pincer morphology ranged from 3 to 66%.<sup>22,32,47</sup> A study which combined different type of athletes (volleyball, soccer, and track & field), found a pincer morphology prevalence of 1%.<sup>33</sup> In elite ice hockey players, Lerebours et al<sup>38</sup> found a prevalence of pincer morphology of 59.8%. Two systematic reviews, by Frank et al<sup>20</sup> and Mascarenhas et al<sup>41</sup> found a prevalence of pincer morphology in athletes of 49.5% and 51.2%, respectively.

### *Pincer morphology: symptomatology*

Comparisons between symptomatic and asymptomatic subgroups were presented in a recent systematic review of Mascarenhas<sup>41</sup>, which included 60 studies. Pincer morphology prevalence in the asymptomatic subgroup, reported in only 1 study, was 57%. In symptomatic individuals, prevalence of pincer morphology was on average 28.5% (standard deviation (SD) ± 19.2)

across studies. The reported prevalence of pincer morphology in asymptomatic individuals in the systematic review by Frank<sup>20</sup> was 67% (range 61 – 76%). That systematic review which included 26 studies did not report on symptomatic individuals. These results are inconsistent with data from Gosvig et al<sup>25</sup> who reported lower prevalence of pincer morphology in men (15.2%) and women (19.4%) in a population-based study. A study by Ahn et al<sup>7</sup> showed pincer prevalence in asymptomatic males and females of 27% and 21%, respectively.

### **Relationship between cam morphology and hip osteoarthritis**

In most studies, cam morphology has been associated with hip OA. The strength of association in several cross sectional and retrospective studies varied between odds ratio (OR)s of 2.2 (95% CI 1.7, 2.8) and 20.6 (95% CI 3.4, 34.8).<sup>12,18,25</sup> The number of well-designed epidemiological studies assessing the relationship between cam morphology and hip OA are limited. Three prospective cohort studies and 2 nested case control studies that included people without hip OA at baseline demonstrated an association between cam morphology and development of hip OA later in life (**TABLE 3**).<sup>2,49,53,62,67</sup> The strength of association varies between ORs of 2.1 (95% CI 1.6, 2.9) and 9.7 (95% CI 4.7, 19.8), primarily depending on the alpha angle threshold used for diagnosis. The positive predictive value (PPV) for developing end-stage OA within 5 years when having cam morphology was 10.9% for an alpha angle greater than 60° and 25.0% for an alpha angle greater than 83°. <sup>2</sup>

### **Relationship between pincer morphology and osteoarthritis**

Pincer morphology does not appear to play a role in the development of hip OA. Three prospective cohort studies defined the presence of pincer morphology by a center-edge angle (CEA) of greater than 33.7° or 40°. <sup>4,62,67</sup> In the CHECK cohort<sup>4</sup>, pincer morphology was measured both laterally (on anteroposterior (AP) pelvic radiographs) and anteriorly (on faux



profile lateral radiographs). Neither anterior pincer morphology nor lateral pincer morphology was associated with development of hip OA within 5 years. Surprisingly, when pincer morphology was present both anteriorly and laterally, a significant protective effect for development of end-stage OA was found (OR 0.34, 95% CI 0.13, 0.87). This is consistent with the data from Chingford cohort<sup>67</sup> that did not identify an association between higher LCEA angles (only measured on AP radiographs) and development of hip OA. In this cohort, the continuous measure of the LCEA was divided into tertiles. Having an LCEA in the highest tertile (greater than 33.7°) was neither associated with development of radiographic hip OA, defined as a Kellgren & Lawrence<sup>34</sup> (K&L) grade greater than 2 ( $P=.64$ ) nor with the need for total hip replacement ( $P=.67$ ) 19 years later. Finally, results from the Rotterdam study<sup>62</sup> also failed to show an increased risk of developing hip OA at 10-year follow-up, with an OR of 1.24 (95% CI 0.93, 1.66) for pincer morphology.

## **Discussion**

Cam and pincer morphology are common findings in the general population but the prevalence rate vary greatly among studied subpopulations. Cam morphology is associated with future development of hip OA while a link between pincer morphology and OA has never been identified in epidemiological studies. It is important to recognise that all studies on the prevalence and its association with OA investigated morphology only, which does not equate to FAI syndrome, which also includes the presence of symptoms and clinical findings.<sup>26</sup>

### *Differences and limitations in quantifying cam morphology*

There is a large variation in the reported prevalence of cam and pincer morphology between subgroups, with some of that variation attributed to the variability in methodology used to determine the presence of cam and pincer morphology. In the literature, while the alpha angle

is an accepted measure to define cam morphology<sup>54</sup>, the angular thresholds that is used varies from 50 to 83°. <sup>5,23,54</sup> Furthermore, alpha angles can be measured by different imaging techniques, including radiographs, computed tomography (CT), and magnetic resonance imaging (MRI). Generally, using radial imaging with multiple measurement points (CT and MRI) around the femoral neck is more likely to detect the presence of cam morphology than 2-dimensional imaging (radiographs) and thus result in higher prevalence.<sup>19</sup> However, the use of multiple measurement points might increase the false positive rate.

#### *Differences in cam morphology prevalence in subgroups*

The differences in the prevalence of cam morphology between subgroups might provide some clues on aetiology. The greatest differences in prevalence are observed between athletes and non-athletes. The high prevalence of cam morphology observed in athletes might be due to repetitive axial loading, especially during skeletal maturation.<sup>3,55,61,64</sup> This might also partly explain the lower prevalence in females, as they mature earlier than males and probably have less exposure to repetitive axial loading during the second growth spurt, when cam morphology usually develops in males. Cam morphology is probably less frequent in the East Asian population, even in those with an athletic background. However, evidence is conflicting and no direct relationship between genetics and cam morphology has been established yet. Finally, whether the isolated presence of cam morphology is associated with, or predictive for symptoms and/or hip pain is unknown. But, we must be cautious in these interpretations, because although subgroups with a higher prevalence of cam morphology have been identified, it should be emphasized that most of these studies suffer from a high risk of bias.<sup>16</sup>

#### *Differences and limitations in quantifying pincer morphology*

The prevalence of pincer morphology is also highly dependent on how it is quantified and the imaging technique used.<sup>4</sup> Pincer morphology can be further defined as having focal or global (acetabular) overcoverage. Focal overcoverage has been defined by several indirect measures such as: the crossover sign, posterior wall sign, and ischial spine sign, which all have generally poor reliability and validity to define true retroversion / pincer morphology.<sup>69</sup> Global overcoverage can be defined by the presence of coxa profunda or protrusio acetabuli or the CEA.<sup>10,50</sup> Coxa profunda and protrusio acetabuli do not seem to be associated with the presence of pincer morphology.<sup>50</sup> Therefore, due to this heterogeneity in definition it is difficult to compare prevalence studies for pincer morphology.

#### *Pincer morphology and hip OA*

The prospective studies on the association between pincer morphology and hip OA all used the LCEA on AP radiographs and are therefore comparable.<sup>4,49,53,62,67</sup> However, none of these epidemiological studies could identify an association between pincer morphology and development of OA. It is also notable that 2 systematic reviews found a higher prevalence of pincer morphology in asymptomatic individuals than in symptomatic patients.<sup>20,41</sup> The reader should also bear in mind that although discussed separately, cam and pincer morphology are frequently found together, also known as a mixed type morphology.<sup>42</sup>

#### *Cam morphology and hip OA*

Despite the reported association between cam morphology and development of hip OA, one should keep in mind that the majority of people with cam morphology will not develop hip OA. Of the hips with cam morphology, between 6% and 25% will develop future OA within 5 to 19 years.<sup>2,53</sup> For cross-sectional and retrospective studies, an important confounder is the fact that the radiographic appearance of OA might mimic cam morphology. For example the

presence of osteophytes on the femoral head and/or flattening of the femoral head may be related to the OA process. This is hard to distinguish when OA and cam morphology are assessed on the same radiographs. This is less of an issue in a few well-designed prospective studies summarized in **TABLE 3**, but these studies have other methodological limitations such as the imaging modalities used and age of the participants.<sup>2,4,49,53,62,67</sup> All of these studies used AP pelvic radiographs and although this is the gold standard to quantify hip OA, it is suboptimal to define the presence of cam morphology. Only the more laterally located cams are seen on AP radiographs and the prevalence is therefore underestimated. The influence of this underestimation on the true association with hip OA is unknown. Further, the studies summarized only included middle aged to older people. The youngest participants included in the CHECK<sup>2</sup> and Chingford<sup>67</sup> cohorts were 45 years of age and the mean age was 55 and 54 years, respectively. The oldest people were included in the Rotterdam study<sup>62</sup>, (minimum age 55 years, mean age 64 years) and in the Johnston county OA cohort study<sup>49</sup> (mean age 62 years). As cam morphology develops during skeletal growth in most cases, it is already present during early adulthood. Therefore, the relationship between cam morphology and hip degeneration between early adulthood and the age of 45 years is unknown. Some indications suggest that this relationship might be stronger in younger people than in middle aged to older people. First, the Rotterdam study showed a stronger relation between cam morphology and OA in people 65 years of age or younger (OR 3.1, 95% CI 2.1, 4.6) while the association disappeared in people greater than 65 years (OR 1.4, 95% CI 0.9, 2.2).<sup>62</sup> Second, features known to be associated with hip OA have been identified in younger populations<sup>11,43,58</sup> with the severity of cam morphology associated with the presence of labral tears and chondral defects.<sup>58</sup> A cross sectional study of asymptomatic participants with a mean age of 20 years showed a decreased in cartilage thickness in those with cam morphology.<sup>60</sup> Finally, from intra-

operative findings it is known that severe cartilage damage can already exist in young people with cam morphology.<sup>13,14</sup> However, well-designed studies in young adults are lacking.

### *Future studies*

Based on the results of this overview, there is a need for standardising criteria to determine the presence of cam and pincer morphology. For cam morphology the alpha angle is most often used and despite its limitations it is probably the best measure to date and future studies should therefore at least report the alpha angle. An alpha angle threshold of 60° has been proposed for AP radiographs<sup>6</sup>, but there is no validated threshold for other radiographic views. To aid future comparison between studies it might be helpful to present results for different alpha angle threshold values. Many people with cam or pincer morphology will not develop any symptoms from this bony variant. Future studies should therefore also focus on characteristics that can differentiate persons with cam and pincer morphology that will become symptomatic and/or develop hip OA. Characteristics that may be worth considering include hip muscle strength, hip range of motion, gait pattern characteristics, the size of cam morphology, and type and amount of physical activities performed. This might lead to the identification of modifiable risk factors to prevent, stop, or slow down disease progression and also help avoid overtreatment. Future studies should also monitor whether treatment for FAI syndrome, nonsurgical or surgical, can stop or slow down the progression towards hip OA.

### **Conclusion**

Cam and pincer morphology is highly prevalent in the general population. Cam morphology is linked to hip OA in the middle-aged population, but no data are available on its relationship among younger people. The association between pincer morphology and hip OA has not been demonstrated in the available prospective cohort studies. The presence of cam and/or pincer

morphology does not always lead to FAI syndrome and subsequent hip OA and future research should focus on identifying factors that may predict who becomes symptomatic (FAI syndrome) in the presence of cam and/or pincer morphology and who subsequently will progress to have hip OA later in life.

## References

1. Agricola R, Bessems JH, Ginai AZ, et al. The development of Cam-type deformity in adolescent and young male soccer players. *Am J Sports Med.* 2012;40(5):1099-1106.
2. Agricola R, Heijboer MP, Bierma-Zeinstra SM, Verhaar JA, Weinans H, Waarsing JH. Cam impingement causes osteoarthritis of the hip: a nationwide prospective cohort study (CHECK). *Ann Rheum Dis.* 2013;72(6):918-923.
3. Agricola R, Heijboer MP, Ginai AZ, et al. A cam deformity is gradually acquired during skeletal maturation in adolescent and young male soccer players: a prospective study with minimum 2-year follow-up. *Am J Sports Med.* 2014;42(4):798-806.
4. Agricola R, Heijboer MP, Roze RH, et al. Pincer deformity does not lead to osteoarthritis of the hip whereas acetabular dysplasia does: acetabular coverage and development of osteoarthritis in a nationwide prospective cohort study (CHECK). *Osteoarthritis Cartilage.* 2013;21(10):1514-1521.
5. Agricola R, Waarsing JH, Arden NK, et al. Cam impingement of the hip: a risk factor for hip osteoarthritis. *Nat Rev Rheumatol.* 2013;9(10):630-634.
6. Agricola R, Waarsing JH, Thomas GE, et al. Cam impingement: defining the presence of a cam deformity by the alpha angle: data from the CHECK cohort and Chingford cohort. *Osteoarthritis Cartilage.* 2014;22(2):218-225.
7. Ahn T, Kim CH, Kim TH, et al. What is the Prevalence of Radiographic Hip Findings Associated With Femoroacetabular Impingement in Asymptomatic Asian Volunteers? *Clin Orthop Relat Res.* 2016;474(12):2655-2661.
8. Allen D, Beaulé PE, Ramadan O, Doucette S. Prevalence of associated deformities and hip pain in patients with cam-type femoroacetabular impingement. *J Bone Joint Surg Br.* 2009;91(5):589-594.
9. Anderson LA, Anderson MB, Kapron A, et al. The 2015 Frank Stinchfield Award: Radiographic Abnormalities Common in Senior Athletes With Well-functioning Hips but Not Associated With Osteoarthritis. *Clin Orthop Relat Res.* 2016;474(2):342-352.
10. Anderson LA, Kapron AL, Aoki SK, Peters CL. Coxa profunda: is the deep acetabulum overcovered? *Clin Orthop Relat Res.* 2012;470(12):3375-3382.
11. Ayeni OR, Banga K, Bhandari M, et al. Femoroacetabular impingement in elite ice hockey players. *Knee Surg Sports Traumatol Arthrosc.* 2014;22(4):920-925.
12. Bardakos NV, Villar RN. Predictors of progression of osteoarthritis in femoroacetabular impingement: a radiological study with a minimum of ten years follow-up. *J Bone Joint Surg Br.* 2009;91(2):162-169.
13. Beck M, Kalhor M, Leunig M, Ganz R. Hip morphology influences the pattern of damage to the acetabular cartilage: femoroacetabular impingement as a cause of early osteoarthritis of the hip. *J Bone Joint Surg Br.* 2005;87(7):1012-1018.
14. Clohisy JC, Baca G, Beaulé PE, et al. Descriptive epidemiology of femoroacetabular impingement: a North American cohort of patients undergoing surgery. *Am J Sports Med.* 2013;41(6):1348-1356.
15. de Bruin F, Reijnen M, Farhang-Razi V, Bloem JL. Radiographic signs associated with femoroacetabular impingement occur with high prevalence at all ages in a hospital population. *Eur Radiol.* 2013;23(11):3131-3139.

16. Dickenson E, Wall PD, Robinson B, et al. Prevalence of cam hip shape morphology: a systematic review. *Osteoarthritis Cartilage*. 2016;24(6):949-961.
17. Diesel CV, Ribeiro TA, Coussirat C, Scheidt RB, Macedo CA, Galia CR. Coxa profunda in the diagnosis of pincer-type femoroacetabular impingement and its prevalence in asymptomatic subjects. *Bone Joint J*. 2015;97-B(4):478-483.
18. Doherty M, Courtney P, Doherty S, et al. Nonspherical femoral head shape (pistol grip deformity), neck shaft angle, and risk of hip osteoarthritis: a case-control study. *Arthritis Rheum*. 2008;58(10):3172-3182.
19. Dudda M, Albers C, Mamisch TC, Werlen S, Beck M. Do normal radiographs exclude asphericity of the femoral head-neck junction? *Clin Orthop Relat Res*. 2009;467(3):651-659.
20. Frank JM, Harris JD, Erickson BJ, et al. Prevalence of Femoroacetabular Impingement Imaging Findings in Asymptomatic Volunteers: A Systematic Review. *Arthroscopy*. 2015;31(6):1199-1204.
21. Fukushima K, Uchiyama K, Takahira N, et al. Prevalence of radiographic findings of femoroacetabular impingement in the Japanese population. *J Orthop Surg Res*. 2014;9:25.
22. Gerhardt MB, Romero AA, Silvers HJ, Harris DJ, Watanabe D, Mandelbaum BR. The prevalence of radiographic hip abnormalities in elite soccer players. *Am J Sports Med*. 2012;40(3):584-588.
23. Gosvig KK, Jacobsen S, Palm H, Sonne-Holm S, Magnusson E. A new radiological index for assessing asphericity of the femoral head in cam impingement. *J Bone Joint Surg Br*. 2007;89(10):1309-1316.
24. Gosvig KK, Jacobsen S, Sonne-Holm S, Gebuhr P. The prevalence of cam-type deformity of the hip joint: a survey of 4151 subjects of the Copenhagen Osteoarthritis Study. *Acta Radiol*. 2008;49(4):436-441.
25. Gosvig KK, Jacobsen S, Sonne-Holm S, Palm H, Troelsen A. Prevalence of malformations of the hip joint and their relationship to sex, groin pain, and risk of osteoarthritis: a population-based survey. *J Bone Joint Surg Am*. 2010;92(5):1162-1169.
26. Griffin DR, Dickenson EJ, O'Donnell J, et al. The Warwick Agreement on femoroacetabular impingement syndrome (FAI syndrome): an international consensus statement. *Br J Sports Med*. 2016;50(19):1169-1176.
27. Hack K, Di Primio G, Rakhra K, Beaulé PE. Prevalence of cam-type femoroacetabular impingement morphology in asymptomatic volunteers. *J Bone Joint Surg Am*. 2010;92(14):2436-2444.
28. Harris JD, Gerrie BJ, Varner KE, Lintner DM, McCulloch PC. Radiographic Prevalence of Dysplasia, Cam, and Pincer Deformities in Elite Ballet. *Am J Sports Med*. 2016;44(1):20-27.
29. Impellizzeri FM, Mannion AF, Naal FD, Hersche O, Leunig M. The early outcome of surgical treatment for femoroacetabular impingement: success depends on how you measure it. *Osteoarthritis Cartilage*. 2012;20(7):638-645.
30. Jung KA, Restrepo C, Hellman M, AbdelSalam H, Morrison W, Parvizi J. The prevalence of cam-type femoroacetabular deformity in asymptomatic adults. *J Bone Joint Surg Br*. 2011;93(10):1303-1307.
31. Kang AC, Gooding AJ, Coates MH, Goh TD, Armour P, Rietveld J. Computed tomography assessment of hip joints in asymptomatic individuals in relation to femoroacetabular impingement. *Am J Sports Med*. 2010;38(6):1160-1165.
32. Kapron AL, Anderson AE, Aoki SK, et al. Radiographic prevalence of femoroacetabular impingement in collegiate football players: AAOS Exhibit Selection. *J Bone Joint Surg Am*. 2011;93(19):e111(111-110).
33. Kapron AL, Peters CL, Aoki SK, et al. The prevalence of radiographic findings of structural hip deformities in female collegiate athletes. *Am J Sports Med*. 2015;43(6):1324-1330.
34. Kellgren JH, Lawrence JS. Radiological assessment of osteo-arthritis. *Ann Rheum Dis*. 1957;16(4):494-502.
35. Khanna V, Caragianis A, Diprimio G, Rakhra K, Beaulé PE. Incidence of hip pain in a prospective cohort of asymptomatic volunteers: is the cam deformity a risk factor for hip pain? *Am J Sports Med*. 2014;42(4):793-797.

36. Laborie LB, Lehmann TG, Engesaeter IO, Eastwood DM, Engesaeter LB, Rosendahl K. Prevalence of radiographic findings thought to be associated with femoroacetabular impingement in a population-based cohort of 2081 healthy young adults. *Radiology*. 2011;260(2):494-502.
37. Larson CM, Sikka RS, Sardelli MC, et al. Increasing alpha angle is predictive of athletic-related "hip" and "groin" pain in collegiate National Football League prospects. *Arthroscopy*. 2013;29(3):405-410.
38. Lerebours F, Robertson W, Neri B, Schulz B, Youm T, Limpisvasti O. Prevalence of Cam-Type Morphology in Elite Ice Hockey Players. *Am J Sports Med*. 2016;44(4):1024-1030.
39. Leunig M, Juni P, Werlen S, et al. Prevalence of cam and pincer-type deformities on hip MRI in an asymptomatic young Swiss female population: a cross-sectional study. *Osteoarthritis Cartilage*. 2013;21(4):544-550.
40. Li Y, Helvie P, Mead M, Gagnier J, Hammer MR, Jong N. Prevalence of Femoroacetabular Impingement Morphology in Asymptomatic Adolescents. *J Pediatr Orthop*. 2017;37(2):121-126.
41. Mascarenhas VV, Rego P, Dantas P, et al. Imaging prevalence of femoroacetabular impingement in symptomatic patients, athletes, and asymptomatic individuals: A systematic review. *Eur J Radiol*. 2016;85(1):73-95.
42. Matsuda DK, Gupta N, Khatod M, et al. Poorer Arthroscopic Outcomes of Mild Dysplasia With Cam Femoroacetabular Impingement Versus Mixed Femoroacetabular Impingement in Absence of Capsular Repair. *Am J Orthop (Belle Mead NJ)*. 2017;46(1):E47-E53.
43. Mayes S, Ferris AR, Smith P, Garnham A, Cook J. Atraumatic tears of the ligamentum teres are more frequent in professional ballet dancers than a sporting population. *Skeletal Radiol*. 2016;45(7):959-967.
44. Mineta K, Goto T, Wada K, et al. CT-based morphological assessment of the hip joint in Japanese patients: association with radiographic predictors of femoroacetabular impingement. *Bone Joint J*. 2016;98-B(9):1167-1174.
45. Monazzam S, Bomar JD, Dwek JR, Hosalkar HS, Pennock AT. Development and prevalence of femoroacetabular impingement-associated morphology in a paediatric and adolescent population: a CT study of 225 patients. *Bone Joint J*. 2013;95-B(5):598-604.
46. Mori R, Yasunaga Y, Yamasaki T, et al. Are cam and pincer deformities as common as dysplasia in Japanese patients with hip pain? *Bone Joint J*. 2014;96-B(2):172-176.
47. Mosler AB, Crossley KM, Waarsing JH, et al. Ethnic Differences in Bony Hip Morphology in a Cohort of 445 Professional Male Soccer Players. *Am J Sports Med*. 2016;44(11):2967-2974.
48. Nardo L, Parimi N, Liu F, et al. Femoroacetabular Impingement: Prevalent and Often Asymptomatic in Older Men: The Osteoporotic Fractures in Men Study. *Clin Orthop Relat Res*. 2015;473(8):2578-2586.
49. Nelson AE, Stiller JL, Shi XA, et al. Measures of hip morphology are related to development of worsening radiographic hip osteoarthritis over 6 to 13 year follow-up: the Johnston County Osteoarthritis Project. *Osteoarthritis Cartilage*. 2016;24(3):443-450.
50. Nepple JJ, Lehmann CL, Ross JR, Schoenecker PL, Clohisy JC. Coxa profunda is not a useful radiographic parameter for diagnosing pincer-type femoroacetabular impingement. *J Bone Joint Surg Am*. 2013;95(5):417-423.
51. Nepple JJ, Riggs CN, Ross JR, Clohisy JC. Clinical presentation and disease characteristics of femoroacetabular impingement are sex-dependent. *J Bone Joint Surg Am*. 2014;96(20):1683-1689.
52. Nepple JJ, Vigdorchik JM, Clohisy JC. What Is the Association Between Sports Participation and the Development of Proximal Femoral Cam Deformity? A Systematic Review and Meta-analysis. *Am J Sports Med*. 2015;43(11):2833-2840.
53. Nicholls AS, Kiran A, Pollard TC, et al. The association between hip morphology parameters and nineteen-year risk of end-stage osteoarthritis of the hip: a nested case-control study. *Arthritis Rheum*. 2011;63(11):3392-3400.
54. Notzli HP, Wyss TF, Stoecklin CH, Schmid MR, Treiber K, Hodler J. The contour of the femoral head-neck junction as a predictor for the risk of anterior impingement. *J Bone Joint Surg Br*. 2002;84(4):556-560.



55. Palmer A, Fernquest S, Gimpel M, et al. Physical activity during adolescence and the development of cam morphology: a cross-sectional cohort study of 210 individuals. *Br J Sports Med.* 2018;52(9):601-610.
56. Philippon MJ, Ho CP, Briggs KK, Stull J, LaPrade RF. Prevalence of increased alpha angles as a measure of cam-type femoroacetabular impingement in youth ice hockey players. *Am J Sports Med.* 2013;41(6):1357-1362.
57. Pollard TC, Villar RN, Norton MR, et al. Femoroacetabular impingement and classification of the cam deformity: the reference interval in normal hips. *Acta Orthop.* 2010;81(1):134-141.
58. Register B, Pennock AT, Ho CP, Strickland CD, Lawand A, Philippon MJ. Prevalence of abnormal hip findings in asymptomatic participants: a prospective, blinded study. *Am J Sports Med.* 2012;40(12):2720-2724.
59. Reichenbach S, Juni P, Werlen S, et al. Prevalence of cam-type deformity on hip magnetic resonance imaging in young males: a cross-sectional study. *Arthritis Care Res (Hoboken).* 2010;62(9):1319-1327.
60. Reichenbach S, Leunig M, Werlen S, et al. Association between cam-type deformities and magnetic resonance imaging-detected structural hip damage: a cross-sectional study in young men. *Arthritis Rheum.* 2011;63(12):4023-4030.
61. Roels P, Agricola R, Oei EH, Weinans H, Campoli G, Zadpoor AA. Mechanical factors explain development of cam-type deformity. *Osteoarthritis Cartilage.* 2014;22(12):2074-2082.
62. Saberi Hosnijeh F, Zuiderwijk ME, Versteeg M, et al. Cam Deformity and Acetabular Dysplasia as Risk Factors for Hip Osteoarthritis. *Arthritis Rheumatol.* 2017;69(1):86-93.
63. Siebenrock KA, Ferner F, Noble PC, Santore RF, Werlen S, Mamisch TC. The cam-type deformity of the proximal femur arises in childhood in response to vigorous sporting activity. *Clin Orthop Relat Res.* 2011;469(11):3229-3240.
64. Siebenrock KA, Kaschka I, Frauchiger L, Werlen S, Schwab JM. Prevalence of cam-type deformity and hip pain in elite ice hockey players before and after the end of growth. *Am J Sports Med.* 2013;41(10):2308-2313.
65. Tak I, Weir A, Langhout R, et al. The relationship between the frequency of football practice during skeletal growth and the presence of a cam deformity in adult elite football players. *Br J Sports Med.* 2015;49(9):630-634.
66. Tannenbaum E, Kopydlowski N, Smith M, Bedi A, Sekiya JK. Gender and racial differences in focal and global acetabular version. *J Arthroplasty.* 2014;29(2):373-376.
67. Thomas GE, Palmer AJ, Batra RN, et al. Subclinical deformities of the hip are significant predictors of radiographic osteoarthritis and joint replacement in women. A 20 year longitudinal cohort study. *Osteoarthritis Cartilage.* 2014;22(10):1504-1510.
68. Van Houcke J, Yau WP, Yan CH, et al. Prevalence of radiographic parameters predisposing to femoroacetabular impingement in young asymptomatic Chinese and white subjects. *J Bone Joint Surg Am.* 2015;97(4):310-317.
69. Zaltz I, Kelly BT, Hetsroni I, Bedi A. The crossover sign overestimates acetabular retroversion. *Clin Orthop Relat Res.* 2013;471(8):2463-2470.

**TABLE 1:** Prevalence of cam morphology in asymptomatic individuals

Study	Group	Definition of cam morphology	No. of individuals (hips)	Mean (range or $\pm$ SD) age in years	Sex (Male/Female, %)	Imaging modality	Prevalence (Male/Female, %)
<b>Agricola et al.<sup>1</sup></b>	Athletes: soccer	AA > 60° and/or flattening or prominence	89 (178) cases, 92 (184) controls	14.8 (12-19) 13.8 (12-19)	All males	AP and FLL radiography	AA: 26, VS: 66 (cases) AA: 17, VS: 18 (controls) (PH)
<b>Agricola et al.<sup>3</sup> (FU 2y)</b>	Athletes: soccer	AA > 60° and/or flattening or prominence	63 (126)	14.43 (12-19)	All males	AP and FLL radiography	AA: 38.9 VS: 69.0 (PH)
<b>Anderson et al.<sup>9</sup></b>	Senior athletes	na	547 (1081)	67 $\pm$ 8	55/45	AP and FLL radiography	66.7 (PH)
<b>Hack et al.<sup>27</sup></b>	Volunteers	AA > 50.5°	200 (400)	29.4 (21.4-50.6)	44/56	MRI	24.7/5.4 (PP)
<b>Jung et al.<sup>30</sup></b>	Abdominal, pelvic or other medial issue	AA > 68° (men) AA > 50° (women)	380 (755)	60.4 (25-92)	28/72	Abdominal or pelvic AP scout CT	28.8/11.7 (PH)
<b>Kang et al.<sup>31</sup></b>	Abdominal trauma or nonspecific abdominal pain	AA > 55°	50 (100)	na (15-40)	54/46	Abdominal CT	10.0 (PH)
<b>Kapron et al.<sup>32</sup></b>	Athletes: collegiate football	AA > 50° and/or HNO < 8mm	67 (134)	21 $\pm$ 1.9	All male	AP and FLL radiography	AA: 72 HNO: 64 (PH)
<b>Kapron et al.<sup>33</sup></b>	Athletes: collegiate	AA > 50° and/or HNO < 8mm	63 (126)	19.6 $\pm$ 1.4	All female	AP and FLL radiography	48 (PH) 60 (PP)

	volleyball, soccer, T&F							
<b>Khanna et al.<sup>35</sup> (FU 4.4y)</b>	Volunteers	AA > 50.5° and second analysis with AA > 60°	BL: (400) FU: 170 (340)	200	FU: 29.5 (25.7-54.5)	45.3/54.7	MRI	FU: 25.9 (PH)
<b>Laborie et al.<sup>36</sup></b>	Follow-up of initially newborns	Pistol-grip deformity, flattening and prominence		2.060 (4120)	18.6 (17.2-20.1)	42.1/57.9	AP and FLL radiography	35.0/10.2 (PP)
<b>Larson et al.<sup>37</sup></b>	Athletes: collegiate football	AA > 55°	125 (239)	na		All male	AP and FLL radiography	65.3 (PH) 75.2 (PP)
<b>Lerebours et al.<sup>38</sup></b>	Athletes: ice hockey	AA ≥ 55°	130 (260)		24.4 ± 4.3	na	AP and FLL radiography	69.4 (PH)
<b>Leunig et al.<sup>39</sup></b>	Females from vocational/gymnasium school, males Swiss army	AA > 50.5°	324 (324)		20.0 ± 0.9 (male) 19.3 ± 1.3 (female)	75.3/24.7	MRI	24.0/0.0 (PP)
<b>Li et al.<sup>40</sup></b>	Children with disorder unrelated to hip	AA ≥ 55°	558 (1116)		14.4 (10-18.2)	49.5/50.5	Pelvic CT	23.9/9.9 (PP)
<b>Mineta et al.<sup>44</sup></b>	Disorder unrelated to hip (Japanese)	AA > 55° and/or FHNO ratio < 0.15	1178 (1178)		58.2 (20-89, ± 14.8)	59/41	Abdominal and pelvic CT	54.4/32.3 (PH)
<b>Mosler et al.<sup>47</sup></b>	Athletes: soccer	AA > 60°	445 (890)		25 ± 4.9	All male	AP pelvic and Dunn view radiography	72 (PP)

<b>Philippon et al.</b> <sup>56</sup>	Athletes: ice hockey	AA $\geq 55^\circ$	61 (na) cases 27 (na) controls	14.5 (10-18, $\pm 2.7$ ) 15.2 (10-18, $\pm 2.7$ )	All male	MRI	75 42 (PP)
<b>Pollard et al.</b> <sup>57</sup>	General population	AA $> 62^\circ$ and AOR $< 0.14$	83 (166)	47.5 (25-69) male 44.4 (22-67) female	47/53	Cross-table lateral radiography	13.0/7.0 (PP)
<b>Reichenbach et al.</b> <sup>59</sup>	Swiss army recruiters	2 = cam, AHNO $<10$ mm, 3 = severe cam, AHNO $>10$ mm	244 (244)	19.9 (18-24)	All male	MRI	24.0 (PP)
<b>Van Houcke et al.</b> <sup>68</sup>	Chinese and Belgian	AA $> 55^\circ$	Chinese: 102 (204) Belgian: 99 (198)	na (18-40)	52.2/47.8	CT	31/17 41/39 (PH)

Abbreviations: AA, alpha angle; AHNO, anterior head-neck offset; AOR, anterior offset ratio; AP, anteroposterior; BL, baseline; CCD, caput-collum-diaphyseal; FHNO, femoral head-neck offset; FLL, frog-leg lateral; FU, follow-up; HNO, head-neck offset; na, not available; PH, per hip; PP, per person; T&F, track & field; VS, visual scoring; y, year

\* If prevalence per gender is not specified, the overall prevalence is presented.

**TABLE 2:** Prevalence of pincer morphology in asymptomatic individuals

Study	Group	Definition of pincer morphology	No. of individuals (hips)	Mean (range or $\pm$ SD) age in years	Sex (Male/Female, %)	Imaging modality	Prevalence (Male/Female, %)
<b>Ahn et al.<sup>7</sup></b>	Korean volunteers	COS, PWS or LCEA $>40^\circ$	200 (400)	34.7 (21-49)	36.5/63.5	AP, Sugioka and $45^\circ$ Dunn radiography	27/21 (PP)
<b>De Bruin et al.<sup>15</sup></b>	Pelvic radiography patients	CEA $>39^\circ$ , AI $<0^\circ$ , CP, PA, AR	262 (522)	na	38/62	AP radiography	63.2 (PH)
<b>Diesel et al.<sup>17</sup></b>	Volunteers	LCEA $>40^\circ$ AI $<0^\circ$ COS CP	226 (452)	36.5 (28-50)	46.3/53.7	AP radiography	10.9/10.9 30.3/41.2 10.9/16.7 60.5/92 (PH)
<b>Gerhardt et al.<sup>22</sup></b>	Athletes: elite soccer	COS	95 (190)	25.4 ( $\pm$ 4.2)	79/21	AP pelvis and FLL radiography	26.7/10 (PP)
<b>Harris et al.<sup>28</sup></b>	Athletes: elite ballet	PWS, COS, ISS, LCEA $>40^\circ$ , CP, PA	47 (94)	23.8 ( $\pm$ 5.4)	45/55	AP pelvis, false profile and Dunn $45^\circ$ radiography	74 (PP)
<b>Kang et al.<sup>31</sup></b>	Abdominal trauma or nonspecific abdominal pain	AV $<15^\circ$ or COS AO/CP (CEA $>40^\circ$ )	50 (100)	na (15-40)	46/54	Abdominal CT	13/1 20 9/7 (PH)
<b>Kapron et al.<sup>32</sup></b>	Athletes: collegiate football	LCEA $>40^\circ$ , AI $<0^\circ$ and/or COS	67 (134)	21 ( $\pm$ 1.9)	All males	AP pelvis and FLL radiography	52 (1 sign) 10 (2 signs) 4 (3 signs) (PH)

<b>Kapron et al.<sup>33</sup></b>	Athletes: collegiate volleyball, soccer, T&F		LCEA >40°  LCEA >40° and AI <0°	63 (126)	19.6 (± 1.4)	All females	AP pelvis and FLL radiography	1 (PH), 2 (PP)  1 (PH), 2 (PP)
<b>Laborie et al.<sup>36</sup></b>	Follow-up initially newborns	of	1 or more finding COS PWS AO	2060 (4120)	18.6 (17.2-20.1)	42.1/57.9	AP and FLL radiography	34.3/16.6 51.4/45.5 23.4/11 14.6/4.9 (PP)
<b>Lerebours et al.<sup>38</sup></b>	Athletes: ice hockey		COS	130 (260)	24.4 ± 4.3	na	AP and FLL radiography	59.8 (PP)
<b>Leunig et al.<sup>39</sup></b>	Females from vocational/grammar school, males Swiss army		AD ≤ 3mm	324 (324)	20.0 ± 0.9 (male) 19.3 ± 1.3 (female)	75.3/24.7	MRI	6/10 (PP)
<b>Li et al.<sup>40</sup></b>	Children with disorder unrelated to hip		LCEA >40°	558 (1116)	14.4 (10-18.2)	49.5/50.5	Pelvic CT	29.7/35.1 (PP)
<b>Mineta et al.<sup>44</sup></b>	Japanese population, reason unrelated to hip		LCEA >40°, AI <0°, COS	1178 (1178)	58.2 (20-89)	59/41	Pelvic CT	41.7/31.3 (PH)
<b>Monazzam et al.<sup>45</sup></b>	Abdominal problems		LCEA ≥ 40°, TA ≤ 0°, AR (AV ≤ 0° and LCEA ≥ 40°)	225 (450)	10.4 (2-19)	45.8/54.2	Pelvic CT	5.8/2.0 4.4/5.3 6.8/4.1 (PH)
<b>Mosler et al.<sup>47</sup></b>	Athletes: elite soccer		LCEA >40°	445 (890)	25 (± 4.9)	All males	AP and Dunn radiography	3.0 (PP)

Abbreviations: AD, acetabular depth; AI, acetabular index; AO, acetabular overcoverage; AR, acetabular retroversion; AV, acetabular version; CEA, center-edge angle; COS, cross-over sign; CP, coxa profunda; FLL, frog-leg lateral; ISS, ischial spine sign; LCEA, lateral

center-edge angle; na, not available; PA, protrusion acetabulae; PH, per hip; PP, per person;  
PWS, posterior wall sign; TA, Tönnis angle

\* If prevalence per gender is not specified, the overall prevalence is presented.

**TABLE 3:** Characteristics of multiple longitudinal studies on relationship between cam/pincer morphology and osteoarthritis, all based on AP radiographs

Study (FU, y)	No. of individuals (hips)	Mean (range or $\pm$ SD) age in years	Sex (Male/Female, %)	Definition cam* and pincer <sup>†</sup> morphology	Cam morphology prevalence (%)	Pincer morphology prevalence (%)	Definition of osteoarthritis	Odds ratio hip osteoarthritis [95% CI]
<b>Agricola et al.<sup>2</sup> (FU 5y)</b>	723 (1411)	55.9 (45-65, $\pm$ 5.2)	20/80	*AA >60° AA >83° AA >83° and IR $\leq$ 20°	11.1	na	End-stage OA: K&L $\geq$ 3 or THR	3.67 [1.68-8.01] 9.66 [4.72-19.78] 25.21 [7.89-80.58]
<b>Agricola et al.<sup>4</sup> (FU 5y)</b>	720 (1391)	55.9 (45-65, $\pm$ 5.2)	21/79	<sup>†</sup> LCEA >40° or ACEA >40°	na	54.6	End-stage OA: K&L $\geq$ 3 or THR	0.34 [0.13-0.87]
<b>Nelson et al.<sup>49</sup> (FU 6y, 12.7y)</b>	120 (239, CA: 71, CO: 168)	63 $\pm$ 8 (CA) 62 $\pm$ 9 (CO)	25/75	*AA >60°    <sup>†</sup> LCEA >40°	Male: 59 (CA) 40 (CO) Female: 47 (CA) 18 (CO)	Male: 10 (CA) 6 (CO) Female: 24 (CA) 17 (CO)	OA: K&L $\geq$ 3 or THR	3.57 [1.17-10.90] in male 4.61 [2.09-10.16] in female NS in male NS in female
<b>Nicholls et al.<sup>53</sup> (FU 19y)</b>	135 (268, CA: 25, CO: 243)	55 (50-60)	All female	*AA  <sup>†</sup> LCEA	na	na	End-stage OA: THR	1.052 per 1° increase NS
<b>Saberi et al.<sup>62</sup> (FU 9.2y)</b>	4438 (2960 RS-I, 1478 RS-II)	65.1 $\pm$ 6.4 (RS-I) 62.9 $\pm$ 6.4 (RS-II)	43/57 (RS-I) 44/56 (RS-II)	*AA >60°  <sup>†</sup> CEA >40°	RS-I: 8.3 (L), 6.4 (R) RS-II: 7.2 (L), 7 (R)	RS-I: 10.9 (L), 8.9 (R) RS-II: 13.5 (L), 8.6 (R)	Incident OA: K&L $\geq$ 2 or THR	2.11 [1.55-2.87] NS
<b>Thomas et al.<sup>67</sup> (FU 19y)</b>	340 (634): OA group 734 (1466): THR group	54.2 (44-67)	All female	*AA >65°  <sup>†</sup> LCEA >33.7°	na	na	OA: K&L $\geq$ 2 End-stage OA: THR	1.05 [1.01-1.09] for OA 1.04 [1.00-1.08] for THR NS for OA NS for THR



Abbreviations: AA, alpha angle; ACEA, anterior center-edge angle; CA, cases, center-edge angle; CI, confidence interval; CO, controls; FU, follow-up; IR, internal rotation; K&L, Kellgren & Lawrence; L, left; LCEA, lateral center-edge angle; na, not applicable; NS, not significant; OA, osteoarthritis; R, right; RS, Rotterdam study; THR, total hip replacement; y, year

<sup>a</sup> If odds ratio per gender is not specified, the overall odds ratio is presented.