

<http://hdl.handle.net/1765/115968>



Physical fitness of children and adolescents with moderate to severe intellectual disabilities

Marieke Wouters, Heleen M. Evenhuis, Thessa I. M. Hilgenkamp.

Disability and Rehabilitation (in press)

ABSTRACT

Background: Low levels of physical fitness are associated with low physical and mental health. The aims of this study were to assess the health-related physical fitness of children with intellectual disability, and study the association of physical activity and motor development with physical fitness.

Methods: One hundred and twenty-eight children with moderate to severe intellectual disability (83 boys; age 2-18y) visiting specialized day program centers engaged in field-based physical fitness tests (body composition, muscular strength, muscular endurance and cardiorespiratory fitness). Scores were compared to reference values, and with linear regression analysis the association between the fitness outcomes and physical activity and motor development was studied.

Results: High rates of overweight (23-25%) and obesity (10-15%) were found. A majority of the participants (71-91%) scored below reference values for muscular strength, endurance and cardiorespiratory fitness tests. Physical activity and motor development were positively associated with scores on several fitness test ($\beta=.27-.44$; $p<.05$).

Conclusions: Children with moderate to severe intellectual disability visiting specialized day program centers have strikingly low physical fitness levels. Policies and interventions to increase the physical fitness for this specific group of children is urgently needed, in which increasing physical activity and motor skills are expected to be effective components.

INTRODUCTION

Low levels of health-related physical fitness have been found in children and adolescents with intellectual disability (ID) [1-5]. These low levels are alarming since children and adolescents with ID already start off with more health problems than typically developing (TD) children [6] and poor physical fitness is a risk factor for cardiovascular diseases, diabetes mellitus and poor mental health [7-10]. Several longitudinal studies show that risk factors of cardiovascular diseases, like hypercholesterolemia, hypertension and overweight transfer from childhood to adulthood [11, 12]. Furthermore, physical fitness is important to perform activities of daily living and low levels of strength and endurance will limit independence in adulthood [13, 14].

Such studies on physical fitness in children and adolescents with ID have mainly been conducted in children with mild to moderate ID. Children and adolescents with more severe levels of ID were usually not well-represented in those studies [15], which is understandable since evaluating these children poses some challenges, like limited understanding of the task. Furthermore, persons with ID experience more musculoskeletal problems than the general population; cerebral palsy, congenital deviation of feet and hip, hypotonia, and scoliosis are more likely to occur [16-18]. These conditions have an effect on the physical ability to perform tasks, but also on the potential to become physically active and fit, and develop motor skills. Other conditions like epilepsy [17], respiratory problems [19], sensorimotor dysfunction [20] can potentially limit these possibilities as well. In clinical practice, caregivers and parents often attribute the low levels of physical fitness, activity and motor skills to the cognitive and physical disabilities of the child or adolescent, and thereby underestimate the potential these youngsters have. Previous research has shown that improving the physical fitness, activity and motor skills in children and adolescents with ID is possible [21-25]. In the authors' experience, current care and support focuses mainly on activities of daily living skills, like getting dressed and eating independently, while there is hardly any focus on improving physical fitness.

Since more severe levels of ID are related to a higher risk of chronic health conditions, this lack of focus on physical fitness in children with more severe levels of ID is even more striking [17, 26]. Therefore, more information is needed on the physical fitness and its correlates, in this group of vulnerable children and adolescents. This information is relevant to prioritize the focus on physical fitness, and to identify child characteristics that are associated with low levels of physical fitness, to be able to target interventions to the most vulnerable groups in this population. Characteristics of interest are sex, age, having Down syndrome, level of ID, physical activity levels and motor development.

In TD children, boys have higher scores on physical fitness tests than girls, and older children have higher scores than younger children [27]. In children and adolescents with ID, these sex and age-effects are seen, but evidence is not that distinct [1, 3, 28, 29]. Having

Down syndrome (DS) and having more severe levels of ID are ID-specific child characteristics associated with low physical fitness [4, 30, 31]. These above-mentioned child characteristics are non-modifiable in order to improve the physical fitness.

Children's physical activity and motor development are parameters that can be influenced and that are positively associated with physical fitness [32-36]. Previous studies have shown that children and adolescents with ID have lower physical activity levels [37-40] and lower age-related motor competence [41] than TD peers. However, little research has been conducted on the contribution of physical activity and motor development to physical fitness in children and adolescents with ID [39]. The few studies that have been undertaken included participants with mild to moderate ID. Results on the relationship between physical fitness and activity were inconclusive: two studies found an association between moderate and/or vigorous physical activity and muscular and/or cardiorespiratory fitness [42, 43] and one did not find a significant relationship [44]. Physical fitness and motor competence were related in youth with ID in two studies [45, 46], but again, no information is available on this relationship for children with more severe ID. Furthermore, none of these studies included physical fitness, physical activity and motor development in the same analysis to study the complex interaction between these characteristics.

Taken together, knowledge on physical fitness levels in children with moderate to severe ID, and the relationship with physical activity and motor development, will help researchers and clinical professionals to better target interventions to increase physical fitness in this specific and vulnerable population. Moreover, more insight in the physical fitness levels and its correlates can motivate policy and decision makers to prioritize these themes in care and school settings. Therefore, the aims of this study were to assess the health-related physical fitness of children with moderate to severe ID visiting specialized day program centers, and to study the association of physical activity and motor development with physical fitness. Low physical fitness levels in this population are hypothesized, and physical fitness is hypothesized to be associated with physical activity and motor development.

METHODS

Participants

Participants were recruited for this cross-sectional study in the seven children's day program facilities of a service provider for people with disabilities in the Netherlands. These day program centers are specialized to support children that are unable to go to a mainstream or special school, due to their severe developmental delay or additional medical or behavioral comorbidity.

Children were invited to participate in the study if they had an age of 2 to 18 years with a moderate or severe level of ID, and sufficient motor capacities to walk independently. The

level of ID was evaluated by the behavioral therapist or psychologist of the child by reviewing existing testing results from the Bayley Scale of Infant and Toddler Development, Third Edition (BSID-III) [47], the Snijders-Oomen Nonverbal Intelligence Test [48], or comparable tests. Children with test-results corresponding an IQ of 20-40 (severe ID) or 40-55 (moderate ID) were selected. Parents or legal representatives of children who met the inclusion criteria received an invitation letter with information about the study, and were invited to sign the informed consent form. If informed consent was provided, parents or legal representatives completed the Physical Activity Readiness Questionnaire (PAR-Q [49] to determine if the participant could participate safely in the study. If there were any concerns, the physician was consulted for advice. Children were not invited if their behavioral therapist, psychologist, physician or nurse practitioner advised against participation in the study because of an expected negative impact on behavior or physical health of the child. Before testing started, the participant was asked to participate in the study. In case of any verbal or non-verbal resistance before or during the testing, the procedure was stopped immediately. This is according the Dutch code of conduct for 'minors' and 'mentally disabled' [50, 51].

Ethical approval was obtained (MEC-2013-491) from the Ethics Committee of the Erasmus Medical Center. The study adheres to the Declaration of Helsinki for research involving human subjects [52].

Measurements

Physical fitness tests

This study focused on the health-related physical fitness components according to the subdivision of the ACSM [53]: body composition, muscular strength, muscular endurance, and cardiorespiratory fitness. Flexibility is another component, however the relationship between flexibility and health has not been confirmed in children [54, 55] and therefore this component was not included in this study.

Appropriate tests for children with a moderate to severe ID were selected based on three steps. In short, first a literature review on the known psychometric properties of field-based physical fitness tests in children with ID was performed [15]. Secondly, the tests included in the review with proper psychometric properties were discussed with a focus group of physical therapists. The physical therapists discussed the feasibility of the tests (functionality, cognitive and motor demands), and face validity. Based on the outcome of the discussion, one or more suitable test per physical fitness component was selected for the pilot study. The pilot study was step three, in which selected tests were studied on feasibility and reliability. This was done in a sample of 39 children with moderate to severe ID, age 2-18 years [56]. Short-term (during one session) and long-term (2-4 weeks in between) test retest reliability was studied. Tests with sufficient reliability (intra class correlation $>.7$) were selected for the current study: body mass index (BMI), waist circumference, overarm throwing, stair climbing, and modified 6-minute walk test (6MWT).

Body composition

BMI was calculated as body weight in kg divided by height in meters squared. BMI-for-age-Z scores (zBMI) were calculated according to the WHO Growth references [57, 58]. Based on these Z-scores, weight status was determined. Participants were classified as underweight when zBMI was $<2SD$. Participants at age 0-5 years with BMI $>2SD$ were classified as overweight, $>3SD$ as obese. For older children (6-18 years), $>1SD$ was classified as overweight, and $>2SD$ as obese [59]. High test-retest reliability results for BMI were found in children with moderate to severe ID ($ICC>.99$) [56].

Waist circumference provides a measure of abdominal adiposity [60]. It was measured half way between the iliac crest and tenth rib, directly on the skin, using a flexible measuring tape accurate at the 0.1 cm level (Seca 201, Hamburg, Germany). High test-retest reliability results were found in this population ($ICC>.99$) [56]. The waist circumferences values were compared to age and sex matched Dutch reference values, in which the +1.3 SD was used as cut-off for overweight, and +2.3 SD was used as cut-off for obesity [61].

Muscular strength

The muscular strength test was selected from the Functional Strength Measurement (FSM), developed for TD children of 4-10 years old [62]. Only one muscular strength test of the FSM, the overarm throwing task, had sufficient test-retest reliability in children with moderate to severe ID ($ICC=.76$) [56] and was selected for this study. In children with mild ID, the overarm throwing test had moderate convergent validity with hand held dynamometry [63]. In this test, the participant stands behind a line and raises a bag of 2, 3 or 4 kg (dependent on the age) behind the head and throws the bag as far as possible. The distance from the line to the distal side of the bag is measured in cm. The actual test consists of three attempts, and the final score is the best score of these three attempts. Some participants needed extra guidance to perform this task correctly. In these cases, the test instructors physically or verbally guided the participants to starting position behind the line, and handed over the bag above their heads.

The muscular strength outcomes were compared to the reference values of 616 Dutch TD children, available for age 4-10 years [64]. The 5th and 95th percentile were used as cut-off values.

Muscular endurance

For this component another test of the FSM [62] was used: the stair climbing test. The participant walks the stairs as fast as possible up and down (three steps up and three down), turns and repeats this as many times as possible in 30 seconds. The amount of steps in 30 seconds is the score. The test consists of three attempts, and the final score is the best score of these three attempts. If the participant had difficulties with turning, the test instructor softly guided the turn.

In children with mild ID, the stair climbing test had moderate convergent validity with hand held dynamometry [63]. A pilot study indicated good test-retest reliability in children with moderate to severe ID (ICC=.92) [56]. However, it also indicated a significant improvement between the test and retest, therefore the children performed the test twice. The best score of the two trials was the final score. The muscular endurance outcomes were compared to the reference values of 616 Dutch TD children of 4-10 years [64]. The 5th and 95th percentile were used as cut-off values.

Cardiorespiratory fitness

The 6MWT is a test of functional capacity, and is recommended by the ACSM to use as field-based test for cardiorespiratory fitness in several populations with low fitness levels [53]. The 6MWT was found to be a valid and reliable tests in adolescents with overweight or obesity and mild-to moderate ID [65]. The test was modified to make it more feasible for children with moderate to severe ID [56]. During the modified 6MWT the participants walk as many meters as possible in six minutes over a 20-meter track, holding a loose hand of the instructor. The heart rate was continuously monitored (Polar RS400, Polar Electro Oy, Kempele, Finland). Peak heart rate was calculated afterwards. The modified 6MWT outcomes were compared to the reference values of 328 TD British children, age 4-11 years. Tenth and 90th percentiles were used as cut-off values [66]. The scores were also compared to an equation which predicts normal scores [67]: $\text{distance} = 11.89 \times \text{age (y)} + 486.1$. This equation was based on 496 Swiss children age 5-16 years.

Physical activity

Participants were asked to wear a of triaxial accelerometers (Actigraph GT3x+) on the right hip, for eight consecutive days. Physical activity was expressed by steps per day. Data were sampled with a frequency of 30Hz, and data was derived with 15 second epochs [68, 69]. Non-wear time was defined as ≥ 20 minutes of consecutive zeros, with no allowance of epochs with counts above zero [70]. Non-wear time was excluded from analysis. Data with at least 4 days of recording with a daily minimum of 8 registered hours were included in the analysis (Rich et al., 2013). In this selection procedure, no distinction was made between week days or weekend days, as no significant differences were found between the physical activity on week days or weekend days (data not shown).

Motor development

The gross motor subscale of the Bayley Scale of Infant and toddler Development, third edition [47] was completed by physical therapists to give insight into the gross motor development of the participants. The Bayley Scale of Infant and toddler Development is designed to measure the developmental status of young children up to 42 months, but it can also be used to assess the development of individuals with severe delays, such as children and adolescents

with ID [71]. Based on clinical experience with children and adolescents with moderate to severe ID and with motor testing, and the experience of consulted physical therapist in the field, the BSID-III was chosen. A score of 42-43 on the gross motor scale corresponds to the motor development of a TD child aged 12 months, a score of 57 to 24 months, 64 points to 36 months, and the maximum score (67-72 points) to 42 months [47].

Adaptive behavior

The relative age corresponding the adaptive behavior was used as a continuous indicator of the level of ID, since ID is a disability characterized by significant limitations both in intellectual functioning and in adaptive behavior [72]. Adaptive behavior was assessed by the Dutch version of the Vineland Adaptive Behavioral Scale [73, 74]. In this scale, three types of skills are covered: conceptual, social and practical skills. The scale was filled in by the caregiver of the child and scored and converted to relative age score by the concerned behavioral therapist or physiologist.

Other child characteristics

Information on autism spectrum disorder, and challenging behavior was provided by the behavioral therapist or psychologist of the participants. Information on age and DS was extracted from the medical file, information for socioeconomic status and origin (Dutch or non-Dutch) was provided by the parents. Socioeconomic status was based on the highest education of the parents: primary and secondary school, and lower or intermediate vocational school were classified as 'low socioeconomic status, higher vocational school or university were classified as 'high socioeconomic status. Children were considered as having a non-Dutch origin when at least one of their parents was born outside the Netherlands [75].

Procedure

The physical fitness tests were performed in a gymnasium at each of the seven-day program facilities during daytime, by one participant at a time. The tests were performed in a fixed order within the test session: body composition, modified 6-minute walk test (6MWT), stair climbing (1st trial), overarm throwing, stair climbing (2nd trial). In this sequence, the load on respectively the upper and lower extremities was taken into account, and familiarization with the tester and the test environment by starting with relatively easy tests as body composition and the modified 6MWT. Resting between tests was allowed. The instructors explained the tests and demonstrated how to perform it correctly. For the more difficult muscular strength and muscular endurance tests, the participants could practice up to five times. When they performed it correctly during this practice phase, the actual test started. During all tests, the instructor encouraged the participants to perform maximally. No standardized encouragement was used since every child had their own preferences and communication style. The test instructors judged if the participant understood the instruction, performed the test with

maximal effort, and according to protocol. When the performance was not meeting these requirements, the reason of drop-out was noted and no score was written down. The gross motor scale of the Bayley Scale of Infant and toddler Development-III was performed at another occasion, within a month of the fitness testing. All instructors (physical therapist or human movement scientist) had many years of experience with testing procedures and working with children with moderate to severe ID.

Analysis

Descriptive statistics were used to describe the participant characteristics. Group differences between the total group and group per test were determined using Pearson's Chi-square tests for dichotomous data, and independent t-tests for interval or ratio data, to investigate selective drop out.

The outcome on the several physical fitness tests were displayed by use of descriptive statistics. In order to compare the outcomes to the reference values, scatter plots were made, with the reference values plotted in the graphs. The percentages of participants that scored worse than the given age-related reference values were calculated. For the participants that fell outside the reference values due to their age, the reference value of the nearest age group was used.

To investigate the association of physical activity and motor development with the physical fitness scores, multiple linear regression analyses were performed. Each physical fitness test was a dependent variable. The following independent variables were entered in one block: sex, age, adaptive behavior (as continuous indicator of level of ID), the Bayley Scale of Infant and toddler Development score (motor development) and steps per day (physical activity). Because of the small number of participants with both valid accelerometer data and a score on either the overarm throwing or stair climbing test, steps per day was not entered in the regression analysis for these two fitness tests. Assumptions of multicollinearity and homoscedasticity were checked before conducting the analysis [76].

The data was analyzed by use of SPSS 24 (IBM statistics), Actilife 6 (Actigraph), and Excel (Microsoft 2016), and alpha level was set at 5%.

RESULTS

Participants

After inviting 219 children and adolescents to participate in the study, a total of 128 children with a moderate or severe ID were included. The flow diagram of the inclusion process can be found in Figure 1.

In Table 1 the characteristics of the participants are described. The total sample consisted of 83 boys and 45 girls, with an average age of 9.6 ± 4.1 y (range 2-18). More than half of the

participants had a severe ID ($n=73$, 57%), with an average adaptive behavior corresponding to the age of 1.4 ± 0.4 y. The children with a moderate ID ($n=54$, 43%) had an adaptive behavior corresponding to 2.7 ± 1.9 y. In total, 44 children had been diagnosed with autism spectrum disorder, 30 with DS. The included children were significantly younger than the non-included children (9.6 ± 4.1 vs 10.5 ± 3.7 y; $t(217)= 2.221$; $p=.03$), but no difference between groups was found with regard to sex ($\chi^2(1)=0.056$; $p=.81$).




Figure 1. Flow diagram of inclusion process

The included children performed at least one physical fitness test according to protocol, 46 participants had scores on all five fitness tests. The main reasons for drop out were challenging behavior, cognitive issues and motor development issues. This was comparable to the drop out that was encountered in the pilot study [56]. Sample size per test ranged from 59 for overarm throwing to 118 for BMI (Table 2). Because of the variable sample size per test, participant characteristics differed per test. However, only for overarm throwing and stair climbing significant differences compared to the total sample were found; the children that correctly performed the overarm throwing test consisted of relatively more children with moderate ID (64%; $\chi^2(1)=7.7$; $p<.01$), had more developed adaptive behavior (2.6 ± 1.8 y; $t(156)=-2.67$; $p=.01$) and a better motor development (63.6 ± 5.9 ; $t(167)=-3.51$; $p<.01$). For stair climbing, the children also had better motor development than the total group (63.6 ± 5.5 ; $t(175)=-3.38$; $p<.01$). The participants' characteristics per test can be found in the Appendix Table A1.

Sixty-eight participants had enough valid accelerometer data to be included in the analysis. For the other participants, data transfer problems occurred ($n=15$), not enough valid data

was on the accelerometer ($n=31$), or the accelerometer got lost ($n=1$). For 58 of the 68 children, at least one weekend day was included. Characteristics of the participants with physical activity data were not significantly different from those of the non-participants [77].

Physical fitness levels

For body composition, BMI and waist circumference was correctly assessed in respectively 118 and 116 children. Average BMI was $19.3 \pm 4.4 \text{ kg/m}^2$, average zBMI 0.82 ± 1.21 (Table 2). Based on these zBMI scores, 3% of the children were classified as having underweight, 25% as having overweight, and 15% as having obesity (Figure 2a). The average waist circumference was $65.4 \pm 15.1 \text{ cm}$ (Table 2). Twenty-seven children (23%) were labeled as overweight, and 12 children (10%) were labeled as obese. Graphs of the scores compared to the reference values can be found in Figure 2b and 2c.

Table 1. Participants' characteristics for the total group.

	N	%	M	SD
N	128			
Sex				
Boys	83	(65)		
Girls	45	(35)		
Age (y)	128		9.6	± 4.1
2-6 y	44	(34)		
7-12 y	44	(34)		
13-18 y	40	(31)		
Level of ID				
Moderate ID	54	(43)		
Severe ID	73	(57)		
Adaptive behavior (y)	111		1.9	± 1.3
Autism Spectrum Disorder	44	(34)		
Down Syndrome	30	(23)		
Challenging behavior	86	(67)		
Low social economic status	46	(36)		
Non-Dutch origin	33	(26)		
Score on Bayley scale of infant development	118		59.5	± 7.4
Steps per day	68		6677	± 2600

Note. ID = intellectual disability.

The muscular strength test was performed by fifty-nine participants according to protocol. Average throwing distance was $155 \pm 84 \text{ cm}$ (Table 2, Figure 2d). Compared to the reference values, 42 participants (71%) scored below the P5 cut-off values.

The muscular endurance test was correctly completed by 63 participants. Their maximum score of the two trials on the stair climbing test was on average 32 ± 17 steps in 30 seconds (Table 2). The majority of the participants achieved the maximum score in the second trial ($n=39$, 62%), 13% ($n=8$) in first trial, and 25% ($n=16$) had the same score in both trials. Compared to the reference values, 57 participants (91%) scored below the P5 cut-off values.

The modified 6MWT, as indicator of cardiorespiratory fitness, was correctly performed by 102 children. They walked an average distance of 349 ± 98 meters in six minutes (Table 2), with an average HR_{peak} of 142 ± 17 beats·min⁻¹ ($n=79$). Compared to the reference values of Lammers, Hislop [66], 77 participants (75%) scored below the P10 cut-off values. None of the children made enough distance to reach the predicted distance by Ulrich, Hildenbrand [67].

In the Appendix Table A2, separated scores for boys and girls on the physical fitness tests can be found.

Table 2. Average score on the physical fitness tests, and percentage participants scoring worse than reference values.

	n	Score on test		Scoring worse than reference values ¹	
		M \pm SD	[95% CI]	%	[95% CI]
BMI (kg/m ²)	118	19.3 \pm 4.4	[18.5-20.1]		
zBMI	118	0.82 \pm 1.21	[0.59-1.04]	40	[31-49]
Waist circumference (cm)	116	65.4 \pm 15.1	[62.6-68.2]	34	[25-42]
Overarm throwing (cm)	59	155 \pm 84	[133-177]	71	[59-83]
Stair climbing (n)	63	32 \pm 17	[28-37]	91	[85-97]
Modified 6MWT (m)	102	349 \pm 98	[330-368]	75	[67-84]


Notes. CI= confidence interval; BMI=body mass index; 6MWT = six minute walk test; ¹ Classified as >2SD (0-5y), >1SD (6-18y) for BMI [59], >1.3SD for waist circumference [61], < P5 for overarm throwing and stair climbing [64], and <P10 for modified 6MWT [66].

Associations with physical fitness


Motor development was positively associated with overarm throwing ($\beta=.27$, $p=.02$), stair climbing ($\beta=.44$, $p<.001$), and the modified 6MWT ($\beta=.34$ $p=.02$). Increasing number of steps per day was only associated with an increasing score on the modified 6MWT ($\beta=.33$, $p<.01$).

DISCUSSION


This study indicates that children with moderate to severe ID visiting specialized day program centers have very low levels of health-related physical fitness. The relationship of the modifiable factors physical activity and motor development with cardiorespiratory fitness and




A. zBMI




B. Waist circumference – males




C. Waist circumference - females



D. Overarm throwing



E. Stair climbing



F. Modified 6mwt

Figure 2. Scatterplots of the individual scores on the physical fitness test, with reference values plotted in the graphs

muscular strength and endurance indicate a potential way to improve physical fitness in this population.

The current findings of low physical fitness levels in this specific group of children compared to TD children are similar to previous findings in children and adolescents with less severe levels of ID. For example, Dutch children aged 8-11 y with borderline to mild ID scored significantly lower on the handgrip, standing broad jump, sit-ups and 20 meter shuttle run

Table 3. Association of fitness outcomes and predictor variables, assessed by a multiple linear regression analysis

	BMI (kg/m ² ; n=54)		zBMI (n=54)		Waist circumference (cm; n=54)		Overarm throwing (cm; n=41)		Stair climbing (n; n=46)		Modified 6MWT (m; n=51)	
	B	SE	B	SE	B	SE	B	SE	B	SE	B	SE
Adjusted R ²	41%		9%		61%		73%		69%		51%	
Constant	18.45	4.90	3.06	1.88	36.15	13.32	-204.69	104.74	-61.95	22.21	-141.59	112.50
Sex ^b	0.28	1.02	.03	-0.14	0.40	-0.06	3.29	2.73	.13	-23.57	17.13	-1.13
Age ^a	0.31	0.11	.33	**	-0.11	0.04	-0.37	*	2.00	0.29	.65	***
Adaptive behavior ^c	1.12	0.32	.47	**	0.19	0.12	.26	1.48	0.86	.17	24.22	5.01
Motor development ^d	-0.05	0.08	-0.09	-0.01	0.03	-0.06	0.10	0.21	.05	4.04	1.59	.27
Physical activity ^e	-0.33	0.20	-0.22	0.14	0.08	-0.28	-0.74	0.51	-0.14	-	-	-

Notes: BMI=body mass index; zBMI=BMI standardized for age; 6MWT= six minute walk test; DS=Down syndrome; ^a in years; ^b 1=male; 2=female; ^c Vineland Z score; ^d Bayley scale of infant development, gross motor scale score; ^e Steps per day, unit per 1000; Bold font indicates significant associations; * p<.05; ** p<.01; *** p<.001; #p=.052.



tests compared to TD peers [1]. Comparable results were seen for adolescents with mild to moderate ID [3], adolescents with DS [2], and children with borderline to mild ID [4]. High overweight rates were seen before as well [3, 5]. The current rates of overweight and obesity are much higher than those of TD Dutch children: 40% vs. 13-15% [78]. A positive association was found between physical activity and cardiorespiratory fitness, suggesting that cardiorespiratory fitness of youth with moderate to severe ID might be improved with increasing the amount of physical activity. These results are in agreement with intervention studies in youth with ID and TD youth which indicated positive effects of physical activity, like exercise, on several physical fitness outcomes [23, 79-83].

The current results show an association between physical fitness and motor development, for the components muscular strength, muscular endurance and cardiorespiratory fitness. This is in agreement with the findings in TD children that high levels of motor skills are related to high levels of physical fitness [84-86]. The current results are comparable to two Asian studies in samples of over 400 children and adolescents with mild to moderate ID [45, 46]. Both studies found statistically significant correlations between motor skills and muscular strength, muscular endurance and CR fitness ($r=.19-.52$; $p<.05$).

In the current study, body composition was not related to physical activity and motor development. This is in line with previous studies in youth with ID [42, 43, 45, 46, 87]. However, in TD children, an association between body composition and physical activity and motor development is seen [88-90]. This difference could be explained by the body composition measures used. The studies in TD children used body fat measures, like skinfold thickness [88-90]. In the studies with participants with ID BMI and waist circumference was measured which are anthropometric measures that are surrogate measures of body fatness. These measures cannot discriminate between fat mass and fat free mass. Another explanation could be the study population. In youth with ID, other factors might be more strongly related to body composition. Factors associated with overweight and obesity in people with ID are nutritional habits, psychotropic medication and altered metabolic rate [91]. However, physical activity interventions did result in weight loss in children and adolescents with ID [92, 93], which suggest that a relationship between physical activity and body composition does exist.

The physical fitness tests used in the current study, were tested on feasibility and reliability in a pilot study of 37 participants [56]. Even though high drop-out rates were seen for the muscular strength and endurance tests both in the current study and in the pilot study, the test-retest reliability was sufficient for group analysis ($ICC >.75$). Further research is required to find appropriate tests for the children and adolescents with moderate to severe ID, who were not able to perform the selected tests. Especially the children with a more severe level of ID, and children with less motor abilities, were less able to perform the muscular strength and muscular endurance tests. For these components more research is needed on tests which demand little cognitive and motor competence.

Implications

Even though children and adolescents with ID have to deal with musculoskeletal problems and other physical challenges [16-18], improvements in physical fitness, physical activity and motor skills are possible [21, 22, 94-100]. For example, motor skill training sessions of 10-15 minutes, three to five times a week, can result in significant improvements in motor functioning in persons with severe to profound ID [22].

In the Netherlands, the specialized day program centers are not obligated to organize physical education, in contrast to regular primary schools. The children and adolescents with ID are physically active during day activities, but do not have physical education sessions supervised by skilled professionals. Considering the low physical fitness levels, associated with physical activity and motor development, found in this study, and the vulnerable population, the government and care organizations should make regular physical education supervised by skilled professionals a central standard for specialized day program centers.

Limitations

The group participants consisted of 128 children and adolescents with moderate to severe ID. However, the sample size per test ranged from 59 to 118 children. Drop-out in the different tests has caused a bias in the muscular strength and muscular endurance tests, in which children with lower adaptive behavior, and less motor development were less likely to perform the tests correctly.

For physical activity steps per day was chosen as outcome measure. This measure reflects mainly the vertical axis activity, and can therefore be seen as participants' walking behavior. This measure was chosen because walking is likely to be the primary activity persons with ID engage in [101, 102]. However, choosing one outcome measure for physical activity is a limitation of the study. Steps per day gives no information on the amount of minutes and intensity of physical activity in which the participants engaged in.

The BSID test was used to indicate motor development, was originally developed for TD children up to 42 months. Potentially, this could have caused a ceiling effect, since the participants in the current study were between the 2 and 18 years old. However, the scores were evenly distributed (range 42-72 points), with only one participant reaching the maximum score (data not shown).

Due to the small sample sizes per test, this study was only powered to control for a few child characteristics in the linear regression analysis. The variables that were controlled for, were sex, age and adaptive behavior (as indicator of level of ID). This was based on previous studies in TD children and children with ID [1, 30]. Preferably, future studies with larger sample size should control for DS, social economic status and origin as well, since previous research has shown that these variables were also associated with physical fitness [103-105]. With regards to the influence of DS, previous research has shown that persons with DS have lower physical fitness values than persons with other causes of ID [31]. Certain characteristics

or comorbidities of DS can explain these differences. One of the explanations of these low fitness values is the altered autonomic function [106]. Since 23% of the current sample had DS, this might have negatively influenced the outcomes. However, univariate analyses indicated that the participants with DS did not score significantly lower than the participants with other causes of ID (data not shown).

The reference values that were used had limitation as well. Only for body composition standardized values for age were calculated. For the other fitness components, only norm-referenced cut-off scores per age were available, and only for a small age-bandwidth. To compare the scores of the participants that fell outside the bandwidth, the reference values of nearest age group were used, but these reference values most likely underestimate the actual 95th percentile. Moreover, it is likely that a sex difference will appear during puberty on these components [107] and the reference values that were used, were not sex-specific.

Due to the cross-sectional design of the study, no comment can be made on the direction of the relationship between physical fitness, physical activity and motor development. In the general population these relationships are suggested to be bidirectional and dynamic [84].

Conclusion

Children with moderate to severe ID visiting specialized day program centers have high rates of overweight and obesity, and low levels of muscular strength, muscular endurance and cardiorespiratory fitness. Policies and interventions to increase the physical fitness for this specific group of children are urgently needed, in which increasing motor skills and physical activity should have a central place.

REFERENCES

1. Hartman, E., et al., *Development of physical fitness in children with intellectual disabilities*. J Intellect Disabil Res, 2015. **59**(5): p. 439-49.
2. Izquierdo-Gomez, R., et al., *Are Poor Physical Fitness and Obesity Two Features of the Adolescent with down Syndrome?* Nutr Hosp, 2013. **28**(4): p. 1348-1351.
3. Salaun, L. and S.E. Berthouze-Aranda, *Physical fitness and fatness in adolescents with intellectual disabilities*. J Appl Res Intellect Disabil, 2012. **25**(3): p. 231-9.
4. Golubovic, S., et al., *Effects of exercise on physical fitness in children with intellectual disability*. Res Dev Disabil, 2012. **33**(2): p. 608-14.
5. Slevin, E., et al., *Obesity and overweight in intellectual and non-intellectually disabled children*. J Intellect Disabil Res, 2014. **58**(3): p. 211-20.
6. Oeseburg, B., et al., *Prevalence of chronic health conditions in children with intellectual disability: a systematic literature review*. Intellect Dev Disabil, 2011. **49**(2): p. 59-85.
7. Ortega, F.B., et al., *Physical fitness in childhood and adolescence: a powerful marker of health*. Int J Obes (Lond), 2008. **32**(1): p. 1-11.
8. Hurtig-Wennlof, A., et al., *Cardiorespiratory fitness relates more strongly than physical activity to cardiovascular disease risk factors in healthy children and adolescents: the European Youth Heart Study*. Eur J Cardiovasc Prev Rehabil, 2007. **14**(4): p. 575-81.
9. Ortega, F.B., et al., *Cardiorespiratory fitness and sedentary activities are associated with adiposity in adolescents*. Obesity (Silver Spring), 2007. **15**(6): p. 1589-99.
10. Anderssen, S.A., et al., *Low cardiorespiratory fitness is a strong predictor for clustering of cardiovascular disease risk factors in children independent of country, age and sex*. Eur J Cardiovasc Prev Rehabil, 2007. **14**(4): p. 526-31.
11. Raitakari, O.T., et al., *Cardiovascular risk factors in childhood and carotid artery intima-media thickness in adulthood: the Cardiovascular Risk in Young Finns Study*. JAMA, 2003. **290**: p. 2277-2283.
12. Andersen, L.B., et al., *The relationship between physical fitness and clustered risk, and tracking of clustered risk from adolescence to young adulthood: eight years follow-up in the Danish Youth and Sport Study*. Int J Behav Nutr Phys Act, 2004. **1**: p. 6.
13. Hilgenkamp, T.I.M., R. van Wijck, and H.M. Evenhuis, *(Instrumental) activities of daily living in older adults with intellectual disabilities*. Research in Developmental Disabilities, 2011. **32**(5): p. 1977-1987.
14. Oppewal, A., et al., *Physical fitness is predictive for a decline in daily functioning in older adults with intellectual disabilities: results of the HA-ID study*. Res Dev Disabil, 2014. **35**(10): p. 2299-315.
15. Wouters, M., H.M. Evenhuis, and T.I.M. Hilgenkamp, *Systematic review of field-based physical fitness tests for children and adolescents with intellectual disabilities*. Res Dev Disabil, 2017. **61**: p. 77-94.
16. Vonken, M.T.H., M.A. Maaskant, and M. van den Akker, *Prevalentie en risicogroepen voor aandoeningen van het bewegingsapparaat bij mensen met een verstandelijke beperking* NTZ, 2006. **2**: p. 98-110.
17. van Schroyensteen Lantman-de Valk, H.M., et al., *Prevalence and incidence of health problems in people with intellectual disability*. J Intellect Disabil Res, 1997. **41** (Pt 1): p. 42-51.
18. Bilo, R.A.C. and H.W.A. Voorhoeve, *Kind in ontwikkeling. Een handreiking bij de observatie van jonge kinderen*. 7th ed. 2007, Houten, Netherlands: Bohn Stafleu van Loghum

19. Seddon, P.C. and Y. Khan, *Respiratory problems in children with neurological impairment*. Archives of Disease in Childhood, 2003. **88**(1): p. 75-78.
20. Wuang, Y.P., et al., *Profiles and cognitive predictors of motor functions among early school-age children with mild intellectual disabilities*. J Intellect Disabil Res, 2008. **52**(12): p. 1048-60.
21. Hocking, J., J. McNeil, and J. Campbell, *Physical therapy interventions for gross motor skills in people with an intellectual disability aged 6 years and over: a systematic review*. Int J Evid Based Healthc, 2016.
22. Houwen, S., A. van der Putten, and C. Vlaskamp, *A systematic review of the effects of motor interventions to improve motor, cognitive, and/or social functioning in people with severe or profound intellectual disabilities*. Res Dev Disabil, 2014. **35**(9): p. 2093-116.
23. Dodd, K.J. and N. Shields, *A systematic review of the outcomes of cardiovascular exercise programs for people with Down syndrome*. Arch Phys Med Rehabil, 2005. **86**(10): p. 2051-8.
24. Frey, G.C., V.A. Temple, and H.I. Stanish, *Interventions to promote physical activity for youth with intellectual disabilities*. Salud Pública de México, 2017. **59**(4): p. 437-45.
25. Shin, I.S. and E.Y. Park, *Meta-analysis of the effect of exercise programs for individuals with intellectual disabilities*. Res Dev Disabil, 2012. **33**(6): p. 1937-47.
26. Moss, S., et al., *Physical morbidity in older people with moderate, severe and profound mental handicap, and its relation to psychiatric morbidity*. Soc Psychiatry Psychiatr Epidemiol, 1993. **28**(1): p. 32-9.
27. Welk, G.J., et al., *Field evaluation of the new FITNESSGRAM(R) criterion-referenced standards*. Am J Prev Med, 2011. **41**(4 Suppl 2): p. S131-42.
28. Lin, J.-D., et al., *Patterns of Obesity among Children and Adolescents with Intellectual Disabilities in Taiwan*. Journal of Applied Research in Intellectual Disabilities, 2005. **18**(2): p. 123-129.
29. Mercer, V.S. and C.L. Lewis, *Hip Abductor and Knee Extensor Muscle Strength of Children with and without Down Syndrome*. Pediatr Phys Ther, 2001. **13**(1): p. 18-26.
30. Skowronski, W., et al., *Eurofit special: European fitness battery score variation among individuals with intellectual disabilities*. Adapt Phys Activ Q, 2009. **26**(1): p. 54-67.
31. Pitetti, K.H., T. Baynard, and S. Agiovlasis, *Children and adolescents with Down syndrome, physical fitness and physical activity*. Journal of Sport and Health Science, 2013. **2**(1): p. 47-57.
32. Malina, R.M., *Physical activity and fitness: pathways from childhood to adulthood*. Am J Hum Biol, 2001. **13**(2): p. 162-72.
33. Hands, B., et al., *The relationship among physical activity, motor competence and health-related fitness in 14-year-old adolescents*. Scand J Med Sci Sports, 2009. **19**(5): p. 655-63.
34. Kristensen, P.L., et al., *The association between aerobic fitness and physical activity in children and adolescents: the European youth heart study*. Eur J Appl Physiol, 2010. **110**(2): p. 267-75.
35. Burgi, F., et al., *Relationship of physical activity with motor skills, aerobic fitness and body fat in preschool children: a cross-sectional and longitudinal study (Ballabeina)*. Int J Obes (Lond), 2011. **35**(7): p. 937-44.
36. Cattuzzo, M.T., et al., *Motor competence and health related physical fitness in youth: A systematic review*. J Sci Med Sport, 2016. **19**(2): p. 123-9.
37. Einarsson, I.T., et al., *Physical activity during school and after school among youth with and without intellectual disability*. Res Dev Disabil, 2016. **56**: p. 60-70.
38. Einarsson, I.T., et al., *Differences in physical activity among youth with and without intellectual disability*. Med Sci Sports Exerc, 2015. **47**(2): p. 411-8.
39. Frey, G.C., H.I. Stanish, and V.A. Temple, *Physical activity of youth with intellectual disability: review and research agenda*. Adapt Phys Activ Q, 2008. **25**(2): p. 95-117.

40. Hinckson, E.A. and A. Curtis, *Measuring physical activity in children and youth living with intellectual disabilities: a systematic review*. Res Dev Disabil, 2013. **34**(1): p. 72-86.
41. Hartman, E., et al., *On the relationship between motor performance and executive functioning in children with intellectual disabilities*. J Intellect Disabil Res, 2010. **54**(5): p. 468-77.
42. Izquierdo-Gomez, R., et al., *Associations of physical activity with fatness and fitness in adolescents with Down syndrome: The UP&DOWN study*. Research in Developmental Disabilities, 2015. **36**: p. 428-436.
43. Matute-Llorente, A., et al., *Physical activity and cardiorespiratory fitness in adolescents with Down syndrome*. Nutricion Hospitalaria, 2013. **28**(4): p. 1151-1155.
44. Shields, N., et al., *An exploratory study of the association between physical activity, cardiovascular fitness and body size in children with Down syndrome*. Developmental Neurorehabilitation, 2017. **20**(2): p. 92-98.
45. Frey, G.C. and B. Chow, *Relationship between BMI, physical fitness, and motor skills in youth with mild intellectual disabilities*. International Journal of Obesity, 2006. **30**(5): p. 861-867.
46. Foley, J.T., et al., *The relationships among fundamental motor skills, health-related physical fitness, and body fatness in South Korean adolescents with mental retardation*. Research Quarterly for Exercise and Sport, 2008. **79**(2): p. 149-157.
47. Bayley, N., *Bayley scales of infant and toddler development, third edition 2006*, San Antonio, TX: Pearson Education, Inc.
48. Tellegen, P.J., et al., *Snijders-Oomen Nonverbal Intelligence Test, SON -R 2½-7, Manual & Research Report*. . 1998, Lisse, the Netherlands: Swets & Zeitlinger.
49. Thomas, S., J. Reading, and R.J. Shephard, *Revision of the Physical Activity Readiness Questionnaire (PAR-Q)*. Can J Sport Sci, 1992. **17**(4): p. 338-45.
50. CCMO. *Code of conduct mentally disabled*. 2002 [cited 2018 October 1]; Available from: <http://www.ccmo.nl/attachments/files/code-of-conduct-mental-disabled.pdf>.
51. CCMO. *Code of conduct minors*. 2002 [cited 2018 October 1]; Available from: <http://www.ccmo.nl/attachments/files/code-of-conduct-minors.pdf>.
52. World Medical Association. *Declaration of Helsinki - Ethical Principles for Medical Research Involving Human Subjects*. 2013 [cited 2018 December 12]; Available from: <https://www.wma.net/what-we-do/medical-ethics/declaration-of-helsinki/doh-oct2013-jama-2/>.
53. American College of Sports Medicine, *ACSM's Guidelines for Exercise Testing and Prescription*. 10th ed. 2017, Philadelphia, PA: Lippincott Williams & Wilkins.
54. Ganley, K.J., et al., *Health-related fitness in children and adolescents*. Pediatr Phys Ther, 2011. **23**(3): p. 208-20.
55. Ruiz, J.R., et al., *Predictive validity of health-related fitness in youth: a systematic review*. Br J Sports Med, 2009. **43**: p. 909-923.
56. Wouters, M., et al., *Feasibility and Reliability of Tests Measuring Health-Related Physical Fitness in Children With Moderate to Severe Levels of Intellectual Disability*. Am J Intellect Dev Disabil, 2017. **122**(5): p. 422-438.
57. de Onis, M., et al., *Development of a WHO growth reference for school-aged children and adolescents*. Bull World Health Organ, 2007(85): p. 661-8.
58. WHO Multicentre Growth Reference Study Group, *WHO Child Growth Standards: Length/height-for-age, weight-for-age, weight-for-length, weight-for-height and body mass index-for-age: Methods and development*. . 2006, World Health Organization: Geneva. p. 312.
59. de Onis, M. and T. Lobstein, *Defining obesity risk status in the general childhood population: which cut-offs should we use?* Int J Pediatr Obes, 2010. **5**(6): p. 458-60.

60. Taylor, R.W., et al., *Evaluation of waist circumference, waist-to-hip ratio, and the conicity index as screening tools for high trunk fat mass, as measured by dual-energy X-ray absorptiometry, in children aged 3-19 y.* Am J Clin Nutr, 2000. **72**(2): p. 490-5.
61. Fredriks, A.M., et al., *Are age references for waist circumference, hip circumference and waist-hip ratio in Dutch children useful in clinical practice?* Eur J Pediatr, 2005. **164**(4): p. 216-22.
62. Aertssen, W.F., G.D. Ferguson, and B.C. Smits-Engelsman, *Reliability and Structural and Construct Validity of the Functional Strength Measurement in Children Aged 4 to 10 Years.* Phys Ther, 2016. **96**(6): p. 888-97.
63. Aertssen, W.F.M., B. Steenbergen, and B.C.M. Smits-Engelsman, *The validity and reliability of the Functional Strength Measurement (FSM) in children with intellectual disabilities.* J Intellect Disabil Res, 2018.
64. Smits-Engelsman, B.C.M. and W.F.M. Verhoef-Aertssen, *Functional Strength Measurement (FSM): Manual.* . 2012, Meteren, the Netherlands: FSM Production.
65. Elmahgoub, S.S., et al., *Reproducibility, validity and predictors of six-minute walk test in overweight and obese adolescents with intellectual disability.* Disabil Rehabil, 2012. **34**(10): p. 846-51.
66. Lammers, A.E., et al., *The 6-minute walk test: normal values for children of 4-11 years of age.* Arch Dis Child, 2008. **93**(6): p. 464-8.
67. Ulrich, S., et al., *Reference values for the 6-minute walk test in healthy children and adolescents in Switzerland.* BMC Pulm Med, 2013. **13**: p. 49.
68. Reilly, J.J., et al., *Objective measurement of physical activity and sedentary behaviour: review with new data.* Arch Dis Child, 2008. **93**(7): p. 614-9.
69. Cliff, D.P., J.J. Reilly, and A.D. Okely, *Methodological considerations in using accelerometers to assess habitual physical activity in children aged 0-5 years.* J Sci Med Sport, 2009. **12**(5): p. 557-67.
70. Esliger, D.W., et al., *Standardizing and Optimizing the Use of Accelerometer Data for Free-Living Physical Activity Monitoring.* Journal of Physical Activity & Health, 2005. **2**(3): p. 366.
71. Pearson Education, *Technical Report. Using the Bayley Scales of Infant and Toddler Development, Third Edition, to Assess Individuals with Severe Delays.* 2008.
72. Schalock, R.L., et al., *Intellectual Disability: Definition, Classification, and Systems of Supports.* 11th ed. Vol. 2010, 2010, Washington, D.C: American Association on Intellectual and Developmental Disabilities
73. de Bildt, A.A. and D.W. Kraijer, *Vineland-Z, sociale redzaamheidsschaal voor kinderen en jeugdigen met een verstandelijke beperking, handleiding [Vineland-Z, social independence scale for children and adolescents with an intellectual disability, manual].* 2003, Leiden: PITS B.V.
74. Sparrow, S.S., D.A. Balla, and D.V. Cicchetti, *Vineland adaptive behavior scales.* 1984, Circle Pines, MN: American Guidance Service.
75. Centraal Bureau voor de Statistiek, *Standaarddefinitie allochtonen [Standard definition immigrants].* 2000, Centraal Bureau voor de Statistiek:: Voorburg, the Netherlands.
76. Field, A., *Discovering Statistics using SPSS.* 2nd ed. 2005, London: Sage Publications Ltd.
77. Wouters, M., H.M. Evenhuis, and T.I.M. Hilgenkamp, *Physical activity levels of children and adolescents with moderate-to-severe intellectual disability.* J Appl Res Intellect Disabil, 2019. **32**(1): p. 131-142.
78. Schonbeck, Y., et al., *Increase in prevalence of overweight in Dutch children and adolescents: a comparison of nationwide growth studies in 1980, 1997 and 2009.* PLoS One, 2011. **6**(11): p. e27608.

79. Casey, A.F. and R. Rasmussen, *Reduction measures and percent body fat in individuals with intellectual disabilities: a scoping review*. Disabil Health J, 2013. **6**(1): p. 2-7.
80. Gonzalez-Aguero, A., et al., *Health-related physical fitness in children and adolescents with Down syndrome and response to training*. Scandinavian Journal of Medicine & Science in Sports, 2010. **20**(5): p. 716-724.
81. Calders, P., S. Elmahgoub, and D. Cambier, *Physical and Metabolic Fitness of Children and Adolescents with Intellectual Disability - How to Rehabilitate?*, in *Latest Findings in Intellectual and Developmental Disabilities Research*, P.U. Tan, Editor. 2012, InTech.
82. Janssen, I. and A.G. Leblanc, *Systematic review of the health benefits of physical activity and fitness in school-aged children and youth*. Int J Behav Nutr Phys Act, 2010. **7**: p. 40.
83. Collins, K. and K. Staples, *The role of physical activity in improving physical fitness in children with intellectual and developmental disabilities*. Res Dev Disabil, 2017. **69**: p. 49-60.
84. Stodden, D.F., et al., *Dynamic relationships between motor skill competence and health-related fitness in youth*. Pediatr Exerc Sci, 2014. **26**(3): p. 231-41.
85. Stodden, D.F., S.J. Langendorfer, and M.A. Robertson, *The association between motor skill competence and physical fitness in young adults*. Res Q Exerc Sport, 2009. **80**(2): p. 223-9.
86. Wrotniak, B.H., et al., *The relationship between motor proficiency and physical activity in children*. Pediatrics, 2006. **118**(6): p. e1758-65.
87. Esposito, P.E., et al., *Physical activity patterns of youth with Down syndrome*. Intellect Dev Disabil, 2012. **50**(2): p. 109-19.
88. Andersen, L.B., et al., *Physical activity and clustered cardiovascular risk in children: a cross-sectional study (The European Youth Heart Study)*. Lancet, 2006. **368**(9532): p. 299-304.
89. Twisk, J.W., et al., *The relation between "long-term exposure" to lifestyle during youth and young adulthood and risk factors for cardiovascular disease at adult age*. J Adolesc Health, 1997. **20**(4): p. 309-19.
90. Ortega, F.B., et al., *Cardiovascular fitness modifies the associations between physical activity and abdominal adiposity in children and adolescents: the European Youth Heart Study*. Br J Sports Med, 2010. **44**(4): p. 256-62.
91. Must, A., et al., *Obesity Prevention for Children with Developmental Disabilities*. Current obesity reports, 2014. **3**(2): p. 156-170.
92. Ordonez, F.J., M. Rosety, and M. Rosety-Rodriguez, *Influence of 12-week exercise training on fat mass percentage in adolescents with Down syndrome*. Med Sci Monit, 2006. **12**(10): p. CR416-9.
93. Elmahgoub, S.M., et al., *The influence of combined exercise training on indices of obesity, physical fitness and lipid profile in overweight and obese adolescents with mental retardation*. Eur J Pediatr, 2009. **168**(11): p. 1327-33.
94. Giagazoglou, P., et al., *Effects of a trampoline exercise intervention on motor performance and balance ability of children with intellectual disabilities*. Res Dev Disabil, 2013. **34**(9): p. 2701-7.
95. Fotiadou, E.G., et al., *The Effect of a Psychomotor Education Program on the Static Balance of Children With Intellectual Disability*. J Strength Cond Res, 2017. **31**(6): p. 1702-1708.
96. Lee, K., M. Lee, and C. Song, *Balance training improves postural balance, gait, and functional strength in adolescents with intellectual disabilities: Single-blinded, randomized clinical trial*. Disabil Health J, 2016. **9**(3): p. 416-22.
97. Lotan, M., et al., *Physical fitness and functional ability of children with intellectual disability: effects of a short-term daily treadmill intervention*. ScientificWorldJournal, 2004. **4**: p. 449-57.

98. Lotan, M., E. Isakov, and J. Merrick, *Improving functional skills and physical fitness in children with Rett syndrome*. J Intellect Disabil Res, 2004. **48**(Pt 8): p. 730-5.
99. Shin, J.Y., et al., *The effects of a home-based intervention for young children with intellectual disabilities in Vietnam*. J Intellect Disabil Res, 2009. **53**(4): p. 339-52.
100. Bishop, J.C. and M. Pangelinan, *Motor skills intervention research of children with disabilities*. Res Dev Disabil, 2018. **74**: p. 14-30.
101. Lin, J.D., et al., *Physical activity and its determinants among adolescents with intellectual disabilities*. Res Dev Disabil, 2010. **31**(1): p. 263-9.
102. Draheim, C.C., D.P. Williams, and J.A. McCubbin, *Prevalence of Physical Inactivity and Recommended Physical Activity in Community-Based Adults With Mental Retardation*. Mental Retardation, 2002. **40**(6): p. 436-444.
103. Pitetti, K.H. and B. Fernhall, *Comparing run performance of adolescents with mental retardation, with and without Down syndrome*. Adapt Phys Activ Q, 2004. **21**(3): p. 219-228.
104. McMurray, R.G., et al., *The influence of physical activity, socioeconomic status, and ethnicity on the weight status of adolescents*. Obes Res, 2000. **8**(2): p. 130-9.
105. Labree, L.J., et al., *Differences in overweight and obesity among children from migrant and native origin: a systematic review of the European literature*. Obes Rev, 2011. **12**(5): p. e535-47.
106. Fernhall, B., G.V. Mendonca, and T. Baynard, *Reduced work capacity in individuals with Down syndrome: a consequence of autonomic dysfunction?* Exerc Sport Sci Rev, 2013. **41**(3): p. 138-47.
107. Wilmore, J.H., D.L. Costill, and W.L. Kenney, *Physiology of Sport and Exercise*. 4th edition ed. 2008, Champaign, United States: Human Kinetics Publishers.

APPENDIX

Table A1. Participants' characteristics (in n (%) or mean \pm standard deviation) per performed test.

	BMI	Waist circumference	Overarm throwing	Stair climbing	Modified 6MWT
N	118	116	59	63	102
Sex					
Boys	75 (64)	73 (63)	36 (61)	43 (66)	65 (64)
Girls	43 (36)	43 (37)	23 (39)	22 (34)	37 (36)
Age (y)					
2-6 y	9.6 \pm 4.0	9.7 \pm 4.1	10.7 \pm 3.7	10.3 \pm 4.0	9.6 \pm 4.0
7-12 y	39 (33)	39 (34)	13 (22)	17 (27)	36 (35)
13-18 y	43 (36)	40 (34)	22 (37)	21 (33)	34 (33)
	36 (31)	37 (32)	24 (24)	25 (25)	32 (31)
Level of ID					
Moderate ID	53 (45)	52 (45)	38 (64) [#]	35 (56)	47 (46)
Severe ID	64 (54)	63 (54)	21 (36) [#]	30 (46)	54 (53)
Adaptive behavior (y)	1.9 \pm 1.4	1.9 \pm 1.4	2.6 \pm 1.8 [*]	2.4 \pm 1.8	2.0 \pm 1.5
Autism spectrum disorder	39 (33)	37 (32)	15 (25)	23 (35)	32 (31)
Down syndrome	30 (25)	29 (25)	17 (29)	15 (23)	24 (24)
Challenging behavior	77 (65)	76 (66)	35 (59)	39 (62)	64 (63)
Low social economic status	43 (36)	43 (37)	23 (39)	25 (38)	37 (36)
Non-Dutch origin	30 (25)	29 (25)	13 (22)	19 (29)	25 (25)
Bayley scale of infant development score	59.9 \pm 7.5	59.8 \pm 7.5	63.6 \pm 5.9 [*]	63.6 \pm 5.5 [*]	59.6 \pm 7.5
Steps per day	6551 \pm 2545 ¹	6630 \pm 2538 ²	7174 \pm 2147 ³	7569 \pm 2436 ⁴	6679 \pm 2508 ⁵

Notes. 6MWT = six-minute walk test; ID = intellectual disability; * significant different from total, assessed by t-test statistics; # significant different from total, assessed by Pearson Chi-square statistics; ¹ n=62; ² n=62; ³ n=27; ⁴ n=31; ⁵ n=59

Table A2. Average score on the physical fitness tests, and percentage participants scoring worse than reference values¹ for boys and girls separated.

	Boys				
	n	Score on test		Scoring worse than reference values ¹	
		M ± SD	95% CI	%	95% CI
BMI (kg/m ²)	75	18.7 ± 4.2	[17.7-19.6]		
zBMI	75	0.72 ± 1.31	[0.42-1.02]	36	[25-47]
Waist circumference (cm)	73	63.3 ± 15.3	[59.7-66.8]	23	[13-33]
Overarm throwing (cm)	36	166 ± 87	[136-195]	61	[44-78]
Stair climbing (n)	42	34 ± 17	[29-40]	83	[81-97]
Modified 6MWT (m)	65	351 ± 99	[326-376]	72	[61-83]
	Girls				
	n	Score on test		Scoring worse than reference values ¹	
		M ± SD	95% CI	%	95% CI
BMI (kg/m ²)	43	20.4 ± 4.6	[19.0-21.8]		
zBMI	43	0.98 ± 1.00	[0.67-1.29]	46	[31-62]
Waist circumference (cm)	43	69.1 ± 14.1	[64.7-73.4]	51	[36-67]
Overarm throwing (cm)	23	138 ± 77	[105-172]	87	[72-100]
Stair climbing (n)	21	29 ± 16	[22-36]	95	[89-100]
Modified 6MWT (m)	37	345 ± 98	[313-378]	81	[68-94]

Notes. BMI=body mass index; 6MWT = six minute walk test; ¹ Classified as >2SD (0-5y), >1SD (6-18y) for BMI [59], >1.3 SD for waist circumference [61], < P5 for overarm throwing and stair climbing [64], and <P10 for modified 6MWT [66].