



## Gestalt Perception in Children With Visual Impairments: Item-Specific Performance and Looking Behavior

Ymie J. van der Zee, Marlou J. G. Kooiker, Marisabel Talamante Ojeda & Johan J. M. Pel

To cite this article: Ymie J. van der Zee, Marlou J. G. Kooiker, Marisabel Talamante Ojeda & Johan J. M. Pel (2019) Gestalt Perception in Children With Visual Impairments: Item-Specific Performance and Looking Behavior, *Developmental Neuropsychology*, 44:3, 296-309, DOI: [10.1080/87565641.2019.1590836](https://doi.org/10.1080/87565641.2019.1590836)

To link to this article: <https://doi.org/10.1080/87565641.2019.1590836>



© 2019 The Author(s). Published with license by Taylor & Francis Group, LLC.



Published online: 17 Mar 2019.



Submit your article to this journal [↗](#)




Article views: 407



View Crossmark data [↗](#)

# Gestalt Perception in Children With Visual Impairments: Item-Specific Performance and Looking Behavior

Ymie J. van der Zee<sup>a</sup>, Marlou J. G. Kooiker<sup>b</sup>, Marisabel Talamante Ojeda<sup>b</sup>,  
and Johan J. M. Pel <sup>b</sup>

<sup>a</sup>Royal Dutch Visio, Centre of expertise for blind and partially sighted people, Rotterdam, The Netherlands;

<sup>b</sup>Vestibular and Ocular Motor Research Group, Department of Neuroscience, Erasmus MC, Rotterdam, the Netherlands

## ABSTRACT

Visual closure is the ability to visualize a complete whole when an incomplete picture is presented. The aim of the present study was to investigate the Kaufman Gestalt closure task in children with cerebral and ocular visual impairments. Looking behavior was assessed by an eye tracker system to quantify the number and duration of fixations. We found that children with visual impairments due to cerebral damage show weaker Gestalt perception and had different looking patterns than children with ocular or without visual impairments. Children with brain damage performed significantly worse on the animate items than the group without brain damage.

## ARTICLE HISTORY

Received 26 November 2018



Revised 21 February 2019

Accepted 24 February 2019

## Introduction

Neuropsychology aims to understand how behavior and cognition are influenced by brain functioning. Especially the effect of mental processes on behavior, such as attention, memory, and perception, are assessed in standardized neuropsychological tests. During such tests, the processing of visual information plays an important role because neuropsychological tests often consist of and employ visual representations. Dependent on the task at hand, a participant can be asked to discriminate details, to name forms and objects and to interpret relations within and between pictures. Thus, besides understanding a given instruction and being able to give a verbal or motor response, participants should be able to create a meaningful representation of the world from visual elements. An underlying premise is that simple visual elements are integrated into a whole, also known as a Gestalt, and that the given representations are visually organized through the same principles, or laws, of perceptual grouping (Han & Humphreys, 1999; Jäkel, Singh, Wichmann, & Herzog, 2016). Examples of Max Wertheimer's classical grouping principals are proximity and visual closure (Han & Humphreys, 1999; Wagemans et al., 2012). The proximity principle means that elements that are closer to each other are more likely to be perceptually grouped than more distant elements. The visual closure principle is the ability to visualize a complete whole when presented with incomplete information or a partial picture; thus, elements that form a closed figure tend to be grouped together (Wagemans et al., 2012). It was shown that visual closure is essential in the identification of incomplete drawings (Russell, Hendrickson, & VanEaton, 1988), of global letters (Pflugshaupt et al., 2016) and of hierarchically organized figures (Huberle & Karnath, 2012).

In adults without neurological impairment, the bilateral temporoparietal junction (TPJ), the bilateral anterior cingulate cortex (ACC), and the precuneus (PC), primarily on the left side, became activated in a recognition task (Huberle & Karnath, 2012). Presumably, perceptual grouping depends

**CONTACT** Johan J. M. Pel  [j.pel@erasmusmc.nl](mailto:j.pel@erasmusmc.nl)  Vestibular and ocular motor research group, Department of Neuroscience, Erasmus MC, PO Box 2040, Rotterdam, CA 3000, The Netherlands

Color versions of one or more of the figures in the article can be found online at [www.tandfonline.com/hdvn](http://www.tandfonline.com/hdvn).

© 2019 The Author(s). Published with license by Taylor & Francis Group, LLC.

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited, and is not altered, transformed, or built upon in any way.

on the integrity of a large brain network and brain damage may result in poorer performance. Indeed, damage in the right occipital lobe (Russell et al., 1988) and the medial parietal lobe, i.e., the precuneus (PC) (Pflugshaupt et al., 2016) resulted in weaker recognition performance. In adults, Gestalt perception, visual illusions or hierarchical letters are used to test aspects of the principle grouping. In children, the subtest Gestalt Closure of the Kaufman Assessment Battery for Children second edition (KABC-II) is frequently used. This test is an object recognition task with incomplete representations and is a subtest of the Simultaneous Processing scale. Despite the importance of these aspects of visual information processing, not many studies have addressed Gestalt perception performance in visually impaired children. A population-based cohort study of very preterm children showed that at an age of 5 years, children born premature had lower scores than full-term controls on multiple scales, including the simultaneous processing scale (Larroque et al., 2008). Exclusion of participants with cerebral palsy (9%), an indication of brain damage, visual impairments (1% of the population had low vision, a visual acuity <3/10) and hearing impairment (<1%) had no influence on the results (Larroque et al., 2008). These data suggested that prematurity combined with brain damage did not explain differences in performances. Yet, the number of children with impairments was too small to draw definite conclusions.

As stated, a perceptual grouping of simple visual elements (e.g., lines and arcs) into simple forms and shapes helps to create meaningful representations of the world. In this process, visual attention plays a very important role. The frontoparietal network is essential for both bottom-up attention (externally driven and involving stimuli that are salient because of their inherent properties relative to the background) and top-down attention (internally driven on prior knowledge, willful plans, and current goals) (Katsuki & Constantinidis, 2014). Based on event-related potential patterns, it was shown that visual objects organized according to Gestalt principles were prioritized over competing non-Gestalt stimuli in a bottom-up and automatic fashion (Marini & Marzi, 2016). However, these electrophysiological responses (~150 ms after stimulus onset) only relate to the first steps of perceptual grouping of simple visual elements and are not directly associated with the process of perceptual grouping that results in perception. Alternatively, visual attention can be quantified from infancy onwards using visually-guided eye movement responses. Differences in interactions between eye movement responses and salience in target areas have been found in children at risk of cerebral and ocular visual impairments compared to typically developing children (Kooiker, van der Steen, & Pel, 2016). Specifically, in children at risk of cerebral visual impairments (CVI), higher reaction times to large and high contrast stimuli were found in combination with poorer fixation stability. These children might have problems with bottom-up regulated visual attention orienting, not only in visually-guided response tasks but also in visual recognition tasks, like Gestalt Closure.

The aim of the present study was to investigate the role of visual attention orienting in the Kaufman Gestalt closure task in children with cerebral and ocular visual impairments. Digitized Gestalt items were shown one-by-one on a monitor with an integrated eye tracker. Each participant was asked to name the complete image that each item represented. For analysis purposes, we grouped the items in animate (i.e., humans and animals) and inanimate objects (i.e., everything else). For each item, eye movement responses were collected to calculate the number of gaze shifts and fixation durations. We hypothesized that children with visual impairments score lower on Gestalt Closure performance than age-matched American controls and that gaze fixation duration on visual elements of the Gestalt (area of interest) in children at risk of CVI is longer than in children with OVI.

## Methods

### Participants

For this study, we used the database of a longitudinal eye-tracking study. Data collection took place between 2012 and 2014 in a cohort of children with visual impairments (Kooiker, Verbunt, van der Steen, & Pel, 2019). The database consisted of 122 clients of Royal Dutch Visio,

a rehabilitation center for visually impaired and blind people. Children were recruited from different regions across The Netherlands: Amsterdam, Grave/Nijmegen, Haren, Huizen, Breda and Rotterdam. During the eye tracking assessments in this cohort, Gestalt Closure was assessed as an additional task. The experimental procedures were approved by the Medical Ethical Committee of the Erasmus University Medical Centre, Rotterdam, The Netherlands (MEC 2012–097) and adhered to the tenets of the declaration of Helsinki (2013). The parents of all children gave written informed consent.

Of the total cohort of 122 patients, 72 patients underwent the Gestalt Closure test. The remaining 51 patients had either a developmental level below 3 years old, or the task was not performed due to a time limitation. The chronological age of the selected patients (42 male, 30 female) ranged from 4y6m to 13y8m ( $M = 9y8m$ ,  $SD = 2y5m$ ). Four patients had syndromes: one CHARGE-syndrome, one Down syndrome, one Noonan syndrome, one Sotos syndrome. Forty-four (60%) had nystagmus as the main diagnosis or as a symptom of another diagnosis, like albinism. Based on the information in the medical records about ocular abnormalities, brain damage, the presence of paresis and visual diagnosis, the children were divided into three different groups: ocular visual impairments (OVI), ocular and cerebral visual impairment (OCVI), or cerebral visual impairments (CVI).

### **Group 1: ocular visual impairment (OVI)**

Of the 72 patients, 38 (23 males, 15 females) were included in group 1. Twenty-three patients had iris or retinal abnormalities: iris coloboma ( $n = 1$ ); aniridia combined with lens luxation and glaucoma ( $n = 1$ ); uveitis and retinopathy of prematurity (ROP;  $n = 1$ ); ocular or oculocutaneous albinism ( $n = 10$ ); cone/rod dysfunction ( $n = 6$ ); CMV retinitis ( $n = 1$ ); retinitis pigmentosa ( $n = 1$ ); retinoschisis ( $n = 1$ ). Ten patients had lens abnormalities: hypermetropia ( $n = 2$ ); high myopia ( $n = 2$ ); cataract ( $n = 6$ ). Four patients had congenital nystagmus and the cause of visual problems was unknown in two patients. A total of 23 patients had nystagmus, 12 patients had strabismus and 21 wore glasses. Visual acuity ranged from 0.05 to 0.80 (Snellen equivalent). Thirty-one patients had a visual acuity equal to or lower than 0.3 (cut-off for low vision), of whom eight could be considered legally blind (visual acuity  $\leq 0.1$ ). Of the remaining seven patients, six had a subnormal visual acuity ( $< 0.8$ ) and one had a normal visual acuity ( $\geq 0.8$ ). Behavioral problems were mentioned in nine patients: three had no diagnosis, one had an autism spectrum disorder (ASD), two had attention deficit hyperactivity disorder (ADHD), one had ASD and ADHD, one had ADHD and oppositional defiant disorder (ODD), one had an attachment disorder. In 11 patients, no estimation of the cognitive level was present. Of the remaining patients, three had a mild cognitive impairment ( $50 < TIQ < 70$ ), four functioned at a borderline cognitive level ( $70 < TIQ < 89$ ), six at a below average level ( $70 < TIQ < 89$ ) and seven at an average level ( $TIQ \geq 90$ ). The chronological age of the OVI group ranged from 4y8m to 13y7m ( $M = 9y9m$ ,  $SD = 2y5m$ ).

### **Group 2: ocular and cerebral visual impairment (OCVI)**

Fourteen patients (7 males, 7 females) were included in group 2. Eight had ocular abnormalities and brain damage. Six had no brain damage but were included based on the clinical signs and diagnosis of CVI, in addition to the presence of ocular abnormalities. In Table 1 the etiological categories for the brain damage and the distribution of cognitive level of the CVI and OCVI group are presented. Present ocular abnormalities were: nystagmus ( $n = 7$ ); myopia ( $n = 1$ ); ocular albinism ( $n = 2$ ); optic nerve atrophy ( $n = 1$ ); cataract ( $n = 1$ ) and retinitis pigmentosa ( $n = 2$ ). Seven children had strabismus and 11 wore glasses. Visual acuity ranged from 0.10 to 0.60. Ten were visually impaired (visual acuity:  $\leq 0.3$ ), one could be considered legally blind (visual acuity:  $\leq 0.1$ ). The other four had a subnormal visual acuity. The (estimated) level of cognitive functioning ranged from a mild intellectual disability ( $50 < TIQ < 70$ ) to an average level ( $90 < TIQ < 110$ ). Behavioral problems

**Table 1.** Etiology of brain damage and estimated cognitive levels in children in the groups OCVI ( $n = 14$ ) and CVI ( $n = 20$ ).

	OCVI	CVI
<b>Etiology of brain damage</b>	<b><i>n</i> (%)</b>	<b><i>n</i> (%)</b>
No imaging data	6 (43)	5 (24)
Perinatal	8 (57)	12 (57)
Asphyxia	3 (21)	3 (15)
Hypoxic-ischemic encephalopathy (HIE) <sup>a</sup>	2 (14)	6 (30)
Dysgenesis/agenesis	2 (14)	2 (10)
Postnatal brain damage		
Acquired brain damage <sup>b</sup>	-	4 (20)
Peri- and/or postnatal		
Hydrocephalus	-	1 (5)
Mild atrophy	1 (7)	-
<b>Estimated cognitive level</b>		
Average (TIQ $\geq 90$ )	4 (29)	4 (20)
Below average (70 < TIQ < 89)	2 (14)	6 (30)
Borderline (70 < TIQ < 89)	4 (29)	3 (15)
Mild impairment (50 < TIQ < 70)	2 (14)	4 (20)
Moderate impairment (TIQ < 50)	-	2 (10)
No data	2 (14)	1 (5)

<sup>a</sup> Hypoxic-ischemic encephalopathy (HIE) includes different categories for perinatal brain damage, i.e., Periventricular leukomalacia (PVL), Intraventricular hemorrhage (IVH), Cerebral Vascular Accident (CVA) or combinations of these categories; <sup>b</sup> Acquired brain damage consists of different categories, i.e., CVA, meningitis, and tumor.

were mentioned in four patients: one had no diagnosis, three had ADHD. The chronological age of the OCVI group ranged from 5y11m to 13y7m ( $M = 9y6m$ ,  $SD = 2y7m$ ).

### Group 3: cerebral visual impairment (CVI)

Twenty patients (12 males, 8 females) were included in group 3. Some patients had multiple causes. Two patients had no objectified brain damage but were included based on the clinical diagnosis of CVI, one patient had a developmental coordination disorder (DCD) and in two patients a paresis was present, suggesting the presence of some type of brain dysfunction or damage. Of the patients with dysgenesis of the brain, one patient had septo-optic dysplasia, a condition characterized by the underdevelopment of the optic nerve and the absence of the septum pellucidum (midbrain structure); one had Arnold-Chiari malformation, a condition of characterized by the malformation of the cerebellum and/or brainstem. One patient had iris coloboma, as a symptom of the CHARGE syndrome combined with hydrocephalus, and one patient had a cone dysfunction.

Eight patients had nystagmus, 14 had strabismus and eight wore glasses. Visual acuity ranged from 0.05 to 1.00. Ten were visually impaired (visual acuity  $\leq 0.3$ ), of whom three could be considered legally blind (visual acuity  $\leq 0.1$ ). Of the remaining 10 patients, 6 had a subnormal visual acuity (visual acuity < 0.8). Behavioral problems were mentioned in five patients: four had no diagnosis, one of them had ASD. The (estimated) level of cognitive functioning ranged from a moderate intellectual disability (TIQ < 50) to an average level (90 < TIQ < 110) and cognitive level was unknown in one patient. The chronological age of the CVI group ranged from 4y6m to 13y10m ( $M = 9y8m$ ,  $SD = 2y5m$ ).

The mean age and the distribution of visual acuity levels of the three different groups did not differ significantly. The distributions of the estimated cognitive level did significantly differ ( $\chi^2(2) = 11.29$ ,  $p < .01$ ) between groups: post-hoc analyses showed a higher mean cognitive level of the OVI group than of the two other groups.

## Procedures

The subtest Gestalt Closure of the Kaufman Assessment Battery for children second edition (KABC-II) consisted of one example item and 37 tests items of increasing difficulty (Kaufman & Kaufman, 2004). The items contained incomplete black figures on a white background and represent 12 animate objects (7 animals including a dinosaur and 5 human faces or activities), and 25 inanimate objects. The normal clinical procedure would be to start with the example item followed by the age-appropriate items. Here, we showed all items on a 24-in. monitor with an integrated infrared eye-tracking system (Tobii T60-XL, Tobii Corporation, Sweden). In-between each item, a neutral display was shown with a fixation point to let the child refocus on the center of the monitor. The system measured gaze position of each eye separately using cornea reflection and compensated for free head movements. Each child sat at approximately 60 cm distance from the monitor. The visual angle towards the monitor was approximately  $30 \times 24$  degrees ( $1280 \times 1024$  pixels). Eye movement responses were recorded with a sampling rate of 60 Hz, and the latency of the system was approximately 25 ms. The experiments were conducted in a quiet room with ambient light conditions. Prior to starting the Gestalt Closure test a standardized five-point calibration procedure of both eyes was performed. Item presentation time was dependent on the child's verbal response, i.e., the examiner pushed the spacebar after the first answer to each item and thereby ended the item presentation. The examiner also noted whether the answer was correct. After four consecutive errors, the assessment was stopped. All data were stored off-line.

## Data analysis

### Performance data

Firstly, we scored items the standard way, i.e., recognition was either correct or incorrect. The items after discontinuation were assumed to be too difficult and therefore scored as incorrect. Raw scores, i.e., the total correct items up to four consecutive incorrect answers, were noted, and the American norm values for the age group 4–14 years old were used to classify the performances (Kaufman & Kaufman, 2004), based on scaled scores ( $M = 10$ ;  $SD = 3$ ): scaled scores 1–5 (expected: 9%) = weak; 6–7 = below average (expected: 16%); 8–11 = average (expected: 50%); 12–13 above average (expected: 16%); 14–19 = superior (expected: 9%). Secondly, a simple item analysis was done by calculating the percentage of patients that correctly recognized each item. This way, we were able to check whether items increased in difficulty as reported in the original population. For the analysis of the subgroups, i.e., the animate and inanimate items, we calculated the percentage of correct items with respect to the total number of items presented in that subgroup.

### Eye tracking data

Gaze data were analyzed off-line using Tobii Studio 3.2 analysis software (Tobii, Danderyd, Stockholm, Sweden). Patients with at least 40% gaze data measured during the assessment were selected for further analysis to ensure the reliability of the eye movement responses. For each stimulus, an area of interest (AOI) was defined that corresponded with the size and the location of the Gestalt item. The eye movement patterns relative to these AOIs were obtained to calculate 1) the mean and total time spent within the AOI and 2) the number of times gaze went in and out of the AOI. A lower mean and/or total time within the AOI might be the result of efficient information processing, i.e., fast recognition, or caused by the presence of nystagmus, attentional problems or a lack of effort.

### Statistical analysis

We used IBM SPSS Statistics version 20 for statistical analyses. Non-parametric tests were used, since data were not normally distributed, even not after data transformation. To compare groups with



ordinal (number of items shown; scaled scores) and skewed data (percentage correct of presented items), Kruskal Wallis and Mann–Whitney U-tests were used. To test for differences in performance while controlling for differences in cognitive level, we used the partial Spearman correlation test. Binomial tests were used to compare the expected percentage of weak, average and superior performers with the actual percentages found in the patient groups. Finally, an explorative correlation analysis was done between weak overall performance, visual acuity, nystagmus, and eye-tracker results.  $p$ -values  $\leq .05$  were considered significant.

## Results

The collection of the performance scores of one patient with albinism in the group OVI ( $n_{OVI} = 37$ , 22 males, 15 females) failed, resulting in a total group of seventy-one tested patients ( $M_{age} = 9y9m$ ,  $SD = 2y5m$ ; 41 males, 30 females).

### Overall performance

#### Scaled scores

Analysis of the scaled scores showed that the distributions of scores significantly differed across groups ( $\chi^2(2) = 8.65$ ,  $p = .01$ ). The group CVI performed significantly worse ( $Mdn = 2$ ) than the group OVI ( $Mdn = 9$ ;  $U = 197.5$ ,  $p < .01$ ). Compared to the normal distribution of the scaled scores in the norm population, half of the patients in the group CVI performed extremely weak. A scaled score of 5 is expected in 9% of the population, yet, this scaled score of 5 or lower was found in 9/37 (24%) of the OVI patients, 5/14 (36%) of the OCVI patients and 12/20 (60%) CVI patients ( $p$ 's  $< .01$ ). Despite these weak performances, still 12/37 (32%) of the OVI patients (7/14) (50% of the OCVI patients and only 12/37 (32%) of the CVI patients performed on an average level ( $p = .05$  and  $p < .01$ ). The percentage of patients that performed at a superior level was not higher than expected. The Spearman rho correlations between the patient groups and 1) the scaled item scores ( $r_s = -.31$ ) and 2) the proportion weak performers ( $r_s = .32$ ) were significant ( $ps = .02$ ). Controlling for the cognitive level, group differences in scaled scores were only near significant ( $r_{s,cognitive\ level} = -.25$ ,  $p = .07$ ), but group differences in the proportion of weak performers remained significant ( $r_{s,cognitive\ level} = .26$ ,  $p = .05$ ). This suggests that the performance of the group CVI remains significantly weaker after correction for differences in cognitive level.

#### Item scores

The item scores showed a monotone decrease with increasing number of presented items, suggesting that test difficulty increased per presented item (Figure 1). Notably, the percentage correct animate items, especially the animal items, deviated from this general pattern. We found that the animal items were more difficult to recognize or to name than the inanimate items. The most difficult items were the dinosaur (item 7; 44% correct) and the elephant (item 18; 39% correct). Analysis of the percentage correct answers of the presented items showed that significant differences in performances were indeed present ( $\chi^2(2) = 7.28$ ,  $p = .03$ ), see also Figure 2. The group CVI ( $Mdn = 59.94$ ) performed significantly worse than the group OVI ( $Mdn = 73.33$ ;  $U = 215.0$ ,  $p = .01$ ). The difference between the groups CVI and OCVI ( $Mdn = 67.95$ ) and OCVI and OVI were not significant. Significant negative correlations ( $r_s = -.34$ ,  $p = .01$ ;  $r_{s,cognitive\ level} = -.29$ ,  $p = .03$ ) were found between the percentage correctly recognized items and the patient groups, indicating that cognitive level does not explain the group difference found.

#### Animate items

Analysis of the number of animate items presented showed that the number of presented animal items differed across groups ( $\chi^2(2) = 6.82$ ,  $p = .03$ ), suggesting differences in performances. In the




Figure 1. Percentage of correct answers per item-group: inanimate objects and animate objects, human (activity) and animals.





Figure 2. Distribution of percentage correct answers given the number presented items in the different patient groups.

groups OVI and the OCVI, only one patient was presented less than seven items (the total number of animal items) (OVI: 2 items; OCVI: 5 items). In the group CVI, seven patients were presented less than seven items; less than three items were presented to five patients, less than five items to one patient and less than six items were presented to three patients. The number of presented items was significantly different between the groups OVI and the CVI ( $U = 290.0$ ,  $p < .01$ ). Analysis of the percentage correct answers of the presented items showed that significant differences in performances were indeed present ( $\chi^2(2) = 13.45$ ,  $p < .01$ ), see also Figure 3. The groups CVI ( $Mdn = 50.00$ ) and OCVI ( $Mdn = 64.29$ ) performed significantly worse than the group OVI ( $Mdn = 85.71$ ;  $U = 171.5$ ,  $p < .01$  and  $U = 158.00$ ,  $p = .03$ ). The difference between the groups CVI and OCVI was not significant. A significant negative correlation ( $r_s = -.43$ ,  $p < .01$ ) was found between the percentage correctly recognized animals and the patient groups. This indicates that children with brain damage tend to perform worse than children without brain damage (OVI) on





**Figure 3.** Distribution of percentage correct answers given the number presented animal items (max 7 items) in the different patient groups.

animal items. Importantly, this correlation remained significant when controlling for differences in cognitive level ( $r_{s,\text{cognitive level}} = -.38, p < .01$ ).

### Inanimate items

To test whether the same pattern of group differences was also present in recognizing the inanimate items, we first selected seven inanimate items with approximately the same performance score as the seven animal items. We selected item 2, 10, 16, 19, 20, 23, and 24. The median difference in performance between these two item sets was  $-1\%$  (range  $-6\%$  to  $+4\%$ ). Thus, compared to the animate items, these selected inanimate items were a bit easier to recognize. Again, the number of presented inanimate items varied from 1 to 7 in the groups OVI and CVI and 2 to 7 in the group OCVI. The distribution of the number of presented inanimate items significantly differed across the groups ( $\chi^2(2) = 11.45, p < .01$ ). In the group CVI, a significantly smaller number of items was presented ( $Mdn = 5$ ) than in the group OVI ( $Mdn = 7; U = 212.5, p < .01$ ). The percentage correct answers, however, to the inanimate items were not significantly different between the groups ( $Mdn_{\text{OVI}} = 85.71, Mdn_{\text{OCVI}} = 71.43, Mdn_{\text{CVI}} = 60.00; p = .13$ ). Correlations between performance and groups were not significant ( $r_s = -.21, r_{s,\text{cognitive level}} = -.19$ ). Thus, children with CVI tend to perform in a similar way as children with OCVI and OVI on inanimate items.

### Gaze patterns

#### Group characteristics

Of the 72 patients, forty-five patients had at least 40% gaze data. The group OVI consisted of 20 patients (12 males, 8 females). The chronological age ranged from 5y0m to 13y7m ( $M = 9y10m, SD = 2y7m$ ). Thirteen patients had iris or retinal abnormalities: iris coloboma ( $n = 1$ ); ocular or oculocutaneous albinism ( $n = 9$ ); cone/rod dysfunction ( $n = 2$ ); retinitis pigmentosa ( $n = 1$ ). Three patients had lens abnormalities: high myopia ( $n = 1$ ), Stickler Syndrome; cataract ( $n = 2$ ). Three patients had congenital nystagmus and the cause of visual problems was unknown in one patient. The group included four patients with a weak performance on the Gestalt test.

The group OCVI consisted of 11 patients (5 males, 6 females). The chronological age ranged from 6y5m to 13y7m ( $M = 9y7m, SD = 2y8m$ ). Etiology of brain damage was: asphyxia ( $n = 3$ ), dysgenesis

( $n = 1$ ), periventricular leukomalacia ( $n = 1$ ), intraventricular hemorrhage ( $n = 1$ ), atrophy ( $n = 1$ ). Imaging data was not available for four patients. Present ocular abnormalities were: nystagmus ( $n = 5$ ); ocular albinism ( $n = 2$ ); optic nerve atrophy ( $n = 1$ ); and retinitis pigmentosa ( $n = 2$ ), unknown ( $n = 1$ ). The group included three patients with a weak performance on the Gestalt test.

The group CVI consisted of 14 patients (10 males, 4 females). The chronological age ranged from 4y6m to 13y10m ( $M = 10y2m$ ,  $SD = 2y6m$ ). Etiology of brain damage was: asphyxia ( $n = 3$ ), Arnold-Chiari malformation ( $n = 1$ ), periventricular leukomalacia ( $n = 2$ ), cerebral vascular incident ( $n = 3$ ), tumor ( $n = 1$ ). No imaging data was present in four patients. This group included eight patients with a weak performance on the Gestalt test.

The mean age and the estimated cognitive level of the three different groups did not differ significantly, see Table 2. Group differences were present in the distributions of the visual acuity ( $\chi^2(2) = 7.27$ ,  $p = .03$ ) and nystagmus ( $\chi^2(2) = 12.22$ ,  $p < .01$ ). Posthoc analyses showed a higher visual acuity and fewer patients with nystagmus in the group CVI than of the two other groups ( $ps \leq .02$ ). Visual acuity in the groups OVI and OCVI were considered comparable.

### Item characteristics

In one CVI patient, no eye-tracking data was available during the presentation of the seven animal and seven selected inanimate items. In the remaining group of 44 patients, eye-tracking data during the presentation of the animal items were present for at least four items in 20/20 OVI patient, 11/11 OCVI patients, and 11/13 CVI patients. To control for individual differences in the number of items, we used mean individual outcomes for group comparisons. Overall, no significant group differences on the number of visits and the median visit duration of each visit within the area of interest (AOI) were found, but we did find a significant difference in the total visit duration ( $\chi^2(2) = 6.38$ ,  $p = .04$ ), see Table 3. This indicates that children with CVI tend to need more time to analyze the Gestalt to formulate an answer. Analysis of individual animal items showed that children with CVI tend to be a bit slower on 6 of 7 items, but differences were not significant. Of the seven selected inanimate items, eye-tracking data of at least four items were available in 20/20 OVI patients, 10/11 OCVI patients, and 12/13 CVI patients. Here, no significant group differences were found, see Table 4.

Finally, explorative correlation analyses were done between weak overall performance, visual acuity, nystagmus and eye tracker results of the animal items within the patient groups. The correlations between weak overall performance, visual acuity, and nystagmus were not significant, except the correlation between weak overall performance and visual acuity in the group

**Table 2.** Group characteristics of children with at least 40% gaze data in the groups OVI ( $n = 20$ ), OCVI ( $n = 11$ ), and CVI ( $n = 14$ ).

Characteristic	OVI <i>n</i> (%)	OCVI <i>n</i> (%)	CVI <i>n</i> (%)	<i>p</i> -value
Glasses	9 (45)	8 (73)	6 (43)	
Strabismus	7 (35)	4 (36)	10 (71)	.09, <i>ns</i>
Nystagmus	15 (75)	11 (100)	5 (36)	<.01
Visual acuity				.03
$\geq 0.8$	1 (5)		4 (29)	
0.5–0.8	4 (20)		3 (21)	
0.4–0.5		3 (27)	3 (21)	
$\leq 0.3$	15 (75)	8 (73)	4 (29)	
Cognitive level				.07, <i>ns</i>
Average (TIQ $\geq 90$ )	7 (35)	4 (36)	1 (7)	
Below average (70 < TIQ < 89)	2 (10)	1 (9)	4 (29)	
Borderline (70 < TIQ < 89)	3 (15)	3 (27)	3 (21)	
Mild impairment (50 < TIQ < 70)	2 (10)	1 (9)	3 (21)	
Moderate impairment (TIQ < 50)			2 (14)	
No data	6 (30)	2 (18)	1 (7)	

*p*-Values < .05 indicate statistically significant group differences.

**Table 3.** Outcomes of eye-tracking data (number of visits of the area of interest, median visit duration, and total visit duration) on animal items in the groups OVI ( $n = 20$ ), OCVI ( $n = 11$ ), and CVI ( $n = 13$ ).

	OVI	OCVI	CVI	<i>p</i> -value
Mean				
# visits AOI	7 (1–33)	10 (3–15)	6 (2–28)	<i>ns</i>
Median visit duration	0.7 (0.1–3.4)	0.7 (0.1–3.3)	1.3 (0.1–4.7)	<i>ns</i>
Total visit duration	3.7 (1.0–6.7)	3.8 (0.5–12.0)	8.4 (0.2–14.4)	.04
Butterfly				
<i>n</i>	20	11	11	
# visits AOI	4 (1–15)	2 (1–12)	3 (1–13)	<i>ns</i>
Median visit duration	0.4 (0.1–3.5)	1.3 (0.1–3.6)	1.2 (0.2–6.1)	<i>ns</i>
Total visit duration	2.4 (0.1–3.5)	2.2 (0.1–5.2)	2.7 (0.8–13.1)	<i>ns</i>
Dog				
<i>n</i>	20	11	13	
# visits AOI	3 (1–19)	2 (1–11)	2 (1–22)	<i>ns</i>
Median visit duration	1.0 (0.1–3.5)	0.6 (0.1–3.4)	1.7 (0.1–5.6)	<i>ns</i>
Total visit duration	2.6 (0.7–7.0)	2.4 (0.1–4.4)	3.0 (0.1–20.7)	<i>ns</i>
Dinosaur				
<i>n</i>	20	11	12	
# visits AOI	8 (1–47)	7 (1–25)	5 (1–63)	<i>ns</i>
Median visit duration	0.7 (0.1–6.2)	0.8 (0.1–11.8)	2.4 (0.3–7.2)	.08
Total visit duration	5.1 (1.8–26.6)	5.5 (0.6–11.8)	6.1 (1.6–30.2)	<i>ns</i>
Fish				
<i>n</i>	20	10	10	
# visits AOI	3.0 (1–90)	5.5 (1–24)	7.0 (1–44)	<i>ns</i>
Median visit duration	0.7 (0.1–4.4)	1.0 (0.1–3.2)	0.80 (0.3–6.4)	<i>ns</i>
Total visit duration	2.8 (0.1–7.6)	2.9 (0.1–22.5)	6.35 (1.18–19.96)	<i>ns</i>
Turtle				
<i>n</i>	20	10	12	
# visits AOI	10 (1–13)	10 (4–34)	8.5 (1–27)	<i>ns</i>
Median visit duration	0.4 (0.1–4.0)	0.3 (0.1–6.0)	0.8 (0.1–7.0)	<i>ns</i>
Total visit duration	3.2 (0.4–17.6)	5.6 (1.4–24.2)	5.3 (0.5–29.0)	.10
Pig				
<i>n</i>	19	10	10	
# visits AOI	7.0 (1–29)	4.0 (1–17)	4.0 (1–12)	<i>ns</i>
Median visit duration	0.3 (0.1–3.8)	0.6 (0.1–4.2)	1.6 (0.3–9.0)	.03
Total visit duration	2.6 (0.1–7.4)	2.9 (0.1–23.0)	4.0 (1.9–18.0)	.12
Elephant				
<i>n</i>	20	10	9	
# visits AOI	7 (2–44)	15 (2–35)	5 (1–94)	<i>ns</i>
Median visit duration	0.4 (0.1–1.1)	0.3 (0.1–1.2)	0.6 (0.2–3.6)	.03
Total visit duration	3.4 (0.1–6.7)	5.0 (0.1–16.3)	5.6 (1.0–43.0)	<i>ns</i>

AOI = area of interest. *p* Values indicate statistically significant differences in eye-tracking parameters between patient groups.

**Table 4.** Outcomes of eye-tracking data (number of visits of the area of interest, median visit duration, and total visit duration) of the seven selected object items in the groups OVI ( $n = 20$ ), OCVI ( $n = 11$ ), and CVI ( $n = 12$ ).

	OVI	OCVI	CVI	<i>p</i> Value
Mean				
# visits AOI	6 (1–34)	5 (2–13)	6 (2–19)	<i>ns</i>
Median visit duration	1.0 (0.2–5.0)	1.3 (0.1–2.6)	1.3 (0.1–4.2)	<i>ns</i>
Total visit duration	3.0 (0.8–7.8)	4.2 (0.6–5.9)	4.7 (1.0–14.8)	<i>ns</i>

AOI = area of interest. *p* Values indicate statistically significant differences in eye-tracking parameters between patient groups.

CVI ( $r_s = -.63$ ,  $p = .02$ ). This suggests that weak performers within the group CVI have lower visual acuities. Within the group OVI we found that the weak performers tend to stay shorter within the AOI ( $r_s = -.52$ ,  $p = .02$ ), children with lower visual acuity visit the AOI more often ( $r_s = -.59$ ,  $p < .01$ ), children with higher visual acuity seem to stay longer within the AOI ( $r_s = .43$ ,  $p = .06$ ) and children with nystagmus need in total more time to analyze and respond ( $r_s = .47$ ,  $p = .03$ ). Within the group CVI we found that weak performers tend to visit the AOI more often

( $r_s = .63$ ,  $p = .03$ ), and need in total more time to analyze and give a response ( $r_s = .72$ ,  $p < .01$ ). This indicates that the weak score of these children was not the result of a lack of effort.

## Discussion

Neuropsychological assessment in children often includes subtests that examine the ability to integrate visual elements into a (meaningful) whole, i.e., perceive Gestalts. This is the first study to address Gestalt perception in visually impaired children. This study about performances and looking behavior on a Gestalt Closure test (KABC-II) in groups of children with visual impairments, i.e., children with brain damage or clinical signs of cerebral visual impairment (CVI), children with ocular abnormalities and brain damage (OCVI) and children ocular abnormalities (OVI), shows that children with visual impairments more often performed weakly than age-matched controls of the American norm population, and that children with brain damage performed worse than children without brain damage. The proportion of weak performers in the groups CVI (60%) and OCVI (36%) was significantly higher than that in the group OVI (24%). Differences remained significant even after controlling for differences in cognitive level. Eye movement patterns revealed that weak performance on animate items in the CVI group was associated with longer fixations and more re-fixations.

Literature on Gestalt performance in children, including fragmented picture recognition, hierarchical figures, and embedded figures, covers many conditions, for example, ASD (Bölte, Holtmann, Poustka, Scheurich, & Schmidt, 2007), schizophrenia (Uhlhaas, Phillips, Mitchell, & Silverstein, 2006; van de Ven, Rotarska-Jagiela, Oertel-Knöchel, & Linden, 2017) intellectual disability syndromes (Hodapp et al., 1992). These reports may provide insights into mechanisms and networks underlying our current results. For example, our finding that children with a (partly) cerebral cause for their visual impairment more often perform weakly on Gestalt Closure is related to findings in children with brain damage in general, who were found to have worse visuomotor Gestalt performance (i.e. Bender Gestalt test) than children without brain damage (Koppitz, 1962). More specific, PVL is an important underlying cause of CVI in children, and children with PVL showed poor results on a Closure subtest of a visual perception battery (Fazzi et al., 2004).

Neuroimaging studies showed that performance in visual closure tasks is associated with activity in object-related processing areas, including the lateral occipital cortex and inferior temporal areas (Doniger et al., 2000; Ploran et al., 2007). For example, patients' impaired perceptual closure performance was associated with a diminished closure-related ERP negativity, of which the neural source has been attributed to visual association areas (Doniger, Foxe, Murray, Higgins, & Javitt, 2002). Furthermore, two studies on Gestalt perception in schizophrenia patients showed a relation of reduced neural connectivity in visual cortical areas and impaired perceptual closure. Although no studies about Gestalt performance in visually impaired children could be found, one study showed that early visual deprivation in patients with congenital cataract aged 9–23 years was related to impairments in holistic (face) processing (Le Grand, Mondloch, Maurer, & Brent, 2004). Our current results in a more general population of children with visual impairments confirm that in children who have experienced some sort of visual deprivation early in life, holistic visual processing, in general, seems to be affected.

Notably, children with brain damage (groups CVI and OCVI) performed significantly worse on the animate items than the group without brain damage (OVI). Differences between groups on the inanimate items were smaller and non-significant, suggesting that the overall difference can mainly be attributed to the weak performance on the animal items. The gaze data collected with the eye tracker reveals that the CVI children with weak performances fixate these Gestalt items longer and even re-fixate them more often. This suggests that this difference in performance may not be attributed to a lack of effort. Further analysis on the seven animal items, items earlier in the test than the seven selected object items but with a comparable difficulty level, shows that weak performers within the group OVI spend less time within the area of interest (AOI) per visit.

There was no relation with visual acuity or nystagmus, which suggests there is another reason why they perform weakly. Maybe children with OVI have learned other strategies, for example, to guess more often to save time and energy or to perform as fast as children without a visual impairment. In order to study this hypothesis, a control group should be added, and reaction times and performances should be compared.

It has been shown that processing of either animate or inanimate pictures can be selectively impaired in patients with damage to anterior visual areas (Caramazza & Shelton, 1998; Hillis & Caramazza, 1991; Warrington & Shallice, 1984). More recently, fMRI studies found clusters of voxel population vectors that were associated with animate and inanimate categories located anterior to retinotopic visual areas (Kriegeskorte et al., 2008; Naselaris, Stansbury, & Gallant, 2012). However, in these studies, images of hundreds of well-defined objects were presented to the subjects. In the present study, visual processing of each item involved perceptually grouping of closely projected elements to visualize a complete object, even when presented incomplete information. To our surprise, we found a significantly worse performance for naming the animate items in children with confirmed or suspected brain damage. We could not find any support for our finding in literature. At this point, we do not know if the differences we found between animate and inanimate object recognition in children with cerebral visual impairments are related to possible damage to the associated processing areas in the brain, e.g., the occipital face area (OFA) and the extrastriate body area (EBA) (Naselaris et al., 2012). The available imaging data lack the detailed information that is required to obtain insight in these specific regions.

This study has some limitations that need to be addressed. Firstly, the Kaufman Gestalt closure task contains a fixed number of 37 items. We applied the classic stopping rule, i.e., we stopped testing when a subject gave four consecutive incorrect answers. As a result, the total number of tested items was different between the subjects. For the between-group comparisons per item, we were able to control for this by calculating the correct items with respect to the total number of items presented within each subject. The comparison of the performance scores between the animate and inanimate items, however, inevitably led to two different subgroups of patients. Secondly, in a majority of the patients, only the global cognitive functioning was scored using total IQ. We are aware of the fact that more exact data on verbal and non-verbal IQ is preferable to control for differences in developmental level between groups. Thirdly, the performance of the children in the present study was compared with the American norm group. We did not test a Dutch age-matched control group, to investigate any cultural differences. Some of the items were typical American illustrations, such as a land map of the US. These items were not judged too strictly – and were not selected for further item analysis. Fourthly, we did not check after the Gestalt closure test whether children could name standard drawings of the same objects. Especially for the animated items, this would have been a valuable control condition. Lastly, the size of the area of interest dependent on the size of the Gestalt closure item. As a result, some AOI's were larger than others but not more than ~20%. Presumably, this had some influence on the number of re-fixations, especially for the small Gestalt closure items. In addition, we attempted to investigate whether a subject fixated the different elements within the Gestalt closure item to create an overview or that this was obtained with one central fixation. Due to the variability in gaze behavior, we were not able to find a consistent pattern. In a future study, we would like to address this question using a fixed set of Gestalt closure items of equal sizes and a higher resolution eye tracker. Lastly, the quality of each eye tracking study depends on the quality of the recorded gaze patterns. In our group, 45 of the 72 patients (63%) had at least 40% valid gaze data. In the present study, the investigator was able to keep track of the gaze tracking quality. Still, we applied this rather strict criterion because we needed good continuous gaze tracking data during item presentation to be able to calculate the presented gaze parameters.

In the present study, we presented data on Gestalt perception performance in combination with looking behavior in visually impaired children. Further research is necessary to address causality of the results: did a different way of looking result in worse perceptual performance, or did poor

perception lead to a different looking pattern? We would like to stress that the analysis of eye movement patterns during perception tasks adds value for the visual testing of ‘non-verbal children’ and may provide a link to later visual training.

## Conclusions

This study showed that Dutch children with visual impairments more often performed weakly on the Kaufman Gestalt closure task than chronologically age-matched American controls, and that children with brain damage performed worse than children without brain damage. Notably, children with brain damage (groups CVI and OCVI) performed significantly worse on the animate items than the group without brain damage (OVI). Based on the looking patterns obtained with the eye tracker, we can conclude that weak Gestalt performance of the children in the CVI group is not the result of intellectual disability or a lack of effort, but instead seems a pure perceptual dysfunction.

## ORCID

Johan J. M. Pel  <http://orcid.org/0000-0001-7618-0413>

## References

- Bölte, S., Holtmann, M., Poustka, F., Scheurich, A., & Schmidt, L. (2007). Gestalt perception and local-global processing in high-functioning autism. *Journal of Autism and Developmental Disorders*, 37, 1493–1504. doi:10.1007/s10803-006-0231-x
- Caramazza, A., & Shelton, J. R. (1998). Domain-specific knowledge systems in the brain the animate-inanimate distinction. *Journal of Cognitive Neuroscience*, 10, 1–34.
- Doniger, G. M., Foxe, J. J., Murray, M. M., Higgins, B. A., & Javitt, D. C. (2002). Impaired visual object recognition and dorsal/ventral stream interaction in schizophrenia. *Archives of General Psychiatry*, 59, 1011–1020.
- Doniger, G. M., Foxe, J. J., Murray, M. M., Higgins, B. A., Snodgrass, J. G., & Schroeder, C. E. (2000). Activation timecourse of ventral visual stream object-recognition areas: High density electrical mapping of perceptual closure processes. *Journal of Cognitive Neuroscience*, 12, 615–621.
- Fazzi, E., Bova, S. M., Uggetti, C., Signorini, S. G., Bianchi, P. E., & Maraucci, I. (2004). Visual-perceptual impairment in children with periventricular leukomalacia. *Brain and Development*, 26, 506–512. doi:10.1016/j.braindev.2004.02.002
- Han, S., & Humphreys, G. W. (1999). Interactions between perceptual organization based on Gestalt laws and those based on hierarchical processing. *Perception and Psychophysics*, 61, 1287–1298.
- Hillis, A. E., & Caramazza, A. (1991). Mechanisms for accessing lexical representations for output: Evidence from a category-specific semantic deficit. *Brain and Language*, 40, 106–144.
- Hodapp, R. M., Leckman, J. F., Dykens, E. M., Sparrow, S. S., Zelinsky, D. G., & Ort, S. I. (1992). K-ABC profiles in children with fragile X syndrome, Down syndrome, and nonspecific mental retardation. *American Journal of Mental Retardation*, 97, 39–46.
- Huberle, E., & Karnath, H. O. (2012). The role of temporo-parietal junction (TPJ) in global Gestalt perception. *Brain Structure and Function*, 217, 735–746. doi:10.1007/s00429-011-0369-y
- Jäkel, F., Singh, M., Wichmann, F. A., & Herzog, M. H. (2016). An overview of quantitative approaches in Gestalt perception. *Vision Research*, 126, 3–8. doi:10.1016/j.visres.2016.06.004
- Katsuki, F., & Constantinidis, C. (2014). Bottom-up and top-down attention: Different processes and overlapping neural systems. *Neuroscientist*, 20, 509–521. doi:10.1177/1073858413514136
- Kaufman, A., & Kaufman, N. (2004). *Kaufman assessment battery for children: Technical manual* (2nd ed.). Circle Pines, MN: American Guidance Service.
- Kooiker, M. J., Verbunt, H. J., van der Steen, J., & Pel, J. J. (2019). Combining visual sensory functions and visuospatial orienting functions in children with visual pathology: A longitudinal study. *Brain and Development*, 41(2), 135–149. doi:10.1016/j.braindev.2018.09.006
- Kooiker, M. J., van der Steen, J., & Pel, J. J. (2016). Development of salience-driven and visually-guided eye movement responses. *Journal of Vision*, 16, 18. doi:10.1167/16.5.18
- Koppitz, E. M. (1962). Diagnosing brain damage in young children with the Bender Gestalt test. *Journal of Consulting Psychology*, 26, 541–546.

- Kriegeskorte, N., Mur, M., Ruff, D. A., Kiani, R., Bodurka, J., & Esteky, H. (2008). Matching categorical object representations in inferior temporal cortex of man and monkey. *Neuron*, 60, 1126–1141. doi:[10.1016/j.neuron.2008.10.043](https://doi.org/10.1016/j.neuron.2008.10.043)
- Larroque, B., Ancel, P. Y., Marret, S., Marchand, L., André, M., & Arnaud, C. (2008). Neurodevelopmental disabilities and special care of 5-year-old children born before 33 weeks of gestation (the EPIPAGE study): A longitudinal cohort study. *Lancet*, 371, 813–820. doi:[10.1016/S0140-6736\(08\)60380-3](https://doi.org/10.1016/S0140-6736(08)60380-3)
- Le Grand, R., Mondloch, C. J., Maurer, D., & Brent, H. P. (2004). Impairment in holistic face processing following early visual deprivation. *Psychological Science*, 15, 762–768. doi:[10.1111/j.0956-7976.2004.00753.x](https://doi.org/10.1111/j.0956-7976.2004.00753.x)
- Marini, F., & Marzi, C. A. (2016). Gestalt perceptual organization of visual stimuli captures attention automatically: Electrophysiological evidence. *Frontiers in Human Neuroscience*, 10, 446. doi:[10.3389/fnhum.2016.00446](https://doi.org/10.3389/fnhum.2016.00446)
- Naselaris, T., Stansbury, D. E., & Gallant, J. L. (2012). Cortical representation of animate and inanimate objects in complex natural scenes. *Journal of Physiology*, 106, 239–249. doi:[10.1016/j.jphysparis.2012.02.001](https://doi.org/10.1016/j.jphysparis.2012.02.001)
- Pflugshaupt, T., Nösberger, M., Gutbrod, K., Weber, K. P., Linnebank, M., & Brugger, P. (2016). Bottom-up visual integration in the medial parietal lobe. *Cerebral Cortex*, 26, 943–949. doi:[10.1093/cercor/bhu256](https://doi.org/10.1093/cercor/bhu256)
- Ploran, E. J., Nelson, S. M., Velanova, K., Donaldson, D. I., Petersen, S. E., & Wheeler, M. E. (2007). Evidence accumulation and the moment of recognition: Dissociating perceptual recognition processes using fMRI. *Journal of Neuroscience*, 27, 11912–11924. doi:[10.1523/JNEUROSCI.3522-07.2007](https://doi.org/10.1523/JNEUROSCI.3522-07.2007)
- Russell, E. W., Hendrickson, M. E., & VanEaton, E. (1988). Verbal and figural gestalt completion tests with lateralized occipital area brain damage. *Journal of Clinical Psychology*, 44, 217–225.
- Uhlhaas, P. J., Phillips, W. A., Mitchell, G., & Silverstein, S. M. (2006). Perceptual grouping in disorganized schizophrenia. *Psychiatry Research*, 145, 105–117. doi:[10.1016/j.psychres.2005.10.016](https://doi.org/10.1016/j.psychres.2005.10.016)
- van de Ven, V., Rotarska-Jagiela, A., Oertel-Knöchel, V., & Linden, D. E. J. (2017). Reduced intrinsic visual cortical connectivity is associated with impaired perceptual closure in schizophrenia. *Neuroimage Clinical*, 15, 45–52. doi:[10.1016/j.nicl.2017.04.012](https://doi.org/10.1016/j.nicl.2017.04.012)
- Wagemans, J., Elder, J. H., Kubovy, M., Palmer, S. E., Peterson, M. A., & Singh, M. (2012). A century of Gestalt psychology in visual perception: I. Perceptual grouping and figure-ground organization. *Psychological Bulletin*, 138, 1172–1217. doi:[10.1037/a0029333](https://doi.org/10.1037/a0029333)
- Warrington, E. K., & Shallice, T. (1984). Category specific semantic impairments. *Brain*, 107, 829–854.
- World Medical Association (2013). World Medical Association Declaration of Helsinki: ethical principles for medical research involving human subjects. *Journal of the American Medical Association*, 310(20), 2191–2194. doi:[10.1001/jama.2013.281053](https://doi.org/10.1001/jama.2013.281053).