



OPEN ACCESS



Check for updates

Colorectal cancer screening with faecal immunochemical testing, sigmoidoscopy or colonoscopy: a microsimulation modelling study

Maaïke Buskermolen,¹ Dayna R Cenin,^{1,2} Lise M Helsingen,^{3,4,5} Gordon Guyatt,⁶ Per Olav Vandvik,⁷ Ulrike Haug,^{8,9} Michael Bretthauer,^{3,4,5} Iris Lansdorp-Vogelaar¹

For numbered affiliations see end of the article.

Correspondence to: M Buskermolen
m.buskermolen@erasmusmc.nl
(ORCID 0000-003-3417-1994)

Additional material is published online only. To view please visit the journal online.

Cite this as: *BMJ* 2019;367:l5383
<http://dx.doi.org/10.1136/bmj.l5383>

Accepted: 14 September 2019

ABSTRACT

OBJECTIVE

To estimate benefits and harms of different colorectal cancer screening strategies, stratified by (baseline) 15-year colorectal cancer risk.

DESIGN

Microsimulation modelling study using Microsimulation SCreening ANalysis-Colon (MISCAN-Colon).

SETTING

A parallel guideline committee (*BMJ* Rapid Recommendations) defined the time frame and screening interventions, including selection of outcome measures.

POPULATION

Norwegian men and women aged 50-79 years with varying 15-year colorectal cancer risk (1-7%).

COMPARISONS

Four screening strategies were compared with no screening: biennial or annual faecal immunochemical

test (FIT) or single sigmoidoscopy or colonoscopy at 100% adherence.

MAIN OUTCOME MEASURES

Colorectal cancer mortality and incidence, burdens, and harms over 15 years of follow-up. The certainty of the evidence was assessed using the GRADE approach.

RESULTS

Over 15 years of follow-up, screening individuals aged 50-79 at 3% risk of colorectal cancer with annual FIT or single colonoscopy reduced colorectal cancer mortality by 6 per 1000 individuals. Single sigmoidoscopy and biennial FIT reduced it by 5 per 1000 individuals. Colonoscopy, sigmoidoscopy, and annual FIT reduced colorectal cancer incidence by 10, 8, and 4 per 1000 individuals, respectively. The estimated incidence reduction for biennial FIT was 1 per 1000 individuals. Serious harms were estimated to be between 3 per 1000 (biennial FIT) and 5 per 1000 individuals (colonoscopy); harms increased with older age. The absolute benefits of screening increased with increasing colorectal cancer risk, while harms were less affected by baseline risk. Results were sensitive to the setting defined by the guideline panel. Because of uncertainty associated with modelling assumptions, we applied a GRADE rating of low certainty evidence to all estimates.

CONCLUSIONS

Over a 15 year period, all screening strategies may reduce colorectal cancer mortality to a similar extent. Colonoscopy and sigmoidoscopy may also reduce colorectal cancer incidence, while FIT shows a smaller incidence reduction. Harms are rare and of similar magnitude for all screening strategies.

Introduction

Colorectal cancer is a public health issue, with an estimated 1.4 million new cases and 700 000 deaths worldwide in 2018.¹ Screening is intended to reduce colorectal cancer incidence and mortality, and its effectiveness has been demonstrated in randomised controlled trials of guaiac faecal occult blood testing (gFOBT) and sigmoidoscopy.²⁻¹¹ Colonoscopy is likely to be at least as effective as sigmoidoscopy since it reaches the whole large bowel, whereas sigmoidoscopy reaches only the distal part of the large bowel. Faecal immunochemical testing (FIT) is likely to be at least as effective as gFOBT, since both tests detect blood in stools, and FIT demonstrates higher sensitivity and specificity.¹² However, because of the lack of published evidence from randomised trials for colonoscopy and

WHAT IS ALREADY KNOWN ON THIS TOPIC

Randomised trials with 15 or more years of follow-up have shown that sigmoidoscopy and guaiac faecal occult blood test (gFOBT) screening reduce colorectal cancer mortality

Colonoscopy and faecal immunochemical test (FIT) are increasingly used for colorectal cancer screening, but there are no published data from randomised trials on the relative effectiveness of the different screening strategies

Although risks and benefits differ across individuals, there is limited evidence for individualised screening recommendations. Basing screening on individual cancer risk, determined in large part by age and sex, may be desirable

WHAT THIS STUDY ADDS

This modelling study found that, over a 15 year period, colorectal cancer mortality reduction may be very similar with a single colonoscopy or sigmoidoscopy, or annual or biennial FIT. Colonoscopy and sigmoidoscopy may result in similar reductions in colorectal cancer incidence, while the effect of FIT on incidence may be smaller or absent

At a baseline cancer risk of 3%, MISCAN-Colon estimates for 1000 individuals over 15 years are:

Colorectal cancer deaths prevented: colonoscopy 6, sigmoidoscopy 5, annual FIT 6, biennial FIT 5

Colorectal cancer cases prevented: colonoscopy 10, sigmoidoscopy 8, annual FIT 4, biennial FIT 1

Harms were similar across screening strategies, with serious colonoscopic complications of <5 per 1000 at highest risk levels

Screening benefits and harms may increase as risk of colorectal cancer increases; benefits more so than harms

FIT screening, it is not known if they are more effective than gFOBT and sigmoidoscopy.¹³

Despite its benefits, colorectal cancer screening can be burdensome, and colonoscopy is associated with rare but serious complications. In addition, screening performance depends on the baseline risk of cancer for individuals. Few studies and guidelines have incorporated how baseline risk affects the balance between benefits, burden, and harms of screening in the past.

To elucidate these issues, we undertook microsimulation modelling as part of the *BMJ* Rapid Recommendations project, a collaborative effort from the MAGIC research and innovation programme (www.magicproject.org) and *The BMJ*. The aim of the Rapid Recommendations is to respond to new, potentially practice changing, evidence and provide trustworthy practice guidelines in a timely manner. The *BMJ* Rapid Recommendations project for colorectal cancer screening was triggered by recent updates from three large randomised trials on sigmoidoscopy screening with follow-up data of 15 years or longer.^{3 5 14} In light of this new evidence, we addressed the potential benefits and harms of colorectal cancer screening with annual or biennial FIT or a single sigmoidoscopy or colonoscopy in the time frame of 15 years. This work informed the parallel guideline published in a multilayered electronic format on bmj.com and MAGICapp. Box 1 shows the articles and evidence linked to this Rapid Recommendation.

Methods

At the request of the guideline panel (Helsingen et al¹⁵), we applied the Microsimulation Screening Analysis-Colon (MISCAN-Colon) model to simulate 15 years of follow-up of population cohorts aged 50–79 years. We estimated the benefits, burden, and harms of the following four screening strategies: annual FIT, biennial FIT, a single sigmoidoscopy, and a single colonoscopy. We compared the four screening strategies with each other and with no screening. We performed the analyses stratified by different levels of baseline colorectal cancer risk. Additionally, we determined screening benefits for men and women separately, and screening harms for different age groups, as requested by the panel.

MISCAN-Colon model

MISCAN-Colon is a well established microsimulation model for colorectal cancer.^{17 18} In brief, MISCAN-Colon simulates life histories of a large group of individuals from birth to death. In addition, the model simulates the development of colorectal cancer through the adenoma carcinoma sequence. As each simulated person ages, one or more adenomas may develop. These adenomas can progress in size increasing from small (≤ 5 mm) to medium (6–9 mm) to large (≥ 10 mm). Some adenomas can develop into preclinical cancer, which may progress through cancer stages I to IV. At any time during the development of the disease, the process may be interrupted because a person dies of

other causes. With screening, colorectal cancer may either be prevented (by the detection and removal of adenomas) or detected at an earlier stage with a more favourable prognosis. In this way, colorectal cancer incidence and/or colorectal cancer mortality may be reduced. The model also estimates harms associated with screening.

The quantification and model assumptions are described in detail in appendix 1. In brief, the age-specific prevalence and multiplicity distribution of adenomas (the distribution of the individual number of adenomas across the population) were calibrated using autopsy studies.^{19–29} The duration of preclinical colorectal cancer (sojourn time) and the adenoma dwell-time (the duration of progression of adenomas) were calibrated using rates of interval cancers (cancers that are diagnosed between screening tests) and surveillance detected cancers (found during surveillance) in randomised gFOBT and sigmoidoscopy trials.^{8 30–35}

For this study, we developed a MISCAN-Colon model version calibrated to the sex-, age-, stage-, and localisation-specific colorectal cancer incidence and survival as observed in Norway during the timeframe of the NORCCAP trial (1999–2011) (appendix 2 (fig 1)), using data provided by the Norwegian Cancer Registry.³⁶ Life expectancy was based on sex-specific lifetables for 2007, the middle of the NORCCAP trial period, from Statistics Norway.³⁷ We validated this model using the results of one of the trigger publications: 15-year follow-up data from the Norwegian Colorectal Cancer Prevention (NORCCAP) trial.⁵ The validation methods and results are described in appendix 3.

Simulated cohorts

We simulated seven population cohorts consisting of men and women aged 50–79 years with a 15-year colorectal cancer risk varying from 1% to 7%, using the same Norwegian sex-specific MISCAN-Colon versions as we used for the validation. The age-specific onset of adenomas in MISCAN-Colon for all ages was adjusted to match the 15-year colorectal cancer risk in the seven cohorts. The modelled risk levels were chosen to cover the majority of individuals under consideration for this study (healthy people aged 50 to 79 years), but still with a manageable number of risk levels. We used the range of risk levels found when applying the QCancer risk prediction model to the UK Biobank cohort as guidance.³⁸ The simulated risk levels from 1% to 7% cover approximately 90% of the colorectal cancer risk levels found in the UK Biobank cohort (personal communication UK Biobank researcher Juliet Usher-Smith). We confirmed that the chosen risk levels also cover the range of risk levels observed in the general population, by comparing the risk levels with the 15 year colorectal cancer risk ranges found in two large population based cancer databases.^{39 40} Data from the UK Biobank were also used to validate the QCancer prediction model for colorectal cancer. The QCancer Calculator is an open-access online tool that aims to predict individual colorectal cancer risk

Box 1: Linked resources for this BMJ Rapid Recommendations cluster

- Helsingen LM, Vandvik PO, Jodal HC, et al. Colorectal cancer screening with faecal immunochemical testing, sigmoidoscopy or colonoscopy: a clinical practice guideline. *BMJ* 2019;367:l5515.15
- Summary of the results from the Rapid Recommendation process
- Jodal HC, Helsingen LM, Anderson JC, et al. Colorectal cancer screening with faecal testing, sigmoidoscopy or colonoscopy: a systematic review and network meta-analysis. *BMJ Open* 2019;0:e032773.16
- Review of all available trials that assessed colorectal cancer screening
- Buskermolen M, Cenin DR, Helsingen LM, et al. Colorectal cancer screening with faecal immunochemical testing, sigmoidoscopy or colonoscopy: a microsimulation modelling study. *BMJ* 2019;367:l5383.
- Modelling study of different modalities for colorectal cancer screening
- MAGICApp (<http://magicproject.org/190220dist>)
- Expanded version of the results with evidence summaries and decision aids for use on all devices

based on risk factors such as medical history, lifestyle factors, and ethnicity.⁴¹ Individuals may predict their 15-year colorectal cancer risk with this calculator (<https://qcancer.org/15yr/colorectal/index.php>). Subsequently, they may use this predicted colorectal cancer risk to look up the best risk-matching MISCAN-Colon predictions of screening outcomes, to get a personal estimate of the magnitude of benefits and harms of colorectal cancer screening. The clinical practice guideline presents relevant details.¹⁵

Screening strategies

For each cohort, we assessed four colorectal cancer screening strategies during a 15-year period: biennial FIT, annual FIT, a single sigmoidoscopy, and colonoscopy. All strategies were compared with no screening. For FIT, we chose a cut-off of 20 µg Hb/g faeces since this is used in many screening programmes.⁴² We assumed that individuals with a positive FIT result and those with at least one adenoma (of any size) diagnosed at sigmoidoscopy screening were referred for colonoscopy.

Sensitivity and specificity of the screening tests were based on diagnostic test accuracy studies (table 1).^{43 47} Age-specific risks for complications associated with colonoscopy were derived from SEER-Medicare data (table 1).^{45 48 49} Only complications requiring hospital admission within 30 days after the colonoscopy were taken into account. Only colonoscopies with polypectomies were considered to cause adverse effects. We included colorectal perforations and bleedings, other gastrointestinal adverse events, cardiovascular adverse events, and mortality related to screening procedure. The number of complications was calculated by multiplying the number of colonoscopies with polypectomies by each complication risk. The simulated cohorts were assumed to have no prior screening.

We simulated surveillance consistent with European Society of Gastrointestinal Endoscopy Guidelines.⁵⁰ Individuals with low risk findings (fewer than three low risk adenomas (<10mm diameter) at primary screening) did not receive any surveillance, whereas

individuals with high risk findings were offered surveillance with colonoscopy after three years, and thereafter colonoscopies repeated at intervals of three to five years depending on the findings.

As our aim is to provide individuals who are considering screening with estimates of the possible benefits and harms of participation, we assumed 100% adherence to screening, follow-up and surveillance for all analyses.

Outcomes

We distinguished the three screening-related outcome groups for the 15-year follow-up time frame chosen by the *BMJ* Rapid Recommendation guideline panel: benefits of screening, screening harms, and screening burden. For benefits of screening, we present model-predicted colorectal cancer incidence and mortality, and all-cause mortality reduction. For screening burden, we present the number of screening tests, number of individuals with at least one colonoscopy (including screening colonoscopies), and number of individuals with at least two colonoscopies (for example, individuals with at least one surveillance colonoscopy). For screening harms, we present risk of screening related colorectal perforations and bleedings, other gastrointestinal adverse events, cardiovascular adverse events, and mortality related to screening procedure.

Sensitivity analyses

As a one-way sensitivity analysis, we assessed results stratified by age and sex; and we assessed outcomes with lifetime follow-up instead of a 15-year follow-up.

Certainty of evidence

We used the GRADE approach to address the certainty of the evidence.⁵¹ GRADE has not yet produced detailed guidance for assessing certainty of evidence in modelling studies. To make our assessment, we considered uncertainty associated with key inputs into the model.

Guideline panel and patient involvement

According to the *BMJ* Rapid Recommendations process, a multiprofessional guideline panel that included three patients who have experienced colorectal cancer screening provided oversight to the study and identified the population and outcomes of interest.

Results

The Rapid Recommendation panel suggests against screening if the risk is below a 15-year colorectal cancer risk of 3% and suggests screening if the risk is above 3%.¹⁵ For simplification, we only describe the estimates for individuals with the 3% risk level. Estimates for all other risk levels are provided in table 2 and figure 1.

Benefits of screening

MISCAN-Colon predicted that all screening strategies reduced colorectal cancer incidence and mortality

across all colorectal cancer risk groups (fig 1a and b, table 2). Colonoscopy showed the largest reduction in colorectal cancer mortality, but the differences between the screening strategies were small and the reduction may be similar for all strategies. For instance, the model predicted that, at 3% risk without

screening, colorectal cancer mortality was nine deaths per 1000 individuals. There were six per 1000 fewer colorectal cancer deaths with colonoscopy and annual FIT (approximately 60% reduction) and five per 1000 fewer deaths with sigmoidoscopy and biennial FIT (approximately 50% reduction).

Table 1 | Key modelling assumptions used in study

Input parameter	Base-case assumption	References
Demography		
All-cause mortality	Norwegian lifetables 2007	Statistics Norway
Natural course of cancer		
Adenoma onset	Age dependent (non-homogeneous Poisson)	*
Adenoma progression:		
State transitions	Age dependent	*
State duration (years, total)	Exp($\lambda=130$)	*
Cancer progression (preclinical):		
Stage transitions	Age-dependent	*
Stage durations (years)	Exp ($\lambda=2.5$)	*
Colorectal cancer incidence (without exposure to screening)	Age-, stage-, and location-dependent	Norwegian Cancer Registry
Colorectal cancer stage distribution	Age- and location-dependent	Norwegian Cancer Registry
Colorectal cancer survival	Age-, stage-, and location-dependent	Norwegian Cancer Registry
Colonoscopy quality		
Sensitivity (%)†:		
Adenomas 0-5 mm	75%	*
Adenomas 6-9 mm	85%	*, van Rijn et al ⁴³
Adenomas ≥10 mm	95%	*, van Rijn et al ⁴³
Malignant neoplasia	95%	*
Specificity (%)	100%	
Complete colonoscopy examination (%)‡	Men 93.5%, women 85.2%	Holme et al ⁴⁴
Complication rates (%)§:		
With polypectomy	Age dependent (50-79 years)	van Hees et al ⁴⁵
Perforations and bleeding	0.2-0.9	
Other GI adverse events	0.2-0.8	
Cardiovascular events	0.1-0.7	
Screening procedure related mortality	0.008-0.04	
Without polypectomy§	-	
Sigmoidoscopy quality		
Sensitivity (%)†:		
Adenomas 0-5 mm	75%	*
Adenomas 6-9 mm	85%	*, van Rijn et al ⁴³
Adenomas ≥10 mm	95%	*, van Rijn et al ⁴³
Malignant neoplasia	95%	*
Specificity (%)¶	98.2%	Buskermolen et al ⁴⁶
Complete examination (%)**	Men 93.2%, women 83.8%	Holme et al ⁴⁴
Faecal immunochemical test (FIT) quality		
Sensitivity (%)†:		
Adenomas 0-5 mm	0	
Adenomas 6-9 mm	11.4	
Adenomas ≥10 mm	15.9	
Malignant neoplasia:		
Short before clinical detection	88.6	
Long before clinical detection	62.6	
Specificity (%)¶	97.6	

Poisson=Poisson distribution; Exp=exponential distribution. GI=gastrointestinal.

*More details available in appendix 1.

†Sensitivity was defined as the probability of detecting an adenoma that was present at the time of test.

‡Colonoscopy was considered complete if the caecum was reached. In the incomplete examinations, the endpoint was assumed to be distributed uniformly over the colon/rectum.

§We assumed that colonoscopy without polypectomy was not associated with a higher risk of complications. The risk of complications for polypectomy was assumed to increase exponentially with age. Perforation and bleeding concerned adverse events requiring blood transfusions; other GI adverse events included paralytic ileus, nausea, vomiting and dehydration, abdominal pain; and cardiovascular adverse events included myocardial infarction or angina, arrhythmias, congestive heart failure, cardiac or respiratory arrest, or syncope, hypotension, or shock. The screening procedure related mortality was derived from estimates of the incidence of perforation and case-fatality for perforation.^{45 48 49}

¶The lack of specificity indicates how many of the tests that did not detect adenomatous lesions resulted in a referral for follow-up colonoscopy. The MISCAN-Colon model is a natural history microsimulation model simulating onset of adenomas in some individuals, that may progress to colorectal cancer in some cases. The model does not explicitly simulate the presence of blood in stool. To simulate FIT screening, we rather use estimates of per-person sensitivity and specificity by disease status based on a study by Imperiale et al. with a cut-off of 20 µg of haemoglobin per g of faeces.⁴⁷ We fitted per-lesion sensitivity and per-person specificity of the model to the per-person sensitivity and specificity estimates found in the study. In the model, the probability for a person to test positive depends on the lack of specificity and the per-lesion sensitivity for the lesions present in that individual.

**Flexible sigmoidoscopy was considered complete if the junction of the sigmoid/colon descendens was reached. In the incomplete examinations, the endpoint was assumed to be distributed uniformly over the rectosigmoid.

Table 2 | Predictions of benefits and harms of screening for individuals aged 50-79 years at varying levels of colorectal cancer risk. All outcomes are given per 1000 screened individuals over 15 years and compared to a scenario with no screening.

Screening strategy	Colorectal cancer					Risk of complications						
	Incidence reduction (%)	No of cases prevented	Mortality reduction (%)	No of deaths prevented	All-cause mortality reduction (%)	No of screening tests	No of individuals with ≥ 1 colonoscopy	No of individuals with ≥ 2 colonoscopies	Perforation and bleeding	Other GI events	Cardio-vascular events	Screen procedure related mortality
1% colorectal cancer risk without screening (10 cases per 1000, with risk of dying of colorectal cancer of 0.3% (3 per 1000))												
Biennial FIT	8	1	53	2	0.4	5441	160	26	0.3	0.3	0.3	0.02
Annual FIT	18	2	62	2	0.5	9464	254	31	0.4	0.4	0.3	0.02
Single sigmoidoscopy	29	3	57	2	0.4	1000	82	28	0.4	0.4	0.3	0.02
Single colonoscopy	35	3	66	2	0.5	1000	1000	33	0.5	0.5	0.4	0.03
2% colorectal cancer risk without screening (20 cases per 1000, with risk of dying of colorectal cancer of 0.6% (6 per 1000))												
Biennial FIT	6	1	51	3	0.8	5288	203	54	0.7	0.7	0.5	0.04
Annual FIT	16	3	60	4	0.9	9091	300	66	0.9	0.9	0.6	0.05
Single sigmoidoscopy	28	6	54	3	0.8	1000	159	57	0.8	0.8	0.6	0.05
Single colonoscopy	34	7	64	4	0.9	1000	1000	68	1.1	1.1	0.8	0.07
3% colorectal cancer risk without screening (30 cases per 1000, with risk of dying of colorectal cancer of 0.9% (9 per 1000))												
Biennial FIT	5	1	50	5	1.1	5134	246	83	1.1	1	0.8	0.06
Annual FIT	15	4	59	6	1.3	8715	347	101	1.3	1.3	1	0.08
Single sigmoidoscopy	27	8	52	5	1.1	1000	237	86	1.3	1.2	0.9	0.07
Single colonoscopy	34	10	63	6	1.4	1000	1000	105	1.7	1.7	1.2	0.1
4% colorectal cancer risk without screening (40 cases per 1000, with risk of dying of colorectal cancer of 1.2% (12 per 1000))												
Biennial FIT	4	2	49	6	1.5	4980	288	112	1.4	1.4	1.1	0.09
Annual FIT	15	6	59	7	1.8	8346	391	138	1.8	1.7	1.3	0.11
Single sigmoidoscopy	27	11	52	6	1.5	1000	312	119	1.7	1.7	1.2	0.1
Single colonoscopy	34	14	63	8	1.8	1000	1000	144	2.3	2.3	1.7	0.14
5% colorectal cancer risk without screening (50 cases per 1000, with risk of dying of colorectal cancer of 1.6% (16 per 1000))												
Biennial FIT	4	2	49	8	1.8	4834	328	142	1.8	1.8	1.3	0.11
Annual FIT	15	7	59	9	2.2	7996	433	175	2.2	2.2	1.6	0.13
Single sigmoidoscopy	27	14	52	8	1.8	1000	382	153	2.2	2.1	1.6	0.13
Single colonoscopy	34	17	63	10	2.4	1000	1000	184	2.9	2.8	2.1	0.17
6% colorectal cancer risk without screening (60 cases per 1000, with risk of dying of colorectal cancer of 1.9% (19 per 1000))												
Biennial FIT	5	3	50	9	2.2	4694	365	171	2.2	2.1	1.6	0.13
Annual FIT	15	9	59	11	2.6	7664	472	211	2.7	2.6	2	0.16
Single sigmoidoscopy	28	17	53	10	2.2	1000	446	189	2.7	2.6	1.9	0.16
Single colonoscopy	34	21	64	12	2.8	1000	1000	226	3.5	3.4	2.5	0.2
7% colorectal cancer risk without screening (70 cases per 1000, with risk of dying of colorectal cancer of 2.2% (22 per 1000))												
Biennial FIT	5	4	50	11	2.6	4558	401	201	2.6	2.5	1.9	0.15
Annual FIT	16	11	59	13	3.0	7346	509	247	3.1	3.1	2.3	0.18
Single sigmoidoscopy	29	20	54	12	2.6	1000	505	228	3.2	3.1	2.3	0.19
Single colonoscopy	35	24	64	14	3.2	1000	1000	268	4	3.9	2.9	0.24

FIT=faecal immunochemical test.

Colonoscopy may be the most effective strategy in reducing colorectal cancer incidence, followed by sigmoidoscopy (fig 1b, table 2). Colonoscopy was estimated to reduce colorectal cancer incidence by 10 colorectal cancer cases and sigmoidoscopy by eight per 1000 individuals (approximately 30% reduction); annual FIT had four and biennial FIT had one incident cancer cases per 1000 screened (approximately 10% and 5% reduction).

The estimated relative effects were similar across the different levels of baseline risk (1% to 7% over

15 years) for both colorectal cancer incidence and mortality, resulting in larger absolute effects for colorectal cancer incidence and mortality for individuals with higher risk. The model assumes that prevented colorectal cancer deaths lead to a corresponding reduction in all-cause mortality. At 3% risk, the estimated relative reduction in all-cause mortality was around 1.5% (table 2), corresponding to a reduction of five per 1000 individuals (all-cause mortality reduced from 328 to 323).




Fig 1 | MISCAN-Colon predictions of colorectal cancer mortality and incidence reduction, screen tests, colonoscopies and complications per 1000 individuals, using biennial or annual faecal immunochemical test (FIT), flexible sigmoidoscopy, or colonoscopy. Results are stratified by colorectal cancer risk. Individuals were followed-up for 15 years

Screening burden

The two FIT strategies required the most screening tests (fig 1c, table 2). The number of FIT rounds for the simulated cohort depended on the age of the individual at the first screening and the screening interval. Individuals ≤ 65 years old at cohort entry received seven rounds with biennial FIT and 15 rounds with annual FIT. However, many individuals received fewer screening rounds: individuals > 65 years at cohort entry because they stopped screening after age 79; individuals who died within 15 years of follow-up; individuals who tested FIT positive and were referred for a colonoscopy and therefore received screening according to the surveillance guidelines. For individuals with a 3% colorectal cancer risk, we predicted that approximately 8700 tests were required with annual FIT screening, and 5100 with biennial FIT screening. Colonoscopy screening resulted in the highest number of individuals receiving at least one colonoscopy, since all individuals received colonoscopy as a screening test (fig 1d, table 2). The number of individuals with at least two colonoscopies (that is, individuals with at least one surveillance colonoscopy) was highest for colonoscopy regardless of colorectal cancer risk (table 2).

Screening harms

The risk of screening related mortality, colorectal perforations and bleedings, other serious gastrointestinal adverse events, or cardiovascular adverse events was proportional to the number of required colonoscopies (fig 1e-g). In the 3% colorectal cancer risk group, the predicted overall complication risk was lowest with biennial FIT screening (2.9 per 1000) and highest with colonoscopy screening (4.6 per 1000) (table 2).

Sensitivity analyses

The model predicted similar reductions in cancer incidence and mortality for the screening tests for men and women for all levels of 15-year baseline risk of colorectal cancer, except for sigmoidoscopy, for which the model predicted that women may benefit slightly less than men (appendix 4 (tables 1-12, fig 1)).

When results were stratified by age, annual FIT was more effective in reducing colorectal cancer mortality than a single colonoscopy in younger individuals (50-59 years old). In older individuals (75-79 years), colonoscopy and sigmoidoscopy were more effective. In addition, the estimated risk of complications increased more with age for the colonoscopy and sigmoidoscopy screening strategies than for the FIT strategies, although the increase in risk in absolute numbers was small (appendix 4 (tables 1-12, fig 2)). For example, at 3% colorectal cancer risk, a single colonoscopy strategy resulted in colorectal perforations and bleedings of 1.6 per 1000 in men aged 50-54 years and 3.4 per 1000 in men aged 75-79 years.

When we considered lifetime follow-up, the model predicted larger absolute reductions in colorectal

cancer incidence and mortality for all screening strategies (appendix 5 (figs 1 and 2)), and less differences between annual FIT and colonoscopy. At younger ages (50-64 years) the model predicted that annual FIT was more effective at reducing colorectal cancer incidence and mortality than a single colonoscopy, although the difference was small. For example, annual FIT screening prevented 43 lifetime colorectal cancer cases in those aged 50-54 compared with 37 prevented cases with a single colonoscopy.

Certainty of evidence

We noted appreciable uncertainty associated with the following model inputs: (a) all colorectal cancers develop through adenomas; (b) differences in colorectal cancer risk are caused by differences in adenoma incidence; and (c) adenoma dwell time. There is no high certainty data to inform the model regarding these model inputs. Therefore, despite the predictive and external validity of the model against the NORCCAP study (appendix 3), the panel considered all model estimates as low certainty evidence.

Discussion

Our microsimulation model analysis suggests that all screening strategies reduce colorectal cancer mortality during 15 years of follow-up, and colonoscopy, sigmoidoscopy, and annual FIT may also reduce colorectal cancer incidence. The extent of absolute risk reduction varies with baseline colorectal cancer risk. Few differences were observed when results were stratified by sex. When outcomes were stratified by age, we observed that FIT screening strategies were estimated to be more effective in younger individuals, while colonoscopy and sigmoidoscopy were more effective in older individuals. FIT screening strategies required the highest overall number of screening tests, and colonoscopy screening resulted in the highest number of colonoscopies, regardless of colorectal cancer risk. Consequently, we observed the highest probability of experiencing a complication in individuals who underwent screening with a single colonoscopy, with increasing risk in older age groups.

Colonoscopy may result in the largest reduction in colorectal cancer incidence, followed by sigmoidoscopy and annual FIT. However, this finding was sensitive to the follow-up setting defined by the guideline panel. The panel chose a 15 year follow-up period because this allowed model predictions to be validated against trial data. With lifetime follow-up, the model predicted that relative incidence and mortality reductions from screening would persist for some more years, resulting in larger absolute numbers of prevented colorectal cancer cases and deaths. This was especially true for FIT, where lifetime follow-up resulted in higher estimates for colorectal cancer incidence reductions, making the test comparable to colonoscopy. However, because longer term follow-up data from trials are lacking, lifetime estimates are more uncertain and were thus not taken into account by the panel.

Strengths and limitations of this study

Model predictions for colorectal cancer mortality and incidence reduction in this study are considerably higher than those observed in randomised screening trials. These seemingly discrepant results can be explained by our assumption of 100% adherence to screening tests, work-up, and surveillance colonoscopies, whereas trial outcomes are the result of real-world adherence patterns, which are considerably lower. When we replicated the NORCCAP trial population including observed adherence patterns, MISCAN-Colon predictions for colorectal cancer incidence and mortality reduction were in line with the trial results (appendix 3). Moreover, when we replicated the design of an Italian cohort study, the model predicted reductions in colorectal cancer incidence and mortality resulting from FIT screening that aligned well with those results observed in that study (data not shown).⁵²

We acknowledge that 100% adherence gives estimates that are higher than what would be expected in a population screening programme, when adherence is never 100%. However, our intention was to inform individuals about expected effectiveness when they (fully) participate in screening rather than considering the impact of a screening programme from a public health perspective and assessing results at the population level. The aim of this study was to support the *BMJ* Rapid Recommendation panel by comparing different screening strategies, stratified by baseline 15-year colorectal cancer risk, using microsimulation modelling. A strength of this work is that we validated our model using the recently published 15-year follow-up results of the NORCCAP trial.⁵ To our knowledge, this is the first modelling study to enable individuals to directly link their individual colorectal cancer risk to their predicted benefits and harms of colorectal cancer screening. Individuals can do this through determining their colorectal cancer risk using a calculator, such as the QCancer calculator,⁴¹ incorporating information on age, sex, ethnicity, and other risk factors. The QCancer (10 year) calculator performed better than other prediction tools when externally validated against the UK Biobank cohort. However, like the other colorectal cancer risk prediction tools, it is far from perfect and, with an area under the curve of 0.67 in men and 0.65 in women, it poorly discriminates between those at a lower and those at a higher risk, which may lead to misclassification of individual baseline colorectal cancer risk.^{16 53}

Our study has several limitations. First, MISCAN-Colon could not replicate the sex-specific differences in colorectal cancer incidence and mortality reduction as observed in the NORCCAP sigmoidoscopy screening trials¹⁶ (appendix 3 (table 1)), despite the sex-specific adjustments to the model.⁵ On the one hand, it may be that the dwell time of adenomas differs between men and women, potentially because the proportion of cancers in the proximal colon is higher in women. In MISCAN-Colon we did not assume sex-specific or localisation-specific differences in duration because of

insufficient information from clinical studies or autopsy studies. On the other hand, the observed difference in screening effectiveness of sigmoidoscopy between men and women is higher in the NORCCAP trial than what was observed in the other sigmoidoscopy studies.¹⁶

Second, at the request of the guideline panel, we modelled only four screening strategies, and, for FIT, applied only one cut-off value (20 µg haemoglobin/g faeces). Applying lower or higher FIT cut-off values may result in higher or lower colorectal cancer screening effectiveness. In view of the results of diagnostic studies, there are also uncertainties regarding the additional benefit of using FIT annually instead of biennially.⁵⁴

Third, it remains unknown whether differences in colorectal cancer risk among the population are caused by variations in the number of adenomas, a faster adenoma progression to malignancy, or a combination of the two. These variations may exist between men and women, different ethnicities, different levels of genetic predisposition or different environments. For this analysis, we assumed that differences in adenoma incidence cause differences in colorectal cancer risk.

Fourth, in MISCAN-Colon we assumed that all cancers developed from precursors via a common pathway. In the model description we refer to this as the adenoma carcinoma pathway with adenoma being the precursor lesions. However, recent evidence suggested that three distinct cancer pathways are relevant: about 60-70% of cancers develop via the conventional adenoma carcinoma sequence, 20-30% via the serrated polyp pathway, and 3% via the Lynch pathway.⁵⁵ In the model, we calibrated the average time it takes for a precursor to develop into colorectal cancer. Therefore, all precursor types are included in the modelled mix of slow and rapid progressing lesions. Modelling one common pathway may have consequences for the modelled results. For instance, we may overestimate the effectiveness of FIT and sigmoidoscopy compared with colonoscopy. Evidence is accumulating that FIT might be less sensitive for serrated polyps, and these polyps are usually located in the right side of the colon.^{39 40 47 56} These polyps may have higher malignant potential than conventional adenomas. However, evidence for the malignant potential of the precursors from the distinct pathways is not yet decisive.

Fifth, this project focuses on the individual's perspective rather than on the perspective of public health professionals deciding on population-based screening programmes, which was the reason for assuming 100% adherence to screening and follow-up for all screening options. In population-based screening programmes, adherence rates of the various screening options can differ widely across countries.⁵⁷⁻⁵⁹ When public health professionals make decisions about population-based screening programmes, they should also include evidence on the country-specific adherence rate to determine which screening option is most suitable. In addition, cost effectiveness analysis should be performed. For public

health professionals, a message from this study still may lie in the finding that benefits of screening do not differ much between the screening options.

Sixth, the estimated reduction in all-cause mortality is not observed in large randomised trials of gFOBT and sigmoidoscopy screening, which have not shown a significant reduction of all-cause mortality with screening. The *BMJ* Rapid Recommendations panel did not regard all-cause mortality estimates from the model as clinically relevant when making their recommendations.

Finally, we did not model probability bands. An important strength of this study is the large number of simulations we have performed, with different screening strategies, background colorectal cancer risks, age groups, and sex. The drawback is that to obtain probability bands for all simulations in this article, would require 840 000 simulations, which is too computationally expensive.

Policy implications and conclusions

Notwithstanding the limitations, this modelling study addresses an important gap in current knowledge on colorectal cancer screening. There is insufficient evidence from clinical studies to determine which screening modality is most effective.¹³ Currently, three large randomised trials are under way to assess the comparative effectiveness of colonoscopy and FIT,⁶⁰ and one on sigmoidoscopy versus FIT.⁵³ Our results indicate that the difference in colorectal cancer mortality reduction between screening modalities is substantially smaller than the differences in colorectal cancer mortality between screening and no screening. To achieve sufficient power to demonstrate these differences, the current colorectal cancer screening trials would require very large sample sizes. It is therefore unlikely that a comparison of all evaluated strategies will ever become available from randomised trials. Comparing screening modalities stratified by baseline colorectal cancer risk, age, and sex is even more complicated. In these cases, clinicians and patients must make choices on the basis of the best available evidence, even if it is of low certainty.

Our belief is that our modelling results are generalisable to individuals across the Western world. Although there may be some differences in life expectancy, age-specific colorectal cancer incidence, colorectal cancer stage distribution, and colorectal cancer survival compared with Norway, we expect that these differences do not significantly affect relative differences between screening modalities.

This study, together with the other publications in this *BMJ* Rapid Recommendations cluster, supports patients and physicians in the process of shared decision making by quantifying screening benefits, harms, and burdens on an individual level. For example, based on a certain colorectal cancer risk threshold, some low risk individuals may conclude that the undesirable consequences of screening outweigh the desirable consequences. Evaluating the modelling results, we predicted that lifetime follow-up of screened

individuals resulted in different estimates of screening effectiveness compared with 15 year follow-up. This additional finding encourages researchers to continue the follow-up in their randomised cohorts to evaluate longer term benefits of screening.

In conclusion, MISCAN-Colon predicted that all screening modalities reduce colorectal cancer mortality during a 15 year follow-up period, regardless of colorectal cancer risk, age, and sex. A single colonoscopy may be the most effective screening modality in preventing colorectal cancer incidence during 15 years follow-up. Colonoscopy screening is also associated with the highest risk for complications, but overall complication risks are low for colorectal screening. These results will contribute to risk-based colorectal cancer screening recommendations in this *BMJ* Rapid Recommendation project.¹⁵

AUTHOR AFFILIATIONS

¹Erasmus MC, University Medical Center Rotterdam, Department of Public Health, Rotterdam, the Netherlands

²Centre for Health Services Research, School of Population and Global Health, The University of Western Australia, Perth, Western Australia, Australia

³Clinical Effectiveness Research Group, Department of Transplantation Medicine, Oslo University Hospital, Oslo, Norway

⁴Clinical Effectiveness Research Group, Institute of Health and Society, University of Oslo, Oslo, Norway

⁵Frontier Science Foundation, Boston, MA, USA

⁶Department of Health Research Methods, Evidence and Impact, McMaster University, ON, Canada

⁷Institute of Health and Society, Faculty of Medicine, University of Oslo, Oslo, Norway

⁸Department of Clinical Epidemiology, Leibniz Institute for Prevention Research and Epidemiology - BIPS, Bremen, Germany

⁹Faculty of Human and Health Sciences, University of Bremen, Bremen, Germany

The study has used data from the Cancer Registry of Norway. The interpretation and reporting of these data are the sole responsibility of the authors, and no endorsement by the Cancer Registry of Norway is intended nor should be inferred.

Contributors: MBr, GG, ILV, MBu, and LMH conceived the study idea and assessed the quality of the body of evidence. MBu conducted data analyses and wrote the first draft of the manuscript. All co-authors interpreted the data analysis and critically revised the manuscript. MBu is the guarantor. The corresponding author attests that all listed authors meet authorship criteria and that no others meeting the criteria have been omitted

Funding: This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

Competing interests: All authors have completed the ICMJE uniform disclosure form and declare no support from any organisation for the submitted work. GG is a member of the GRADE working group. ILV has received funding for monitoring, evaluation and cost effectiveness analysis of different colorectal cancer screening strategies, from the US National Cancer Institute, European Commission, ZonMw, The Dutch Cancer Society, Netherlands Institute of Public Health, and NHMRC Australia. Her institution has received payments for her lectures and educational events on colorectal cancer screening for Erasmus MC and Utrecht Medical Center. She has contributed in writing a guidebook on colorectal cancer screening for the International Agency for Research in Cancer (IARC) and has received travel expenses and per diem from IARC. She has not directly contributed to other guidelines, but her work has served as input to the deliberations of other guideline-issuing committees such as USPSTF and ACS. MBr is an author on the NORCCAP trial, one of the trigger trials of this guideline with long term follow-up of sigmoidoscopy screening. He is the principal investigator on the ongoing Nordic trial comparing no screening with colonoscopy screening. No results have been published from this study yet. He has taken part in writing a guidebook on colorectal cancer screening for

the IARC 2017-18. There are no other relationships or activities that could appear to have influenced the submitted work.

Ethical approval: Not required.

Data sharing: No additional data available.

MBU and ILV affirm that the manuscript is an honest, accurate, and transparent account of the study being reported; no important aspects of the study have been omitted.

This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

- Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, Jemal A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin* 2018;68:394-424. doi:10.3322/caac.21492
- Holme Ø, Bretthauer M, Frøheim A, Odgaard-Jensen J, Hoff G. Flexible sigmoidoscopy versus faecal occult blood testing for colorectal cancer screening in asymptomatic individuals. *Cochrane Database Syst Rev* 2013;(9):CD009259. doi:10.1002/14651858.CD009259.pub2
- Atkin W, Wooldrage K, Parkin DM, et al. Long term effects of once-only flexible sigmoidoscopy screening after 17 years of follow-up: the UK Flexible Sigmoidoscopy Screening randomised controlled trial. *Lancet* 2017;389:1299-311. doi:10.1016/S0140-6736(17)30396-3
- Segnan N, Armaroli P, Bonelli L, et al, SCORE Working Group. Once-only sigmoidoscopy in colorectal cancer screening: follow-up findings of the Italian Randomized Controlled Trial-SCORE. *J Natl Cancer Inst* 2011;103:1310-22. doi:10.1093/jnci/djr284
- Holme Ø, Løberg M, Kalager M, et al, NORCCAP Study Group. Long-term effectiveness of sigmoidoscopy screening on colorectal cancer incidence and mortality in women and men: a randomized trial. *Ann Intern Med* 2018;168:775-82. doi:10.7326/M17-1441
- Hardcastle JD, Chamberlain JO, Robinson MH, et al. Randomised controlled trial of faecal-occult-blood screening for colorectal cancer. *Lancet* 1996;348:1472-7. doi:10.1016/S0140-6736(96)03386-7
- Scholefield JH, Moss SM, Mangham CM, Whynes DK, Hardcastle JD. Nottingham trial of faecal occult blood testing for colorectal cancer: a 20-year follow-up. *Gut* 2012;61:1036-40. doi:10.1136/gutjnl-2011-300774
- Jørgensen OD, Kronborg O, Fenger C. A randomised study of screening for colorectal cancer using faecal occult blood testing: results after 13 years and seven biennial screening rounds. *Gut* 2002;50:29-32. doi:10.1136/gut.50.1.29
- Lindholm E, Brevinge H, Haglund E. Survival benefit in a randomized clinical trial of faecal occult blood screening for colorectal cancer. *Br J Surg* 2008;95:1029-36. doi:10.1002/bjs.6136
- Mandel JS, Church TR, Bond JH, et al. The effect of fecal occult-blood screening on the incidence of colorectal cancer. *N Engl J Med* 2000;343:1603-7. doi:10.1056/NEJM200011303432203
- Schoen RE, Pinsky PF, Weissfeld JL, et al, PLCO Project Team. Colorectal-cancer incidence and mortality with screening flexible sigmoidoscopy. *N Engl J Med* 2012;366:2345-57. doi:10.1056/NEJMoa1114635
- Shapiro JA, Bobo JK, Church TR, et al. A comparison of fecal immunochemical and high-sensitivity guaiac tests for colorectal cancer screening. *Am J Gastroenterol* 2017;112:1728-35. doi:10.1038/ajg.2017.285
- Lauby-Secretan B, Vilahur N, Bianchini F, Guha N, Straif K, International Agency for Research on Cancer Handbook Working Group. The IARC perspective on colorectal cancer screening. *N Engl J Med* 2018;378:1734-40. doi:10.1056/NEJMs1714643
- Miller EA, Pinsky PF, Schoen RE, Prorok PC, Church TR. Effect of flexible sigmoidoscopy screening on colorectal cancer incidence and mortality: long-term follow-up of the randomised US PLCO cancer screening trial. *Lancet Gastroenterol Hepatol* 2019;4:101-10. doi:10.1016/S2468-1253(18)30358-3
- Helsing LM, Vandvik PO, Jodal HC, et al. Colorectal cancer screening with faecal immunochemical testing, sigmoidoscopy or colonoscopy: a clinical practice guideline. *BMJ* 2019;367:l5515. doi:10.1136/bmj.l5515
- Jodal HC, Helsing LM, Anderson JC, et al. Colorectal cancer screening with faecal testing, sigmoidoscopy or colonoscopy: a systematic review and network meta-analysis. *BMJ Open* 2019;0:e032773. doi:10.1136/bmjopen-2019-032773
- Loeve FBR, van Ballegooijen M, van Oortmarssen GJ, Habbema JDF. Final Report MISCANCOLON, Microsimulation Model for Colorectal Cancer: Report to the National Cancer Institute Project. No. NO1-CN55186. Department of Public Health, Erasmus University, 1998.
- van Hees F, Zauber AG, van Veldhuizen H, et al. The value of models in informing resource allocation in colorectal cancer screening: the case of The Netherlands. *Gut* 2015;64:1985-97. doi:10.1136/gutjnl-2015-309316
- Blatt L. Polyps of the colon and rectum: incidence and distribution. *Dis Colon Rectum* 1961;4:277-82. doi:10.1007/BF02616606
- Arminksi TC, McLean DW. Incidence and distribution of adenomatous polyps of the colon and rectum based on 1,000 autopsy examinations. *Dis Colon Rectum* 1964;7:249-61. doi:10.1007/BF02630528
- Bombi JA. Polyps of the colon in Barcelona, Spain. An autopsy study. *Cancer* 1988;61:1472-6. doi:10.1002/1097-0142(19880401)61:7<1472::AID-CNCR2820610734>3.0.CO;2-E
- Chapman I. Adenomatous polypi of large intestine: incidence and distribution. *Ann Surg* 1963;157:223-6. doi:10.1097/00000658-196302000-00007
- Clark JC, Collan Y, Eide TJ, et al. Prevalence of polyps in an autopsy series from areas with varying incidence of large-bowel cancer. *Int J Cancer* 1985;36:179-86. doi:10.1002/ijc.2910360209
- Jass JR, Young PJ, Robinson EM. Predictors of presence, multiplicity, size and dysplasia of colorectal adenomas. A necropsy study in New Zealand. *Gut* 1992;33:1508-14. doi:10.1136/gut.33.11.1508
- Johannsen LGK, Mømsen O, Jacobsen NO. Polyps of the large intestine in Aarhus, Denmark. An autopsy study. *Scand J Gastroenterol* 1989;24:799-806. doi:10.3109/00365528909089217
- Rickert RR, Auerbach O, Garfinkel L, Hammond EC, Frasca JM. Adenomatous lesions of the large bowel: an autopsy survey. *Cancer* 1979;43:1847-57. doi:10.1002/1097-0142(197905)43:5<1847::AID-CNCR2820430538>3.0.CO;2-L
- National Cancer Institute. Surveillance, Epidemiology, and End Results (SEER) Program. SEER* Stat Databases: Incidence—SEER 9 Regs Limited-Use, Nov 2002 Sub (1973-2002). 2003.
- Vatn MH, Stalsberg H. The prevalence of polyps of the large intestine in Oslo: an autopsy study. *Cancer* 1982;49:819-25. doi:10.1002/1097-0142(19820215)49:4<819::AID-CNCR2820490435>3.0.CO;2-D
- Williams AR, Balasooriya BA, Day DW. Polyps and cancer of the large bowel: a necropsy study in Liverpool. *Gut* 1982;23:835-42. doi:10.1136/gut.23.10.835
- Lansdorp-Vogelaar I, van Ballegooijen M, Boer R, Zauber A, Habbema JD. A novel hypothesis on the sensitivity of the fecal occult blood test: Results of a joint analysis of 3 randomized controlled trials. *Cancer* 2009;115:2410-9. doi:10.1002/cncr.24256
- Rutter CM, Knudsen AB, Marsh TL, et al. Validation of models used to inform colorectal cancer screening guidelines: accuracy and implications. *Med Decis Making* 2016;36:604-14. doi:10.1177/0272989X15622642
- Atkin WS, Edwards R, Kralj-Hans I, et al, UK Flexible Sigmoidoscopy Trial Investigators. Once-only flexible sigmoidoscopy screening in prevention of colorectal cancer: a multicentre randomised controlled trial. *Lancet* 2010;375:1624-33. doi:10.1016/S0140-6736(10)60551-X
- Hardcastle JD, Chamberlain JO, Robinson MHE, et al. Randomised controlled trial of faecal-occult-blood screening for colorectal cancer. *Lancet* 1996;348:1472-7. doi:10.1016/S0140-6736(96)03386-7
- Scholefield JH, Moss S, Sufi F, Mangham CM, Hardcastle JD. Effect of faecal occult blood screening on mortality from colorectal cancer: results from a randomised controlled trial. *Gut* 2002;50:840-4. doi:10.1136/gut.50.6.840
- Mandel JS, Church TR, Ederer F, Bond JH. Colorectal cancer mortality: effectiveness of biennial screening for fecal occult blood. *J Natl Cancer Inst* 1999;91:434-7. doi:10.1093/jnci/91.5.434
- Norwegian Cancer Registry. 2018 <https://www.kreftregisteret.no/en/>.
- National Institute for Public Health and the Environment. National monitoring of the colorectal cancer screening programme 2014. 2017. https://www.rivm.nl/en/Documents_and_publications/Common_and_Present/Publications/Disease_prevention_and_healthcare/bowel_cancer/National_Monitoring_of_the_Colorectal_Cancer_Screening_Programme.
- Hippisley-Cox J, Coupland C. Development and validation of risk prediction algorithms to estimate future risk of common cancers in men and women: prospective cohort study. *BMJ Open* 2015;5:e007825. doi:10.1136/bmjopen-2015-007825
- Heigh RI, Yab TC, Taylor WR, et al. Detection of colorectal serrated polyps by stool DNA testing: comparison with fecal immunochemical testing for occult blood (FIT). *PLoS One* 2014;9:e85659. doi:10.1371/journal.pone.0085659
- Rex DK, Ahnen DJ, Baron JA, et al. Serrated lesions of the colorectum: review and recommendations from an expert panel. *Am J Gastroenterol* 2012;107:1315-29, quiz 1314, 1330. doi:10.1038/ajg.2012.161
- ClinRisk. Welcome to the QCancer (15yr, colorectal) risk calculator. 2017. <https://qcancer.org/15yr/colorectal/index.php>.

- 42 Schreuders EH, Ruco A, Rabeneck L, et al. Colorectal cancer screening: a global overview of existing programmes. *Gut* 2015;64:1637-49. doi:10.1136/gutjnl-2014-309086
- 43 van Rijn JC, Reitsma JB, Stoker J, Bossuyt PM, van Deventer SJ, Dekker E. Polyp miss rate determined by tandem colonoscopy: a systematic review. *Am J Gastroenterol* 2006;101:343-50. doi:10.1111/j.1572-0241.2006.00390.x
- 44 Holme Ø, Schoen RE, Senore C, et al. Effectiveness of flexible sigmoidoscopy screening in men and women and different age groups: pooled analysis of randomised trials. *BMJ* 2017;356:i6673. doi:10.1136/bmj.i6673
- 45 van Hees F, Zaubler AG, Klabunde CN, Goede SL, Lansdorp-Vogelaar I, van Ballegooijen M. The appropriateness of more intensive colonoscopy screening than recommended in Medicare beneficiaries: a modeling study. *JAMA Intern Med* 2014;174:1568-76. doi:10.1001/jamainternmed.2014.3889
- 46 Buskermolen M, Gini A, Naber SK, Toes-Zoutendijk E, de Koning HJ, Lansdorp-Vogelaar I. Modeling in colorectal cancer screening: assessing external and predictive validity of MISCAN-Colon microsimulation model using NORCCAP trial results. *Med Decis Making* 2018;38:917-29. doi:10.1177/0272989X18806497
- 47 Imperiale TF, Ransohoff DF, Itzkowitz SH, et al. Multitarget stool DNA testing for colorectal-cancer screening. *N Engl J Med* 2014;370:1287-97. doi:10.1056/NEJMoa1311194
- 48 Warren JL, Klabunde CN, Mariotto AB, et al. Adverse events after outpatient colonoscopy in the Medicare population. *Ann Intern Med* 2009;150:849-57. W152. doi:10.7326/0003-4819-150-12-200906160-00008
- 49 Gatto NM, Frucht H, Sundararajan V, Jacobson JS, Grann VR, Neugut AI. Risk of perforation after colonoscopy and sigmoidoscopy: a population-based study. *J Natl Cancer Inst* 2003;95:230-6. doi:10.1093/jnci/95.3.230
- 50 Hassan C, Quintero E, Dumonceau JM, et al, European Society of Gastrointestinal Endoscopy. Post-polypectomy colonoscopy surveillance: European Society of Gastrointestinal Endoscopy (ESGE) Guideline. *Endoscopy* 2013;45:842-51. doi:10.1055/s-0033-1344548
- 51 Guyatt GH, Oxman AD, Vist GE, et al, GRADE Working Group. GRADE: an emerging consensus on rating quality of evidence and strength of recommendations. *BMJ* 2008;336:924-6. doi:10.1136/bmj.39489.470347.AD
- 52 Ventura L, Mantellini P, Grazzini G, et al. The impact of immunochemical faecal occult blood testing on colorectal cancer incidence. *Dig Liver Dis* 2014;46:82-6. doi:10.1016/j.dld.2013.07.017
- 53 Usher-Smith JA, Harshfield A, Saunders CL, et al. External validation of risk prediction models for incident colorectal cancer using UK Biobank. *Br J Cancer* 2018;118:750-9. doi:10.1038/bjc.2017.463
- 54 van Roon AH, Goede SL, van Ballegooijen M, et al. Random comparison of repeated faecal immunochemical testing at different intervals for population-based colorectal cancer screening. *Gut* 2013;62:409-15. doi:10.1136/gutjnl-2011-301583
- 55 East JE, Atkin WS, Bateman AC, et al. British Society of Gastroenterology position statement on serrated polyps in the colon and rectum. *Gut* 2017;66:1181-96. doi:10.1136/gutjnl-2017-314005
- 56 Chang LC, Shun CT, Hsu WF, et al. Fecal immunochemical test detects sessile serrated adenomas and polyps with a low level of sensitivity. *Clin Gastroenterol Hepatol* 2017;15:872-879.e1. doi:10.1016/j.cgh.2016.07.029
- 57 Hol L, van Leerdam ME, van Ballegooijen M, et al. Screening for colorectal cancer: randomised trial comparing guaiac-based and immunochemical faecal occult blood testing and flexible sigmoidoscopy. *Gut* 2010;59:62-8. doi:10.1136/gut.2009.177089
- 58 Segnan N, Senore C, Andreoni B, et al, SCORE3 Working Group-Italy. Comparing attendance and detection rate of colonoscopy with sigmoidoscopy and FIT for colorectal cancer screening. *Gastroenterology* 2007;132:2304-12. doi:10.1053/j.gastro.2007.03.030
- 59 Liang PS, Wheat CL, Abhat A, et al. Adherence to competing strategies for colorectal cancer screening over 3 years. *Am J Gastroenterol* 2016;111:105-14. doi:10.1038/ajg.2015.367
- 60 Robertson DJ, Kaminski MF, Bretthauer M. Effectiveness, training and quality assurance of colonoscopy screening for colorectal cancer. *Gut* 2015;64:982-90. doi:10.1136/gutjnl-2014-308076

Appendices. 1: MISCAN-Colon model description.
 2: Calibration of MISCAN-Colon to Norwegian setting.
 3: Validation of MISCAN-Colon against NORCCAP study.
 4: MISCAN-Colon predictions with 15 year follow-up stratified for colorectal cancer risk, age, and gender.
 5: MISCAN-Colon predictions with lifetime follow-up stratified for colorectal cancer risk, age, and gender.