

Diet quality in childhood: the Generation R Study

Nguyen AN*, van der Velde LA*, Schoufour JD, Geelen A., Jaddoe VWV, Franco OH, Voortman T.

Diet quality in childhood: the Generation R Study.
European journal of nutrition. 2019;58(3): 1259-1269.

ABSTRACT

Objective: We aimed to evaluate diet quality of 8-year-old children in the Netherlands, to identify sociodemographic and lifestyle correlates of child diet quality, and to examine tracking of diet quality from early to mid-childhood.

Methods: For 4,733 children participating in a population-based cohort, we assessed dietary intake using a validated food-frequency questionnaire at a median age of 8.1 years (interquartile range:8.0-8.2)(2011-2014). Based on dietary guidelines, we developed and validated a food-based diet quality score for children consisting of ten components (score 0-10): sufficient intake of vegetables; fruit; whole-grains; fish; legumes; nuts; dairy; oils & soft fats; and low intake of sugar-containing-beverages; and high-fat and processed meat.

Results: We observed a mean (\pm SD) diet quality score of 4.5 (\pm 1.2) out of a maximum of 10. On average, intake of legumes, nuts, and oils or soft fats was below recommendations, whereas intake of sugar-containing beverages and high-fat or processed meat was higher than recommended. The main factors associated with higher diet quality were higher maternal educational level ($\beta=0.29$, 95%CI:0.21,0.37 versus low education), higher household income ($\beta=0.15$, 95%CI:0.05,0.025), no maternal smoking ($\beta=0.13$, 95%CI:0.25,0.02 versus current smoking), and less screen time ($\beta=0.31$, 95%CI:-0.38,-0.24)– all independent of each other. For children with available dietary data at age 1 year ($n=2,608$), we observed weak tracking of diet quality from early to mid-childhood (Pearson's $r=0.19$, $k=0.11$ for extreme quartiles).

Conclusion: Overall diet quality of 8-year-old children is not conform dietary guidelines, especially for children having more screen time, children of lower educated and smoking mothers, and from lower-income households.

INTRODUCTION

A healthy diet during childhood is important for healthy growth and development ¹, and may contribute to the prevention of obesity and chronic diseases later in life ^{2,3}. Furthermore, dietary habits in childhood have been shown to track over time and are an important predictor of diet quality in adulthood ⁴. Therefore, it is important to examine children's dietary intake, to identify potential gaps between their actual and recommended intake, and to study determinants of diet, in order to develop targeted interventions focusing on groups with a high risk of poor dietary habits early in life.

Given the complexity of the human diet and the strong interactions between intake of different foods and nutrients, measuring overall dietary patterns is recommended as a complementary approach to measuring the intake of only single foods or nutrients ^{5,6}. One way to study overall diet is by predefined diet quality scores, which are usually based on dietary guidelines ^{5,7}. Although the use of diet quality indices in children has increased over the past years, Marshall et al. suggested in their systematic review that more prospective cohort studies evaluating diet quality in children and its impact on health are needed ⁸. A few studies from different countries including the UK, Brazil, and the US, assessed diet quality among school-age children ⁹⁻¹¹. However, because dietary habits and guidelines may differ between countries and cultures, it is important to use a diet quality score that assesses recommendations specific for the study population ¹². Previously, we developed a food-based diet quality score specifically for preschool children ¹³. However, to date, no diet quality score is available for school-age children in the Netherlands, and factors related to diet quality have not been studied in this age category. Furthermore, previous studies reported tracking of diet from mid-childhood or adolescence to adulthood ^{2,14}, but information on changes in diet quality from early childhood to mid-childhood is scarce. This information is needed in order to establish whether dietary interventions could be efficient early in life ².

Therefore, we aimed to evaluate overall diet quality of 8-year-old children participating in a large population-based cohort in the Netherlands. For this aim we developed a new food-based diet quality score based on current Dutch dietary guidelines ¹⁵, and we assessed the construct validity of this new diet quality score. This score can be applied in future studies to evaluate diet quality, to investigate associations between diet quality and health, and to support future dietary advice and interventions. Furthermore, we aimed to identify which parental and child sociodemographic and lifestyle factors, such as educational level, physical activity, and screen time correlate with diet quality of children and we aimed to investigate associations between diet quality at the ages of 1 and 8 years. This information can help to identify the best target groups and time frame for interventions to improve diet quality in children.

METHODS

Study design and study population

This study was embedded in the Generation R Study, a multiethnic population-based prospective cohort from fetal life onward in Rotterdam, the Netherlands. Women living in the city of Rotterdam were enrolled during pregnancy. Children participating in this study were born between April 2002 and January 2006. The study was approved by the Medical Ethics Committee of Erasmus Medical Center and written informed consent was obtained from parents of all participating children¹⁶. A dietary questionnaire was sent to mothers who provided consent for follow-up when their child was around the age of 8 years ($n=7,662$). The questionnaire was returned for 4,787 children (62.5%). After exclusion of subjects with invalid dietary data ($n=54$), defined as a reported energy intake below 650 ($n=47$) or above 3,700 kcal/d ($n=7$), valid dietary data were available for 4,733 children (**Figure 2.4.1**). Of all children with dietary data at the age of 8 years, 2,608 children also had dietary data available at their age of 1 year.




Figure 2.4.1. Flowchart of participants included in the study

Dietary assessment

Dietary intake was assessed at a median age of 8.1 years (interquartile range (IQR): 8.0-8.2) using a validated semi-quantitative food-frequency questionnaire (FFQ)¹⁷. The FFQ was completed by the parents of the child, using the last four weeks as reference period. As explained in detail previously¹⁷, the FFQ was developed based on results from a national food consumption survey in the Netherlands¹⁸, which resulted in the selection of 71 food items relevant for the energy intake of 2 to 12-year-old children. Questions concerned the frequency of consumption and portion sizes of these foods,

and for 27 food items additional questions were included about specific types or brands and preparation methods. Portion sizes were inquired for in natural units, household units, or grams; and parents were asked to measure the volume of glasses and cups regularly used by their child. Dietary intake data were cleaned and corrected based on detection of missing values, outliers of quantities, and inconsistencies using standardized algorithms developed for the FFQ¹⁷. Information on frequencies, types, and portion sizes was converted into grams of individual food items per day based on standard Dutch portion sizes, using SAS VoVris (Vovris V2.4, TNO, 1999-2006). Energy and nutrient intakes from foods were calculated using data from the Dutch Food Composition Table (NEVO 2001) with SAS Veves (Veves V2.2, TNO, 1999-2003).

The FFQ has been validated for energy intake among 4 to 6-year-old Dutch children (n=30) using the doubly labelled water method¹⁷. The Pearson's correlation between energy intake as estimated from the FFQ and energy expenditure measured with doubly labelled water was 0.62, indicating a reasonable capacity to rank subjects with respect to energy intake. Furthermore, no relevant intake-related bias was observed in the Bland-Altman plot. These findings indicate that the FFQ is a valid instrument for the assessment of energy intake in children¹⁷.

Construction of the diet quality score for school-age children

We constructed a food-based diet quality score based on dietary recommendations for children from the Netherlands Nutrition Center¹⁵, thereby also taking into account the Dutch Guidelines for a Healthy Diet of 2015¹⁹, on which the Nutrition Center recommendations were based. We included ten components (i.e., food groups) in the diet quality score, of which eight were adequacy components (i.e., adequate intake is recommended) and two were moderation components (i.e., low intake is recommended). Only recommended food items were included for the adequacy components, and recommended food items were excluded for the moderation components (**Table 2.4.1**). The components, food items included, and their cut-off values were determined *a priori*, based on the recommendations of the Netherlands Nutrition Center for 8-year-old children¹⁵, and were as follows: fruit (≥ 150 g/d), vegetables (≥ 150 g/d), whole-grains (≥ 90 g/d), fish (≥ 60 g/w), legumes (≥ 84 g/w), nuts (≥ 15 g/d), dairy (≥ 300 g/d), oils and soft or liquid fats (≥ 30 g/d), sugar-containing beverages (≤ 150 g/d) and high-fat and processed meat (≤ 250 g/w) (Table 2.4.1).

Scoring of diet quality was performed by calculating the ratio of reported and recommended intake for each component¹³. These component scores were truncated at 1. For the adequacy components, this resulted in a minimum score of 0 points when these food items were not consumed and a maximum score of 1 point when the amount of the cut-off value or more was consumed. For example, a fruit intake of 60 g/d resulted in a score of 0.4 (60 divided by 150 g/d) for this component. For the moderation compo-

nents, this scoring system was reversed, with higher scores reflecting lower intakes (e.g., a sugar-containing beverages intake of 75 g/d resulted in a score of 0.5 (1-(75 divided by 150 g/d)). Scores of the individual components were summed, resulting in a total score for diet quality ranging from 0 to 10 on a continuous scale, with a higher score indicating a better overall diet quality.

Table 2.4.1. Components, cut-off values and included and excluded food items of the diet quality score

Component	Cut-off value ^a	Foods included in the diet quality score	Foods not included in the diet quality score
Fruit	≥ 150 g/d	Fresh fruit, frozen fruit, dried fruit (up to 20 g/d ^b), canned fruit without added sugar	Fruit juice, dried fruit (>20 g/d), fruit products with added sugar
Vegetables	≥ 150 g/d	Fresh vegetables, frozen vegetables, canned vegetables ^c	-
Whole-grains	≥ 90 g/d	Brown/ whole grain bread or crackers, whole-grain rice, whole-wheat pasta, whole grain breakfast cereals without added sugar	White bread or crackers, white rice, white pasta, breakfast cereals with added sugar
Fish	≥ 60 g/w	Fish, canned fish, shellfish	Fish products containing <70% fish (e.g. battered fish)
Legumes	≥ 84 g/w	Fresh, dried or canned legumes ^c	-
Nuts	≥ 15 g/d	Nuts, peanuts, peanut butter	Coated nuts
Dairy	≥ 300 g/d	Unsweetened, skimmed and semi-skimmed milk, yoghurt, or quark; dairy products without added sugar; buttermilk; low-fat cheese	Full-fat milk, yoghurt, or quark, dairy products with added sugar, full-fat cheese, whipped cream, ice cream
Oils and soft or liquid margarines	≥ 30 g/d	Vegetable oils, soft margarine (≤ 30% saturated fat of total fat), liquid cooking and frying fats	Hard margarine (>30% saturated fat of total fat), hard cooking and frying fats, butter
Sugar-containing beverages	≤ 150 g/d ^d	Soft drinks, fruit juice, lemonade, fruit juice concentrates	Milk based sugar-containing beverages ^e
High-fat and processed meat	≤ 250 g/w	Processed and/or high fat meat (>5% saturated fat)	Unprocessed low fat meat (≤5% saturated fat)

^aBased on the recommendations of the Netherlands Nutrition Center and Health Council of the Netherlands ^{15,19}.

^bAccording to the guidelines, a maximum of 20 g/d of dried fruit was included in the fruit component.

^cCanned products with added salt or sugar were not excluded, as no distinction was made in the FFQ.

^dNo quantitative recommendation was available for sugar-containing beverages, we chose a cut-off of 1 glass/d.

^eAs milk-based beverages with added sugar can provide valuable nutrients and have a greater satiating effect compared to sweetened drinks containing sugar only, these were not included in the sugar-containing beverages component.

g/d= gram/ day; g/w= gram/ week.

Diet quality in infancy

To assess tracking of diet quality from infancy to mid-childhood, we used a previously defined diet quality score for infants. As described in detail elsewhere ¹³, information on dietary intake at the age of 1 year was collected with a semi-quantitative FFQ, which

was developed specifically for this age group. This diet quality score consisted of ten components: vegetables; fruit; bread and cereals; rice, pasta, potatoes, and legumes; dairy; meat; fish; oils and fats; candy and snacks; and sugar sweetened beverages. The scoring system for this diet quality score was similar to that of the diet quality score for 8-year-old children. The score ranged from 0 to 10 on a continuous scale with higher scores reflecting better adherence to dietary guidelines¹³.

Assessments of sociodemographic and lifestyle factors

Several sociodemographic and lifestyle factors were assessed for both the children and their parents. Information on date of birth and sex of the child was obtained from medical records and hospital registries. Ethnicity of the child was based on the country of birth of the parents, which was obtained with questionnaires at enrollment. If both parents were born in the Netherlands, the child was considered to have a Dutch ethnic background. If one parent was born outside of the Netherlands, the country of birth of that parent determined the child's ethnicity. If both parents were born abroad, the country of birth of the mother determined the ethnicity of the child^{16,20}. Ethnicity was categorized according to the largest ethnic groups in our study population, which were Dutch, Moroccan, Turkish, Surinamese and Antillean, other Western, and other non-Western¹³.

During follow-up visits of the participants to our research center at a median ages of 6.0 years (IQR 5.9-6.2) and 9.7 years (IQR 9.6-9.9), we measured several child and maternal factors. As most measurements at these time points were strongly correlated, we used measurements taken at age 9.7 years for the main analyses in the current study, as this age was closest to the age of our dietary assessment at age 8.1 years. Child's height and weight were measured to calculate their body mass index (BMI) (kg/m²), which was categorized into 'underweight', 'normal weight', or 'overweight' according to the Cole-criteria²¹. Questionnaires were used to assess time spent playing sports (i.e., any organized sports outside school hours), which was categorized into <2, 2 to 4, or ≥4 hours per week, and time spent watching television and/or on the computer (screen time), which was dichotomized into <2 or ≥2 hours per day²². At the same visits, we measured mothers' height and weight to calculate BMI, which we categorized into 'underweight' (<18.5), 'normal weight' (≥18.5 to 25), or 'overweight' (≥25)²¹. Information on other parental factors was assessed with questionnaires. Maternal smoking habits were categorized into: 'never smoker', 'past smoker', or 'current smoker'. Maternal educational level was dichotomized into 'no higher education' or 'higher education', with higher education defined as completed higher vocational training or more. Net household income was dichotomized into <2800 or ≥2800 euros per month²³.

Statistical analysis

Child and parental characteristics were described as median (IQR) for continuous variables or percentage for categorical variables. Total diet quality score was described as mean with standard deviation (SD) and as percentage of children with a maximum score. Component scores and intake per component were described as median (IQR) and as percentage of children with a maximum score. Pearson's correlations were used to assess correlations between the individual components of the diet quality score.

Linear regression models were used to identify sociodemographic and lifestyle correlates of diet quality. In these models, we examined children's age, sex, ethnicity, BMI, physical activity, and screen time; and maternal age, BMI, marital status, educational level, and smoking habits; and household income. The basic model was adjusted for total energy intake only; the multivariable model was additionally adjusted for all other sociodemographic and lifestyle variables that were examined in order to assess whether they were independent of each other. Associations of the diet quality score with intake of nutrients associated with a healthy diet were examined in order to assess the construct validity of the diet score (i.e., the degree to which the diet quality score measures a healthy diet)²⁴. These associations were evaluated using Pearson's correlations and partial Pearson's correlations, controlling for energy intake.

Pearson's correlations were also used to assess the association between diet quality at age 1 year and 8 years. Tracking of diet quality score from age 1 year to age 8 years was assessed by determining to which extent children maintained their rank in the categories 'lowest 25%', 'middle 50%', or 'highest 25%'. For this, a 3x3 matrix was constructed and a linear weighted Kappa statistic (k) was computed²⁵, with $k < 0$ indicating poor agreement, and $k 0.81-1.0$ indicating almost perfect agreement²⁶.

Because the FFQ was developed and validated for a Dutch population, sensitivity analyses were performed among children with a Dutch ethnic background only (n=3,143). As non-response analysis, descriptive characteristics of children with valid dietary data (n=4,733) were compared to children with missing dietary data but who were eligible for dietary assessment (i.e., those who still participated in the study at the age of 8 years and who received the FFQ) (n=2,929). To reduce potential bias associated with missing data in our study, multiple imputation of missing data on sociodemographic and lifestyle factors was performed and 10 independent datasets were created. Because similar effect estimates were found in analyses with imputed and unimputed data, pooled results after the multiple imputation were presented. All statistical analyses were performed using SPSS version 21.0 (IBM Corp., 2012, Armonk, NY). A two-sided P-value of 0.05 was considered statistically significant.

RESULTS

Subject characteristics

Characteristics of the children and their parents are described in **Table 2.4.2**. The majority of children had a Dutch ethnic background (66.4%). At the 9-year visit, median BMI of the children was 16.9 (IQR 15.7-18.4), with 80.7% of children having a normal weight, 7.2% underweight, and 12.1% overweight. More than half (51.6%) of the children had a screen time of ≥ 2 hours per day. Median BMI of the mothers was 24.5 (IQR 22.3-27.5), with 55.1% of mothers having a normal weight and 44.1% being overweight. The majority of mothers was highly educated (62.8%) and had never smoked (52.9%).

Characteristics were similar before and after multiple imputation. Of the 7,662 children whose parents received the FFQ, children with missing dietary data ($n=2,929$) more often had a non-Dutch ethnic background and their mothers were on average younger, lower educated, had a higher BMI, and a lower household income.

Diet quality

Our diet quality score approximated a normal distribution with a mean (SD) of 4.5 (1.2). None of the children reached the maximum possible diet quality score of 10. Median scores on most diet quality score components were around or below 0.5 out of a possible maximum of 1 (**Table 2.4.3**). For the adequacy components, median intakes of vegetables, legumes, nuts, dairy, and oils or soft or liquid fats were well below the cut-off values in our study population. For example, median daily vegetable intake was 79 g (IQR 49-123), whereas 150 g is recommended, resulting in a median component score of 0.53 (IQR 0.32-0.82) out of 1 for vegetables. Intakes of the two moderation components (sugar-containing beverages and high-fat and processed meat) exceeded the recommended intake in most children, with corresponding low scores. Median sugar-containing beverages intake, for example, was 323 g/d (IQR 180-524), with a median score of 0.0 (IQR 0.0-0.0) and only 12.8% of the children having a score above zero. Components with the highest median scores were whole-grains (1.0 (IQR 0.72-1.0)), fruit (0.74 (IQR 0.51-1.0)), and fish (0.63 (IQR 0.0-1.0)). Correlations between the diet score components ranged from -0.13 to 0.08. The scores were comparable between boys and girls (**Supplemental table 2.4.1**).

Associations between the diet quality score and several nutrients was assessed to examine construct validity. We observed a positive correlation between the diet quality score and the intake of protein, mainly plant protein ($r=0.41$), dietary fiber ($r=0.58$), and $n-3$ fatty acids ($r=0.24$), and inversely correlated with intake of saturated fat, and monosaccharides and disaccharides ($r=-0.11$) (energy-adjusted, all $p < 0.01$). The score was also positively correlated with intake of all of the examined micronutrients (energy-adjusted, $r=0.16$ to 0.55 , all $p < 0.01$) (**Supplemental table 2.4.2**).

Table 2.4.2. Characteristics of study participants and their parents (n=4,733)

Child characteristics	Median (IQR), Percentage
Boy (%)	49.9
Ethnicity (%)	
Dutch	66.4
Other Western	9.4
Moroccan	3.6
Turkish	5.1
Surinamese and Antillean	7.2
Other non-Western	8.4
Age at FFQ (y)	8.1 (8.0-8.2)
BMI (kg/m ²)	16.9 (15.7-18.4)
Underweight (%)	7.2
Normal weight (%)	80.7
Overweight (%)	12.1
Playing sports (h/w)	
<2	31.9
2-4	40.7
≥4	27.4
Screen time (h/d)	
≥2	51.6
Parental characteristics	
Age mother at 9-year visit (y)	42.0 (39.0-44.6)
BMI mother	24.5 (22.3-27.5)
Underweight (%)	0.8
Normal weight (%)	55.1
Overweight (%)	44.1
Marital status mother (%)	
Married/ partner	88.1
Educational level mother (%)	
Higher education	62.8
Smoking mother (%)	
Never smoker	52.9
Past smoker	33.7
Current smoker	13.4
Household income per month (%)	
≥2800 €	67.6

Values are medians (IQR) for continuous variables, and percentages for categorical variables, on the basis of imputed data (n=10 imputations).

IQR= Interquartile range; FFQ=Food-frequency questionnaire; y= year; BMI=Body mass index; kg/m²= kilogram/ meter²; h/w= hours/ week; h/d= hours/ day.

Table 2.4.3. Cut-offs values, actual intakes, and scores of the different diet quality score components

Component	Cut-off values	Unit	Intake (n=4,733)	Score (n=4,733)	% with a maximum score
Fruit	≥ 150	g/d	111 (77-167)	0.74 (0.51-1.00)	29.4
Vegetables	≥ 150	g/d	79 (48-123)	0.53 (0.32-0.82)	16.3
Whole-grains	≥ 90	g/d	98 (65-131)	1.0 (0.72-1.00)	57.4
Fish ^a	≥ 60	g/w	38 (0-83)	0.63 (0.00-1.00)	36.0
Legumes ^a	≥ 84	g/w	18 (0-70)	0.21 (0.00-0.83)	21.1
Nuts ^a	≥ 15	g/d	3 (0-10)	0.20 (0.00-0.64)	10.6
Dairy	≥ 300	g/d	164 (54-298)	0.55 (0.18-0.99)	24.9
Oils and soft or liquid fats	≥ 30	g/d	11 (2-17)	0.37 (0.071-0.57)	2.7
Sugar-containing beverages ^a	≤ 150	g/d	323 (180-524)	0.00 (0.00-0.00)	2.1
High-fat and processed meat ^a	≤ 250	g/w	323 (218-453)	0.00 (0.00-0.13)	0.2

Values are median (IQR) Maximum score per component: 1.

Collinearity between the diet score components: r -0.13 to 0.077.

^aA score of 0 was obtained by 27% of the participants for the fish component, 46.5% for the legumes component, 27.3% for the nuts component, 87.8% for the sugar-containing beverages component, and 68.2% for the meat component.

g/d= gram/ day; g/w= gram/ week; IQR= Interquartile range.

Sociodemographic and lifestyle factors and the diet quality score

Associations between sociodemographic and lifestyle factors and the diet quality score are shown in **Table 2.4.4**. In the multivariable model, children with underweight had a lower diet quality score than children with a normal weight. Children with a screen time of ≥ 2 hours per week had a 0.31 points lower diet quality score (95%CI -0.38; -0.24) than children with a screen time < 2 hours per week, and children who played sports for 2 to 4 hours per week had a 0.10 points higher diet quality (95%CI 0.02; 0.19) than those who played sports < 2 hours per week, although no significant difference was found between playing sports for < 2 hours per week versus ≥ 4 hours per week. In the multivariable model, Moroccan children had a 0.29 points higher diet quality score (95%CI 0.10; 0.48) than children with a Dutch ethnicity, whereas in the basic model, without adjustment for the other factors, Turkish as well as Surinamese and Antillean children had a lower diet quality score than children with a Dutch ethnicity. Children's sex or age at dietary assessment was not associated with the diet quality score in the basic and multivariable model.

Parental socioeconomic status was also associated with children's diet quality: children of higher educated mothers or from households with a higher income had a higher diet quality score (Table 2.4.4). Independent of these socioeconomic factors, children of overweight mothers and children of mothers who were current smokers had a lower diet quality score than children of normal-weight or never-smoking mothers, respectively. Sensitivity analyses among Dutch children only showed similar effect estimates.

Table 2.4.4. Associations between sociodemographic and lifestyle factors and the diet quality score

	Basic model ^a		Multivariable model ^b	
	β (95 % CI)	p-value	β (95 % CI)	p-value
Child characteristics				
Sex				
Boy	<i>Reference</i>			
Girl	-0.01 (-0.04; 0.03)	0.78	-0.03 (-0.10; 0.04)	0.41
Ethnicity				
Dutch	<i>Reference</i>			
Other Western	0.01 (-0.10; 0.13)	0.83	0.03 (-0.08; 0.14)	0.61
Moroccan	0.001 (-0.18; 0.18)	1.00	0.29 (0.10; 0.48)	0.002
Turkish	-0.43 (-0.59; -0.28)	< 0.001	-0.11 (-0.28; 0.06)	0.21
Surinamese and Antillean	-0.21 (-0.34; -0.08)	0.002	0.05 (-0.08; 0.19)	0.47
Other non-Western	-0.11 (-0.23; 0.01)	0.08	0.08 (-0.04; 0.21)	0.19
Age at FFQ (y)	-0.02 (-0.16; 0.12)	0.77	0.06 (-0.08; 0.20)	0.37
Energy intake (100 kcal/ d)	0.10 (0.09; 0.10)	<0.001	0.10 (0.09; 0.11)	<0.001
BMI				
Normal weight	<i>Reference</i>			
Underweight	-0.13 (-0.26; 0.003)	0.06	-0.16 (-0.29; -0.03)	0.014
Overweight	-0.17 (-0.27; -0.06)	0.003	0.01 (-0.11; 0.12)	0.92
Playing sports (h/w)				
<2	<i>Reference</i>			
2-4	0.17 (0.07; 0.26)	<0.001	0.10 (0.02; 0.19)	0.017
≥ 4	0.09 (-0.02; 0.20)	0.11	0.04 (-0.06; 0.15)	0.43
Screen time (h/d)				
<2	<i>Reference</i>			
≥ 2	-0.39 (-0.46; -0.32)	<0.001	-0.31 (-0.38; -0.24)	<0.001
Parental characteristics				
Age mother at 9-year visit (y)	0.01 (0.01; 0.02)	<0.001	0.001 (-0.01; 0.01)	0.71
BMI mother				
Normal weight	<i>Reference</i>			
Underweight	0.35 (-0.04; 0.75)	0.08	0.30 (-0.08; 0.68)	0.12
Overweight	-0.22 (-0.29; -0.14)	<0.001	-0.10 (-0.17; -0.02)	0.01
Marital status mother				
Married/ partner	<i>Reference</i>			
No partner	-0.23 (-0.34; -0.12)	<0.001	-0.03 (-0.15; 0.09)	0.61
Educational level mother				
No higher education	<i>Reference</i>			
Higher education	0.44 (0.37; 0.51)	<0.001	0.29 (0.21; 0.37)	<0.001
Smoking mother				
Never smoker	<i>Reference</i>			

Table 2.4.4. Associations between sociodemographic and lifestyle factors and the diet quality score (continued)

	Basic model ^a		Multivariable model ^b	
	β (95 % CI)	p-value	β (95 % CI)	p-value
Past smoker	0.05 (-0.03; 0.14)	0.23	0.05 (-0.03; 0.13)	0.26
Current smoker	-0.26 (-0.38; -0.14)	<0.001	-0.13 (-0.25; -0.02)	0.027
Household income per month				
<2800 €	Reference			
≥2800 €	0.32 (0.23; 0.40)	<0.001	0.15 (0.05; 0.25)	0.004

^aValues are regression coefficients with 95% confidence intervals from linear regression analyses adjusted for total energy intake.

^bValues are regression coefficients with 95% confidence intervals from multivariable linear regression analyses including all variables presented in the table.

IQR= Interquartile range; FFQ=Food-frequency questionnaire; y= year; BMI=Body mass index; kcal/ d= kilocalorie/ day; h/w= hours/ week; h/d= hours/ day.

Tracking of diet quality from early to mid-childhood

For children with dietary data at both the ages of 1 year and 8 years (n=2,608), we observed a Pearson's correlation of $r=0.19$ for the diet quality score between both ages ($p<0.01$). Significant correlations were also found between the seven individual diet quality score components that were examined at both time points ($r=0.11$ to 0.23 , all $p<0.01$) (**Supplemental table 2.4.3**). A linear weighted kappa showed slight agreement between the diet quality scores at both ages ($k=0.11$ (95%CI: 0.08; 0.14) for their rank in the lowest 25%, middle 50%, or highest 25% of the scores (Supplemental table 2.4.4).

DISCUSSION

We developed a food-based diet quality score based on Dutch dietary guidelines to estimate overall diet quality of children. Using this score, we evaluated diet quality of over 4,700 children at the age of 8 years in a population-based cohort in the Netherlands. We observed that diet quality in this population was suboptimal and none fully adhered to the guidelines. Factors that correlated with a higher diet quality in this group were, amongst others, a higher socioeconomic status and no maternal smoking. We observed only weak tracking of diet quality between the ages of 1 year and 8 years.

Interpretation of findings and comparison with previous research

Diet quality was suboptimal in our study population of 8-year-old children. This is consistent with studies in the US, Brazil, and the UK that showed less than optimal diet quality in similarly aged children⁹⁻¹¹. Compared to the American population aged 7-9 years⁹, level of adherence was similar for the fruit and vegetable components, however, for the dairy component adherence was lower in our study population. This might be

explained by the difference in scoring, as we only included recommended food items in the dairy component, whereas in the US-based study all dairy items were included⁹.

As expected from previous studies²⁷, socioeconomic status was positively associated with diet quality. A strong association was observed particularly for maternal educational level, independent of household income and other factors. Previous studies indicated that individuals with a higher educational level could have more nutritional knowledge^{4,27,28}; our study suggests that this also translates to the diet provided to their children. Furthermore, families with a higher income may be more able to buy healthy, more expensive, food products^{29,30}, explaining our association of household income with child diet quality, independent of educational level. A previous study among households in Canada showed that access to dairy, fruit and vegetables, which are food groups that positively contribute to our diet score, may be constrained by low income irrespective of educational level³¹. Unfortunately we did not assess food security in our study, which may partly explain the association between socioeconomic status and diet quality found in our study.

The negative association of maternal smoking with diet quality score is consistent with previous research among 515 children aged 2 to 17 years in the U.S., which showed that children from low-income families with parents who smoked, had a poorer diet quality than children from low-income families with nonsmoking parents³². We also observed a negative association between maternal overweight and child's diet quality score, which is in line with a previous study among 1,640 children aged 3 years in the UK³³. These findings for maternal smoking and overweight suggest that an unhealthy lifestyle of the mother negatively influences their child's diet quality, independent of socioeconomic status.

Independent of these maternal factors, we also found an association between children's lifestyle and their diet quality. Being more physically active and having less screen time were associated with a higher diet quality in children, although for physical activity this association was not observed for the group with the highest levels of physical activity. These findings are consistent with previous research, which showed that sedentary behavior is associated with a less healthy diet^{27,33}. Finally, in our fully adjusted models, we observed that children with a Moroccan ethnic background had a higher diet quality score than those with a Dutch ethnicity. Children with a Moroccan ethnicity had a higher intake of fish, legumes, and nuts and a lower high-fat and processed meat intake, suggesting a more Mediterranean-style diet³⁴.

In our study population, we found only weak tracking of the diet quality score and its individual components between the ages of 1 year and 8 years. Studies on tracking of diet from early life to later childhood are limited³⁵. One previous study found moderate to fair tracking of the intake of fruit, vegetables, and sugar-sweetened beverages from the age of 18 months to 7 years³⁵. A review by Nicklaus & Remy (2013), showed moder-

ate tracking of eating habits after the first year of life³⁶. Combined, these results suggest that tracking of diet may start after the age of 1 year.

Methodological considerations of the diet quality score

The diet quality score was positively associated with intake of micronutrients, indicating adequate construct validity, since dietary recommendations are, amongst others, developed to provide a sufficient supply of nutrients²⁴. We included both healthy and unhealthy components in the score, which may better capture overall diet quality than including healthy or unhealthy components only, as eating healthy foods is not necessarily inversely related to eating unhealthy foods³⁷. Further research is needed to examine whether this combined score is indeed associated with child health. Another strength of our diet score is the use of cut-off values based on current dietary recommendations instead of using a population-specific cut-off value such as a population-specific median intake, which may not be related to an actual healthy intake level²⁴. Finally, a strength of our diet quality scoring system is that we used a continuous scale, which provides more detail and is more accurate in ranking children with respect to diet quality than a dichotomous scoring system²⁴.

Constructing an overall diet quality index involves many choices²⁴. Although it may have been preferred to ascribe greater weights to components that have a greater effect on health, not enough information on the overall health effects of individual components was available, so we chose not to apply any weighting. In addition to the number and weights of components, another aspect to consider is the type of components included in the diet index. Most diet indices are based on intake of nutrients or food groups, or a combination of these, and some indices also include measures of dietary variety^{24,38}. We chose to construct our diet quality score on the basis of intake of food groups only, in line with the Dutch dietary guidelines, but we also observed positive associations of the diet score with intake of micronutrients, suggesting it represent an overall healthy diet. When diet quality score components are similar to each other or when they are strongly correlated, they contribute more heavily to the score²⁴. However, in our diet quality score, we observed low correlation between the diet score components (r -0.13 to 0.08). Finally, because our diet quality score is based on Dutch recommendations, important food groups may be absent for children with another ethnic background. However, the Dutch recommendations are comparable to recommendations in other countries³⁹. Furthermore, a systematic review conducted by Gilbert & Khokhar (2008) showed that after moving to a Western country, the majority of ethnic groups change their eating habits to a more Western diet⁴⁰. Also, we did not find major differences in diet score between the different ethnic groups in our population, suggesting that the Dutch recommendations and our diet score were also suitable for the study participants with another ethnicity.

Strengths and limitations

Major strengths of the Generation R study, in which we applied our diet quality score, are the population-based prospective cohort design and the large number of subjects. Also, we had information available on many parental and child sociodemographic and lifestyle factors for which we could examine their correlation with diet quality. However, there may be other correlates of diet quality that were not assessed in our study. Unfortunately, we had no detailed information on physical activity of the children. We used the amount of time participating in organized sports as a proxy for physical activity, which may not be an optimal measure, because it does not take into account other sources of physical activity. In addition, not all correlates were assessed at the same moment as dietary assessment, which may have influenced the associations. However, there was a high correlation of most variables throughout childhood, and we expect that any changes in these correlates are only limited. Therefore, we chose the time point that was the closest to our moment of dietary assessment. Furthermore, non-response analyses showed that non-responders to the FFQ more often had characteristics associated with a lower diet quality score, such as a lower educational level, suggesting that diet quality might be even lower in the children in Rotterdam than observed in the study population for which we had data available.

Another limitation of the study was the assessment of dietary intake with an FFQ. Limitations of FFQs in general are that they contain a limited amount of food items, and recollection of the consumed foods and portion sizes can be sources of error⁴¹. The FFQ used in our study was validated against the doubly labelled water method, regarded as the golden standard for the determination of total energy expenditure in free-living subjects, and this validation showed a reasonable capacity of the FFQ to rank subjects with respect to energy intake¹⁷. However, the FFQ was not validated for the intake of specific foods or food groups. Finally, for our analyses on tracking of diet quality, a limitation was that diet quality was not scored in exactly the same manner at the ages of 1 and 8 years and that no data on dietary intake were available for the period in between these two age categories.

Implications

The results of our study suggest that overall diet quality among 8-year-old children in our study population in an urban multi-ethnic setting in the Netherlands is suboptimal and that none of the children fully adhered to the dietary guidelines. This is undesirable as a healthy diet is important for healthy growth and development of the child¹. However, future research is needed to assess whether a higher overall adherence to the dietary guidelines is indeed associated with a better health and evaluate diet quality in other populations of children. Although the observed effect estimates for the correlates of diet quality were relatively small on an individual level, these may be relevant for

public health strategies. We observed for example that children from higher educated mothers had a 0.3 higher diet quality score (scale 0 to 10) than children whose mothers had not completed higher education. Most of the observed correlates of diet quality in our study are in line with previous research. Consistent with other studies, we found low socioeconomic status to be a strong predictor of a lower diet quality²⁷, emphasizing the need to target child dietary interventions especially to families with a lower socioeconomic status. Interventions should focus on promotion of healthy food products and increase the accessibility of these foods for these groups. Additionally, interventions should also discourage the consumption of unhealthy food products. Adherence to the recommendations was particularly low for the moderation components in our study population, underlining the importance of discouraging the intake of sugar-containing beverages and high-fat and processed meat. As previous evidence showed tracking of diet between mid-childhood and adulthood¹⁴, dietary interventions targeted at children are expected to not only improve diet quality during childhood, but also their diet quality into adulthood. However, we observed only weak tracking of diet quality from early to mid-childhood. Therefore, further research is needed to establish the optimal age and also the best target groups (e.g., children, parents, and/or schools) for dietary interventions in order to improve long-term diet quality.

Conclusion

To conclude, in this large population-based cohort in the Netherlands, we observed that diet quality of 8-year-old children was suboptimal, which indicates that they do not meet the current dietary guidelines. Particularly the intake of legumes, nuts, and oils or soft or liquid fats was too low, whereas the intake of sugar-containing beverages and high-fat and processed meat was too high. Main sociodemographic and lifestyle factors that correlated with a higher diet quality were a higher maternal education, a higher household income, no maternal smoking, and less time spent on watching television or using a computer of the child. Tracking of diet quality from the age of 1 year to 8 years was weak.

REFERENCES

1. Boumtje PI, Huang CL, Lee J-Y, Lin B-H. Dietary habits, demographics, and the development of overweight and obesity among children in the United States. *Food Policy*. 2005;30(2):115-128.
2. Demory-Luce D, Morales M, Nicklas T, Baranowski T, Zakeri I, Berenson G. Changes in food group consumption patterns from childhood to young adulthood: the Bogalusa Heart Study. *J Am Diet Assoc*. 2004;104(11):1684-1691.
3. Florence MD, Asbridge M, Veuglers PJ. Diet quality and academic performance. *J Sch Health*. 2008;78(4):209-215; quiz 239-241.
4. Mikkilä V, Räsänen L, Raitakari OT, Pietinen P, Viikari J. Consistent dietary patterns identified from childhood to adulthood: the cardiovascular risk in Young Finns Study. *Br J Nutr*. 2005;93(06):923-931.
5. Hu FB. Dietary pattern analysis: a new direction in nutritional epidemiology. *Curr Opin Lipidol*. 2002;13(1):3-9.
6. Nicklas T. Assessing diet quality in children and adolescents. *J Am Diet Assoc*. 2004;104(9):1383-1384.
7. Voortman T, Leermakers ET, Franco OH, et al. A priori and a posteriori dietary patterns at the age of 1 year and body composition at the age of 6 years: the Generation R Study. *Eur J Epidemiol*. 2016;31(8):775-783.
8. Marshall S, Burrows T, Collins CE. Systematic review of diet quality indices and their associations with health-related outcomes in children and adolescents. *Journal of Human Nutrition and Dietetics*. 2014;27(6):577-598.
9. Carlson A, Lino M, Gerrior S, Basiotis PP. Report card on the diet quality of children ages 2 to 9. *Family Economics and Nutrition Review*. 2003;15(2):52.
10. de Farialli CP, Cadel NV, Zandonadev E. Socioeconomic predictors of child diet quality. *Rev Saúde Pública*. 2010;44(5):785-732.
11. Jennings A, Welch A, van Sluijs EM, Griffin SJ, Cassidy A. Diet quality is independently associated with weight status in children aged 9–10 years. *J Nutr*. 2011;141(3):453-459.
12. van Lee L, Geelen A, van Huysduynen EJ, de Vries JH, van't Veer P, Feskens EJ. The Dutch Healthy Diet index (DHD-index): an instrument to measure adherence to the Dutch Guidelines for a Healthy Diet. *Nutr J*. 2012;11(1):49.
13. Voortman T, Kiefte-de Jong JC, Geelen A, et al. The development of a diet quality score for preschool children and its validation and determinants in the Generation R Study. *J Nutr*. 2015;145(2):306-314.
14. Craigie AM, Lake AA, Kelly SA, Adamson AJ, Mathers JC. Tracking of obesity-related behaviours from childhood to adulthood: A systematic review. *Maturitas*. 2011;70(3):266-284.
15. Netherlands Nutrition Center. *Richtlijnen Schijf van Vijf (Wheel of Five guidelines)*. The Hague: Netherlands Nutrition Center (Voedingscentrum);2016.
16. Kooijman MN, Kruihof CJ, van Duijn CM, et al. The Generation R Study: design and cohort update 2017. *Eur J Epidemiol*. 2016;31(12):1243-1264.
17. Dutman AE, Stafleu A, Kruizinga A, et al. Validation of an FFQ and options for data processing using the doubly labelled water method in children. *Public Health Nutrition*. 2011;14(03):410-417.
18. Netherlands Nutrition Center. *Zo eet Nederland: resultaten van de Voedselconsumptie Survey 1997-1998 (Results of the Dutch Food Consumption Survey 1997-1998)*. The Hague: Netherlands Nutrition Center (Voedingscentrum);1998.

19. Health Council of the Netherlands. *Richtlijnen Goede Voeding 2015 (Guidelines for a healthy diet 2015)*. The Hague: The Health Council of the Netherlands (Gezondheidsraad);2015.
20. Voortman T, van den Hooven EH, Heijboer AC, Hofman A, Jaddoe VW, Franco OH. Vitamin D deficiency in school-age children is associated with sociodemographic and lifestyle factors. *J Nutr*. 2015;145(4):791-798.
21. Cole TJ, Bellizzi MC, Flegal KM, Dietz WH. Establishing a standard definition for child overweight and obesity worldwide: international survey. *Bmj*. 2000;320(7244):1240.
22. Strasburger VC, Hogan MJ, Mulligan DA, et al. Children, adolescents, and the media. *Pediatrics*. 2013;132(5):958-961.
23. Netherlands Bureau for Economic Policy Analysis. *Onzekere wereld; Nederlandse economie stabiel (Uncertain world; Dutch economy stable)* The Hague: Netherlands Bureau for Economic Policy Analysis (Centraal Planbureau);2016.
24. Waijers PM, Feskens EJ, Ocke MC. A critical review of predefined diet quality scores. *Br J Nutr*. 2007;97(2):219-231.
25. Lowry R. Kappa as a Measure of Concordance in Categorical Sorting. 2016; <http://vassarstats.net/kappa.html>. Accessed 01-06, 2016.
26. Landis JR, Koch GG. The measurement of observer agreement for categorical data. *Biometrics*. 1977;33(1):159-174.
27. Darmon N, Drewnowski A. Does social class predict diet quality? *Am J Clin Nutr*. 2008;87(5):1107-1117.
28. Parmenter K, Waller J, Wardle J. Demographic variation in nutrition knowledge in England. *Health Educ Res*. 2000;15(2):163-174.
29. Eikenberry N, Smith C. Healthful eating: perceptions, motivations, barriers, and promoters in low-income Minnesota communities. *J Am Diet Assoc*. 2004;104(7):1158-1161.
30. Story M, Kaphingst KM, Robinson-O'Brien R, Glanz K. Creating healthy food and eating environments: policy and environmental approaches. *Annu Rev Public Health*. 2008;29:253-272.
31. Kirkpatrick S, Tarasuk V. The relationship between low income and household food expenditure patterns in Canada. *Public Health Nutrition*. 2003;6(06):589-597.
32. Johnson RK, Wang M, Smith MJ, Connolly G. The association between parental smoking and the diet quality of low-income children. *Pediatrics*. 1996;97(3):312-317.
33. Fisk CM, Crozier SR, Inskip HM, et al. Influences on the quality of young children's diets: the importance of maternal food choices. *Br J Nutr*. 2011;105(2):287-296.
34. El Rhazi K, Nejari C, Romaguera D, et al. Adherence to a Mediterranean diet in Morocco and its correlates: cross-sectional analysis of a sample of the adult Moroccan population. *BMC Public Health*. 2012;12(1):1.
35. Bjelland M, Brantsaeter AL, Haugen M, Meltzer HM, Nystad W, Andersen LF. Changes and tracking of fruit, vegetables and sugar-sweetened beverages intake from 18 months to 7 years in the Norwegian Mother and Child Cohort Study. *BMC Public Health*. 2013;13(1):793.
36. Nicklaus S, Remy E. Early origins of overeating: tracking between early food habits and later eating patterns. *Current Obesity Reports*. 2013;2(2):179-184.
37. Anderson SE, Ramsden M, Kaye G. Diet qualities: healthy and unhealthy aspects of diet quality in preschool children. *Am J Clin Nutr*. 2016;103(6):1507-1513.
38. Huybrechts I, Vereecken C, De Bacquer D, et al. Reproducibility and validity of a diet quality index for children assessed using a FFQ. *Br J Nutr*. 2010;104(1):135-144.

39. Food and Agriculture Organization of the United Nations. Food-based dietary guidelines. 2016; <http://www.fao.org/nutrition/education/food-dietary-guidelines/home/en/>. Accessed 28-12-2016.
40. Gilbert PA, Khokhar S. Changing dietary habits of ethnic groups in Europe and implications for health. *Nutr Rev.* 2008;66(4):203-215.
41. Newby PK, Hu FB, Rimm EB, et al. Reproducibility and validity of the Diet Quality Index Revised as assessed by use of a food-frequency questionnaire. *Am J Clin Nutr.* 2003;78(5):941-949.

Supplemental table 2.4.1. Median intake per diet quality score component and median score per diet quality score component (maximum score is 1) for boys (n=2364) and girls (n=2369) separately

Component	Unit	Cut-off values	Intake		Score	
			Boys	Girls	Boys	Girls
Fruit	g/d	≥ 150	111 (77-165)	111 (77-167)	0.74 (0.51-1.00)	0.74 (0.51-1.00)
Vegetables	g/d	≥ 150	77 (47-122)	80 (51-123)	0.51 (0.31-0.82)	0.53 (0.34-0.82)
Whole-grains	g/d	≥ 90	101 (65-135)	98 (65-130)	1.00 (0.72-1.00)	1.00 (0.72-1.00)
Fish	g/w	≥ 60	38 (0-95)	38 (0-79)	0.63 (0.00-1.00)	0.63 (0.00-1.00)
Legumes	g/w	≥ 84	35 (0-70)	18 (0-70)	0.42 (0.00-0.83)	0.21 (0.00-0.83)
Nuts	g/d	≥ 15	4 (0-11)	2 (0-8)	0.29 (0.00-0.70)	0.14 (0.00-0.53)
Dairy	g/d	≥ 300	182 (56-319)	152 (54-284)	0.61 (0.19-1.00)	0.51 (0.18-0.95)
Oils and soft or liquid fats	g/d	≥ 30	11 (2-22)	11 (2-17)	0.37 (0.07-0.74)	0.37 (0.07-0.56)
Sugar-containing beverages	g/d	≤ 150	335 (193-557)	311 (169-506)	0.00 (0.00-0.00)	0.00 (0.00-0.00)
High-fat and processed meat	g/w	≤ 250	340 (232-482)	306 (203-430)	0.00 (0.00-0.07)	0.00 (0.00-0.19)

Values are median (IQR).

Energy intake (kcal/d) was 1531 (IQR 1299-1763) among boys and 1391 (IQR 1184-1606) among girls.

g/d= gram/ day; g/w= gram/ week; IQR= Interquartile range; kcal/d= kilocalorie/ day.

Supplemental table 2.4.2. Associations between the diet quality score and intake of nutrients

	Pearson correlation (r) Unadjusted	Pearson correlation (r) Adjusted for energy intake
Macronutrients:		
Fat (g)	0.26	-0.02
Saturated fat (g)	0.20	-0.11
N-3 fatty acids (mg)	0.26	0.24
Protein (g)	0.40	0.29
Animal protein (g)	0.24	0.05
Plant protein (g)	0.49	0.41
Carbohydrates (g)	0.24	-0.11
Monosaccharides and disaccharides (g)	0.15	-0.11
Dietary fiber (g)	0.62	0.58
Micronutrients:		
Vitamin B 1 (mg)	0.46	0.36
Vitamin B 2 (mg)	0.38	0.25
Vitamin B 3 (niacin) (mg)	0.42	0.31
Vitamin B 6 (mg)	0.43	0.32
Vitamin B 12 (µg)	0.30	0.18
Vitamin C (mg)	0.29	0.16
Vitamin D (µg)	0.40	0.30
Calcium (mg)	0.39	0.27
Copper (mg)	0.54	0.52
Iron (mg)	0.45	0.36
Magnesium (mg)	0.55	0.55
Phosphorus (mg)	0.48	0.44
Selenium (µg)	0.37	0.24
Zinc (mg)	0.41	0.31

All *r*-values were statistically significant at the 0.05 level, except for total fat (adjusted for energy intake): $p=0.16$.
g= gram; *mg*= milligram; μg = microgram.

Supplemental table 2.4.3. Associations between the diet quality score and its components at the age of 1 y and at the age of 8 y (n=2,608)

Component	Pearson correlation (r)
Fruit	0.20
Vegetables	0.11
Whole-grains	0.19
Fish	0.23
Dairy	0.14
Oils and unsaturated fats	0.19
Sugar-containing beverages	0.12
Total	0.19

All significant at 0.01 level

The components 'legumes' and 'nuts' were not incorporated in the diet quality score at age 1 year, and the 'meat' component was an adequacy component (low-fat unprocessed meat) in the preschool score and a moderation component (high-fat and processed meat) in the childhood score, therefore no tracking was assessed for these components

Supplemental table 2.4.4. Tracking matrix constructed for the calculation of weighted Kappa statistic for diet quality score at age 1 y and age 8 y among children with dietary data at both time points (n=2,607)

	Lowest 25% at 8 y	Middle 50% at 8 y	Highest 25% at 8 y
Lowest 25% at 1 y	201	339	112
Middle 50% at 1 y	302	653	348
Highest 25% at 1 y	104	317	231

The entry in a specific cell indicates the number of subjects belonging to the corresponding classes at age 1 y and age 8 y

Additional supplemental material for this chapter can be found online:

https://static-content.springer.com/esm/art%3A10.1007%2Fs00394-018-1651-z/MediaObjects/394_2018_1651_MOESM1_ESM.docx