



ORIGINAL ARTICLE

Basic and Translational Allergy Immunology



WILEY

Induction of IgG₂ and IgG₄ B-cell memory following sublingual immunotherapy for ryegrass pollen allergy

Jorn J. Heeringa^{1,2,3} | Craig I. McKenzie¹ | Nirupama Varese^{1,4} |
Mark Hew^{4,5} | Amy T. C. M. Bakx¹ | Pei M. Aui¹ | Jennifer M. Rolland^{1,4} |
Robyn E. O'Hehir^{1,4} | Menno C. van Zelm^{1,4}

¹Department of Immunology and Pathology, Central Clinical School, Monash University, Melbourne, Vic., Australia

²Department of Immunology, Erasmus MC, University Medical Center, Rotterdam, the Netherlands

³Department of Pediatrics, Erasmus MC, University Medical Center, Rotterdam, the Netherlands

⁴Department of Respiratory Medicine, Allergy and Clinical Immunology (Research), Central Clinical School, Monash University, and Alfred Hospital, Melbourne, Vic., Australia

⁵School of Public Health and Preventive Medicine, Monash University, Melbourne, Vic., Australia

Correspondence

Menno C. van Zelm, Department of Immunology and Pathology, Central Clinical School, Monash University, 89 Commercial Road, Melbourne, Vic. 3004, Australia.
Email: menno.vanzelm@monash.edu

Funding information

Sophia Children's Hospital Fund, Grant/Award Number: S698; Num Pon Soon Charitable Trust; National Health and Medical Research Council, Grant/Award Number: GNT1117687

Abstract

Background: While treatment for atopic rhinitis is aimed mostly to relieve symptoms, only allergen-specific immunotherapy (AIT) is targeted to modify the natural history of allergic diseases. This results in sustained clinical tolerance, even when treatment has stopped. The immunomodulatory effects of AIT are attributed mainly to increased regulatory T-cell function and increased allergen-specific IgG₄, yet little is known about the effect on the memory B-cell compartment.

Objective: We aimed to examine the effects of AIT on the IgE- and IgG subclass-expressing memory B cells.

Methods: We recruited 29 patients with atopic seasonal rhinoconjunctivitis and performed a longitudinal analysis of the peripheral immune compartment before, during, and after sublingual immunotherapy (SLIT) for allergy to temperate grass pollen, predominantly to ryegrass pollen (RGP; *Lolium perenne*). Using flow cytometry on peripheral blood mononuclear cells and serum immunoassays, we analyzed the effects of a 4 months preseasonal treatment regimen comprising two or three courses in consecutive years on circulating IgE⁺ and IgG⁺ memory B cells and allergen-specific Ig levels.

Results: SLIT increased RGP-specific serum IgG₂ and IgG₄, as well as the frequencies of IgG₂⁺ and IgG₄⁺ memory B cells, whereas no effect was observed on the IgE⁺ memory B-cell compartment. Furthermore, SLIT enhanced proportions of regulatory T cells specific to RGP. These changes were associated with clinical improvement.

Conclusion: Our data provide evidence for immunological effects of SLIT on B-cell memory. Skewing responses toward IgG₂ and IgG₄ subclasses might be a mechanism to suppress IgE-mediated allergic responses.

Abbreviations: AIT, allergen-specific immunotherapy; FeNO, fractional exhaled nitric oxide; IL, interleukin; RGP, ryegrass pollen; SCIT, subcutaneous immunotherapy; SHM, somatic hypermutation; SLIT, sublingual immunotherapy; SPT, skin prick test; Th1/2, T helper 1/2; Treg, regulatory T cell; VAS, visual analog score.


Jorn J. Heeringa and Craig I. McKenzie equal contribution.

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

© 2019 The Authors. *Allergy* published by John Wiley & Sons Ltd.

KEYWORDS

B cells, flow cytometry, IgE, immunotherapy and tolerance induction, Rhinitis



GRAPHICAL ABSTRACT

This study examines the effect of ryegrass pollen AIT on B-cell responses in a population of 29 patients with allergic rhinitis. Successful immunotherapy for ryegrass pollen allergy increases allergen-specific IgG₂ and IgG₄ serum levels, and proportions of IgG₂⁺ and IgG₄⁺ expressing memory B cells. Skewing toward the anti-inflammatory IgG₂ and IgG₄ subclasses might be a mechanism to suppress IgE-mediated allergic responses.

1 | INTRODUCTION

Rhinoconjunctivitis and other IgE-mediated allergies are an increasing disease burden globally.¹ Most therapies for allergies are directed at relieving symptoms, but allergen-specific immunotherapy (AIT) is the only current therapy that modifies the natural course of allergic diseases. Subcutaneous immunotherapy (SCIT) and sublingual immunotherapy (SLIT) are both proven effective treatments for grass pollen-induced rhinoconjunctivitis.²⁻⁴ The therapeutic effect is maintained beyond the conclusion of treatment.⁵⁻⁷ In patients with allergic rhinitis, AIT can prevent the onset of new sensitizations⁸ and decrease the likelihood of developing asthma.⁹ The immunomodulatory properties of AIT affect local and systemic immune responses, with an impact on the number and function of mast cells, basophils, antigen-presenting cells, T cells, and B cells.^{10,11}

Allergic patients manifest sensitization by means of allergen-specific IgE bound to effector cells, particularly mast cells and basophils.¹² The underlying mechanism is thought to be a shifted T-cell balance toward a T helper 2 (Th2) phenotype, and these cells produce interleukin (IL)-4 and IL-13 that direct allergen-specific B cells to produce IgE.¹³ Furthermore, Th2 cells produce IL-5 which promotes the involvement of eosinophils in the pathogenesis of allergic diseases.¹⁴ In contrast, Th1 responses are promoted by IFN- γ and skew away from a Th2 phenotype.¹⁵ Effective immunotherapy

has been shown to reverse the Th2 dominance and to result in anergy of allergen-specific T cells,^{16,17} induction of regulatory T cells (Treg),¹⁸⁻²¹ and production of blocking antibodies of the IgG and IgA isotypes.^{22,23} Specifically, TGF- β and IL-10 produced by Treg are pivotal for the successful immune deviation in AIT.^{24,25}

The tolerogenic functions of IL-10 are extensive, but mainly encompass the inhibition of mast cell activity,²⁶ suppression of IL-5 production by Th2 cells,²⁷ and cell death induction in eosinophils.²⁸ Furthermore, IL-10 in combination with IL-4 and IL-13 directs B-cell immunoglobulin class switching to IgG₄ instead of IgE.²⁹ Indeed, one of the known effects of AIT is an increase in allergen-specific serum IgG₄ and an increased serum IgG₄/IgE antibody ratio that is associated with clinical efficacy.³⁰

SCIT and SLIT have distinct immunomodulatory capabilities that appear related to the different routes of administration. Sublingual administration results in fewer systemic adverse effects, but some studies indicate diminished clinical and immunological efficacy compared with subcutaneous administration.^{2,31} SLIT results in increased numbers of FoxP3⁺ Treg both in the oral epithelium and in the peripheral blood.^{23,32} Further systemic alterations are more diverse. Some studies report an initial increase in allergen-specific IgE serum levels, followed by a decrease after 1 month.³³ Furthermore, allergen-specific IgG₂, IgG₄, and IgA serum levels are reported to increase in as little as 1 day after the start of therapy.³³⁻³⁵ However, other studies detected no systemic alterations with regard to

allergen-specific lymphoproliferation, cytokine secretion, or Ig serum levels.^{36,37}

IgG₂ and IgG₄ heavy chain constant regions are encoded by genes in the *IGH* locus. Ig class switching to IgG₂ and IgG₄ frequently occurs indirectly following a switch from IgM to the more proximal IgG₃ and IgG₁ genes rather than directly from IgM to IgG₂ or IgG₄.³⁸ Given the higher loads of somatic hypermutation (SHM) in variable regions of IgG₂ and IgG₄ transcripts, it has been suggested that B cells expressing these transcripts have spent more time in the germinal center response.³⁹ In addition, the majority of IgG₂- and IgG₄-expressing B cells co-express CD27, and their frequencies increase with age.^{40,41} Hence, it appears that these Ig class switches occur following repeated exposure to the same antigen.

Since AIT has been shown to have long-lasting beneficial effects, it is important to determine whether this is the result of changes in immunological memory. We here address this question in our cohort of patients with moderate-to-severe seasonal allergic rhinitis, studied longitudinally before, during, and after SLIT for grass pollen allergy.⁴² As published previously,^{42,43} SLIT in our cohort resulted in allergic rhinitis symptom relief and conferred significant protection from epidemic thunderstorm asthma, making this an ideal cohort to examine the effects of a 4-month treatment regimen and the subsequent effects of two further courses of treatment over 3 years on circulating IgE⁺ and IgG subclass-expressing memory B cells and allergen-specific Ig levels.

2 | METHODS

2.1 | Study design

Using an open-label longitudinal design (ClinicalTrials.gov identifier: NCT02014623), 29 participants were recruited for treatment with a commercial 5-grass pollen SLIT tablet (Oralair®; Stallergenes) using a 4-month (May–September) regimen completed prior to the Australian pollen season, for 3 consecutive years (2014–2016; subject numbers at each time point shown in Figure 1A). Treatment with Oralair® involved dissolution under the tongue (at least 2 minutes) followed by swallowing the residue. The treatment regimen comprised the following: day 1–1 tablet 100 IR (index of reactivity); day 2–2 tablets 100 IR; and day 3 to day 120–1 daily tablet 300 IR. Blood samples were collected immediately before initial treatment (May 2014) and after the first 4 months of treatment (September 2014), followed by annual collections in May 2015 and May 2016 (prior to commencement of 2nd and 3rd courses of SLIT), and May 2017 (Figure 1A).

2.2 | Participant characteristics

Participants were recruited from The Alfred Hospital Allergy Clinic, Melbourne, Victoria, Australia. All had well-characterized moderate-to-severe seasonal allergic rhinitis (plus or minus asthma) due to RGP allergy with positive serum RGP-specific IgE (≥ 0.35 kU/L; ImmunoCAP, Phadia). Exclusion criteria were a




FIGURE 1 Study design and clinical parameters of allergic rhinitis decreased after SLIT. A, Timeline of SLIT for grass pollen allergy between May 2014 and 2017. Time points indicate blood sampling. B, Allergic rhinitis symptoms by visual analog scale measured during peak pollen season. C, Wheal diameter (in mm) from skin prick test (SPT) with RGP. D, Fractional exhaled nitric oxide (FeNO) measured immediately prior to starting SLIT. E, Total IgE in serum. Each dot represents one individual; red lines indicate median values. Statistical analysis was performed between baseline and each follow-up time point to assess changes induced by SLIT using the Wilcoxon signed-rank test; * $P < .05$, ** $P < .01$, and *** $P < .001$

co-existing immunodeficiency, previous immunotherapy within the last 5 years, ongoing immunotherapy with other allergens, and treatment with continuous oral corticosteroids and/or β -blockers. The use of usual medications for allergic rhinitis was permitted, including antihistamines and topical corticosteroids. Alfred Hospital Research and Ethics Committee approval and written informed consent from each participant were obtained prior to inclusion (project number 514/13). Twenty-nine participants (12 males) were recruited for treatment, with a mean age of 35 years (range 18–59 year) and mean serum RGP-specific IgE of 52 kU/L. Five withdrew after the first (baseline) time point ($n = 4$ tongue swelling, upset stomach; $n = 1$ failed to attend) leaving 24 participants who commenced SLIT (Table S1). At subsequent time points, three participants failed to attend after 4 months of treatment and a further seven participants were excluded after 1 year ($n = 2$ opted to receive SCIT, $n = 2$ opted to receive sublingual drops, and $n = 3$ withdrew). Blood samples for serum and flow cytometric analysis were obtained from $n = 24$ patients in May 2014 prior to starting SLIT, $n = 24$ in September 2014, $n = 21$ in May 2015, $n = 14$ in May 2016, and $n = 14$ in May 2017. Details on sample numbers for each analysis are included in Table S2. A further 5 RGP-allergic subjects who did not receive SLIT (ie, received usual medication alone) were included as untreated patients at baseline and 4 months.

2.3 | Clinical parameters of allergic rhinitis

Allergic rhinitis symptoms during the peak RGP season were recorded by the participants using a visual analog score (VAS; scale, 0–100). Fractional exhaled nitric oxide (FeNO, in parts per billion [ppb]; HypoAirFeNO) was measured according to the manufacturer's instructions (NIOX, Uppsala, Sweden). FeNO was measured immediately before the start of SLIT therapy outside of the grass pollen season to minimize effects of daily fluctuations in pollen levels.

2.4 | Quantification of serum total IgE and allergen-specific IgE, IgG₂, and IgG₄

Serum total IgE, RGP-specific-IgE, and -IgG₄ levels were measured by ImmunoCAP. Serum RGP-specific IgG₂ antibodies were measured by in-house ELISA, as described previously.⁴⁴ Briefly, ELISA plate wells were coated with an aqueous RGP extract (Stallergenes Greer), blocked with 2% bovine serum albumin in PBS (Sigma-Aldrich), and incubated with serial dilutions of serum samples. Separate wells were coated with serial dilutions of purified human IgG₂ (Sigma-Aldrich, #I5404) to generate a standard curve for quantification of IgG₂ in serum samples. Bound IgG₂ was detected using biotinylated anti-hIgG₂ (clone HP6002; Thermo Scientific) followed by Pierce High Sensitivity Streptavidin-HRP (Thermo Scientific). ELISA was developed using TMB (Thermo Scientific), and the reaction stopped with 1 mol/L HCl. Absorbance (OD 450 nm) was measured using a FLUOstar Optima plate reader (BMG Labtech).

2.5 | In vitro RGP stimulation of PBMC, Treg staining, and measurement of cytokines

PBMC were isolated by Ficoll-paque density centrifugation. Fresh PBMC were used for in vitro culture, and the remaining cells stored in liquid nitrogen. PBMC were labeled with CFSE (0.5 μ mol/L CFSE/ 10^7 PBMC; Molecular Probes) and cultured with an aqueous RGP extract (50 μ g/mL; Stallergenes Greer) or tetanus toxoid (20 Lfu/mL; Statens Serum Institut, Copenhagen, Denmark). On day 7, cells were stained with CD4-PE Cy7, CD25-PE (both from BD Biosciences), FoxP3-APC (eBioscience), and aqua live/dead dye (Life Technologies). The Treg gating strategy is shown in Figure S1. Data were acquired using an LSR-II flow cytometer (BD Biosciences).

The levels of IFN- γ , IL-5, IL-10, and IL-13 in 7-day culture supernatants were determined using a Luminex human premixed multi-analyte kit (R&D Systems Inc) according to the manufacturer's instructions. Due to changes in IL-5 production observed after 3 years of SLIT, IL-13 was also assessed at the same time point to further investigate Th2 cytokine production. Tetanus toxoid was included as a control antigen to determine RGP specificity. "No antigen" values were subtracted from test values.

2.6 | B-cell subset analysis by flow cytometry

One million thawed PBMC were incubated with 11-color antibody cocktails against B-cell markers for 15 minutes at room temperature in 100 μ L total volume (Table S3). Flow cytometric analyses were performed on a 4-laser LSRFortessa (BD Biosciences), and data were analyzed using FACSDiva V8.0 (BD Biosciences). B-cell subsets were defined as described previously.^{41,45,46} Briefly, within the CD19⁺ B-cell population, the proportions were determined of plasmablasts (CD27⁺CD38^{high}), transitional (CD27⁺CD38^{high}), naive mature (CD27⁺IgM⁺IgD⁺), natural effector memory B cells (CD27⁺IgM⁺IgD⁺), and IgM-only memory B cells (CD27⁺IgM⁺IgD⁻). Furthermore, we analyzed Ig switched CD27⁺CD38^{dim} and CD27⁺CD38^{dim} memory B cells expressing IgA, IgE, IgG, or each of the 4 IgG subclasses.

2.7 | Molecular analysis of Ig gene rearrangements

RNA was isolated from PBMC from a limited cohort of 5 subjects treated with SLIT (Table E1; patient no. 1, 11, 13, 15, and 19) with a GenElute mammalian RNA kit (Sigma-Aldrich) and reverse transcribed to cDNA with random primers (Invitrogen Life Technologies, Waltham, MA). Rearranged IgG transcripts were amplified in a multiplex PCR approach using 4 different IGHV family leader forward primers in combination with an IGHG-consensus reverse primer.⁴⁷ PCR products were cloned into a pGEMT easy vector (Promega), amplified by colony PCR, and sequenced by the Micromon facility of Monash University on an Applied Biosystems 3730s DNA Analyzer (Thermo Scientific). Obtained sequences were analyzed using the IMGT database (<http://www.imgt.org>) to assign the IGHV, IGHJ, and IGHJ gene alleles and to identify SHM. For each unique clone, the position and frequency of

mutations were determined within the entire *IGHV* gene (FR1-CDR1-FR2-CDR2-FR3). SHM was determined as variations on the best-matched V-gene and represented as the percentage of mutations of the total sequenced V-gene nucleotides. The IgG subclasses were determined using the IGH reference sequence (NG_001019).

2.8 | Statistical analysis

Differences in symptom scores, serum Ig values, cytokines, and B- and T-cell subsets before, during, and after treatment were analyzed with the Wilcoxon signed-rank test. All analyses were two-tailed, and differences were considered statistically significant if P -values were $<.05$. Due to missing values at year 1, 2, and 3 measures, it was not possible to use repeated measures ANOVA. Therefore, we performed pairwise analysis between time point 0 and each follow-up sample. Differences in IgG subclass usage of unique *IGH* transcripts were statistically analyzed with the chi-squared test. Statistical analysis was performed using GraphPad Prism software, version 7.01 (GraphPad Software).

3 | RESULTS

3.1 | SLIT reduces symptoms of allergic rhinitis

To study the clinical effects of SLIT, we assessed the severity of symptoms for allergic rhinitis using a VAS. Before the start of treatment, participants reported a median VAS of 80 mm for the 2013 pollen season (Figure 1B). In the first pollen season after commencing SLIT, participants experienced fewer symptoms (median VAS 40 mm, $P < .001$), and these remained low for the second and third seasons following repeat SLIT courses (median VAS 20 mm at 2 years, $P < .001$; median VAS 35 mm at 3 years, $P < .01$), confirming sustained clinical efficacy.

The level of RGP sensitization was monitored by skin prick tests (SPT) with wheal diameters (mm) positively correlating with symptoms of allergic disease.⁴⁸ SLIT significantly decreased SPT wheal diameter in response to SPT with RGP extract within 1 year of commencing therapy (Figure 1C). Wheal size remained low on retesting after the third year of SLIT. In addition, the severity of airway inflammation and bronchial hyperreactivity was assessed by measurement of FeNO.^{49,50} SLIT significantly decreased FeNO from baseline at 1, 2, and 3 years after commencing SLIT (Figure 1D). The significant decreases in SPT wheal diameter and




FIGURE 2 Sublingual immunotherapy (SLIT) alters in vitro Treg proliferation and cytokine production in response to RGP. PBMC stimulated with RGP were assessed for A, Treg proliferation and production of B, IL-10, C, IFN- γ , and D, IL-5 were determined for all patients included at $t = 0, 1, 2, 3$ y. E. Paired analysis of IL-5 and F. IL-13 at $t = 0$ and $t = 3$ y. The lower limit of detection of IL-10 levels (0.5 pg/mL) in panel B is depicted by a dashed line and datapoints representing undetectable levels are placed below it. Statistical analysis was performed between baseline and each follow-up time point to assess changes induced by SLIT using the Wilcoxon signed-rank test; $*P < .05$, $**P < .01$, $***P < .001$ and $****P < .0001$




FIGURE 3 RGP-specific IgG₄ and IgG₄⁺:IgE⁺ memory B-cell ratio increased after a 4-mo course of SLIT. A, Gating strategy for Ig isotype and IgG subclass-expressing memory B cells by flow cytometry. B, RGP-specific IgG₂ and IgG₄ in sera and C, proportion of IgG₂⁺ and IgG₄⁺ memory B cells as a percentage of IgG⁺ population after 4 mo of SLIT. D, Proportion of IgE⁺ memory B cells as a percentage of total CD19⁺ B cells after 4 mo of SLIT. E, Ratio of IgG₄⁺ to IgE⁺ B-cell percentages after 4 mo of SLIT (from C and D). Each dot represents one individual; red lines indicate median values. Statistical analysis was performed between baseline and each follow-up time point to assess changes induced by SLIT using the Wilcoxon signed-rank test. **P* < .05 and *****P* < .0001

FeNO were consistent with decreased symptoms of allergic rhinitis 1 year after commencing treatment as well as after the second and third successive years of SLIT. The reduction in VAS and FeNO after SLIT did not correspond with any changes in total serum IgE levels (Figure 1E).

3.2 | Induction of RGP-specific Treg and IL-10 production following SLIT

Successful AIT has been associated with proliferation of allergen-specific Treg.⁵⁰ To assess Treg proliferation in response to allergen, we stimulated PBMC with RGP and measured the proliferation of activated Treg (CD4⁺CD25⁺FoxP3⁺) using CFSE. RGP-induced proliferation of Treg was enhanced by SLIT after 4 months and remained raised throughout subsequent years (Figure 2A).

Given the role of cytokines in skewing T-cell responses, we quantified IL-5, IL-10, IL-13, and IFN- γ production from RGP-stimulated PBMC. After 3 years of SLIT, T cells produced significantly more IL-10 and IFN- γ and significantly less IL-5 and IL-13 (Figure 2B-E). SLIT did not alter the PBMC cytokine response to tetanus toxoid (Figure S2). Taken together, the data from our study suggest SLIT enhances Treg response to allergen within 4 months without impacting the Treg response to other antigens. Furthermore, SLIT may alter the Th1/Th2 cytokine profile away from pro-allergic Th2 cytokines toward a regulatory and Th1-biased response.

3.3 | Increased IgG₄ serum levels and IgG₄⁺ memory B-cell frequencies after 4 months SLIT

To study the short-term effects of SLIT on the immune system, we analyzed serum Ig levels and B-cell subsets before and directly




FIGURE 4 Persistent increase in IgG₄ and late rise in IgG₂ after three 4-mo SLIT courses. A, RGP-specific IgE, IgG₂ and IgG₄ in serum. B, Proportions of IgE⁺, IgG₂⁺, and IgG₄⁺ memory B cells in peripheral blood. IgE⁺ memory B cells presented as a percentage of CD19⁺ B cells. IgG₂⁺ and IgG₄⁺ memory B cells presented as a percentage of IgG⁺ memory B cells. Baseline data for IgE⁺ and IgG₄⁺ B cells are the same as those in Figure 3C and D. Statistical analysis was performed between baseline and each follow-up time point to assess changes induced by SLIT using the Wilcoxon signed-rank test; **P* < .05 and ***P* < .01




FIGURE 5 Sublingual immunotherapy (SLIT) increases frequency of unique IgG2 transcripts. A, Schematics of the human IGH locus depicting the positioning of the constant gene regions relative to the rearranged VDJ exon. B, Somatic hypermutation frequencies of unique IgG transcripts obtained from 5 patients before (2014) and after SLIT (2017) and grouped per IgG subclass. C, Relative isotype distribution of unique IgG transcripts. Central number indicates total unique IgG sequences identified. Significance was determined by chi-squared test; **P = .0011

after the first 4 months of therapy. The gating strategies for flow cytometric detection of memory B cells expressing the four IgG subclasses are shown in Figure 3A. We observed that after 4 months of immunotherapy, RGP-specific serum IgG₂ increased from a median of 2.46 to 5.08 μg/mL (Figure 3B). Furthermore, all participants showed an increase in RGP-specific serum IgG₄ from a median of 0.37 μg/mL pretreatment to 1.16 μg/mL post-treatment after 4 months of SLIT. This was accompanied by a significant increase in the frequency of IgG⁺ memory B cells (CD19⁺CD38^{dim}) expressing IgG₄ (Figure 3C). The increase in the IgG₄⁺ memory B-cell frequencies was not directly correlated with the increase in RGP-specific serum IgG₄ ($p > .05$). SLIT did not change IgE⁺ memory B-cell frequencies (Figure 3D). However, the increase in IgG₄⁺ memory B cells resulted in a significantly higher IgG₄⁺/IgE⁺ memory B-cell ratio following 4 months of treatment (Figure 3E). The frequencies of all other B-cell subsets, including transitional, naive mature, memory, and plasmablasts, remained unchanged after 4 months of SLIT (Figure S3). Thus, 4 months SLIT quite specifically affected allergen-specific IgG₄ serum levels and the frequencies of IgG₄-expressing memory B cells.

3.4 | SLIT has persistent long-term effects on IgG₂ and IgG₄ memory B cells

In addition to short-term effects of SLIT, we studied the longer-term effects of SLIT, that is, 1, 2, and 3 years after the start of the first treatment course. SLIT did not significantly alter serum RGP-specific IgE levels (Figure 4A). RGP-specific IgG₂ levels increased after a total of 3 courses of SLIT (Figure 4A). RGP-specific IgG₄ increased after each consecutive course of treatment at 1, 2, and 3 years (Figure 4A). Similar to RGP-specific IgE and IgG₂ antibodies, frequencies of IgE⁺ memory B cells were unchanged by SLIT, while IgG₂⁺ memory B cells were significantly increased 3 years after commencing SLIT (Figure 4B). Frequencies of IgG₄⁺ memory B cells were increased 2 years after commencing SLIT.

3.5 | Molecular analysis of Ig gene rearrangements

Given that SLIT increased IgG₂ and IgG₄ antibodies and memory B-cell proportions, we investigated whether these changes were reflected in the proportions of unique IgG transcripts from blood B cells for a subgroup of 5 participants with an increased percentage of IgG₂⁺ memory B cells after SLIT. Median frequencies of somatic hypermutations did not differ between the proximal IgG₁ and the distal IgG₂ subclasses (Figure 5A and B) nor were significantly different after 3 years SLIT. However, after 3 years SLIT the relative usage of the IgG₂ and IgG₄ subclasses were significantly increased at the expense of IgG₁ (Figure 5C). Taken together, these data demonstrate that repeat courses of SLIT for grass pollen allergy induce allergen-specific IgG₂ and IgG₄ responses, evidenced by an increase in IgG₂⁺ and IgG₄⁺ B-cell proportions and skewing toward unique IgG₂ and IgG₄ transcripts.

4 | DISCUSSION

We here report that SLIT for grass pollen allergy not only has long-term beneficial clinical effects, but also results in sustained systemic effects on the immune system. SLIT induced a rapid and prolonged increase in RGP-specific serum IgG₄ accompanied by an increase in the frequency of peripheral blood IgG₄⁺ memory B cells. Furthermore, repeat courses of SLIT resulted in a similar increase in RGP-specific IgG₂ in serum corresponding with increased frequency of IgG₂⁺ memory B cells in the blood.

Currently, grass pollen SLIT is recommended as a preseasonal and co-seasonal course starting 4 months prior to the hay fever season, confirmed by meta-analyses as clinically effective.² Yet, long-term treatment regimens are costly and discourage treatment adherence.⁵¹ As patients are exposed to grass pollens during the spring season, we reasoned that a 4 months preseasonal treatment regimen would avoid the risk of adding to excessive and

unpredictable allergen loads during the Melbourne Spring. Based on our analysis of symptom scores, this approach is highly effective.^{42,43} Prolonged treatment (duration > 12 months) is known to have beneficial effects on symptom and medication scores.⁵² The fact that some immunological effects are delayed, only occurring after the second or third treatment year as observed for serum RGP-specific IgG₂ levels, or continuing to rise after consecutive treatment as for serum RGP-specific IgG₄ levels, supports these premises.

In particular, we observed a marked increase in RGP-specific IgG. Previously, allergen-specific immunotherapy, either SCIT or SLIT, has already been demonstrated to result in increased allergen-specific IgG₄ serum levels.^{53,54} Increased allergen-specific IgG₄ has been postulated as one of the explanations for the beneficial effects of immunotherapy and has been observed as a natural effect in beekeepers exposed to bee venom for prolonged periods,⁵⁵ yet the exact desensitizing effect of specific IgG₄ in immunotherapy remains unclear. Allergen-specific IgG₄ can competitively inhibit IgE from binding to allergens and may subsequently reduce allergic responses by preventing FcεR-mediated activation of granulocytes.⁵⁶ Furthermore, IgG₄ antibody has been proposed to inhibit inflammatory responses by preventing C1q complement activation and binding to the inhibitory receptor FCγRIIb (CD32b).^{57,58}

SLIT also increased RGP-specific IgG₂ after three consecutive courses of SLIT. This suggests repeated or high-dose exposure to RGP from SLIT is required to enhance RGP-specific IgG₂ beyond that which is generated from annual RGP exposure during the pollen season. Furthermore, sublingual administration of RGP may have preferentially induced an IgG₂ response not seen from environmental exposure through the airway.

The immune mechanisms by which allergen-specific IgG₂ may contribute to the benefits of immunotherapy remain unclear. IgG₂ has been shown to inhibit histamine release from basophils by activating FCγRIIb and may reduce allergic symptoms by this mechanism.⁵⁹ In a similar manner to IgG₄, IgG₂ may also bind allergen and prevent effector cell degranulation by masking IgE epitopes.

The source of increased serum IgG₂ and IgG₄ levels is IgG₂⁺ and IgG₄⁺-producing plasma cells, respectively. As the majority of serum IgG is produced by bone marrow residing plasma cells, we were unable to assess these. However, we did assess the more immature plasmablasts in blood, finding that their frequencies were not affected by SLIT. Further characterization of Ig isotypes and IgG subclasses was not possible due to our approach for membrane staining, since the majority of plasmablasts (especially those producing IgG) lack surface Ig expression. Hence, we focused on the analysis of memory B cells, which are abundant in the blood due to their circulatory nature, and their capacity to quickly differentiate into plasma cells in subsequent antigen responses.⁶⁰

We observed that SLIT drives increased frequencies of IgG₂⁺ and IgG₄⁺ memory B cells, whereas there was no effect on frequencies of IgE⁺ memory B-cell subsets. The latter observation can explain the absence of a decline in IgE serum levels as also demonstrated

by others.^{52,61,62} Since we observed the increase in IgG₄⁺ memory B cells after 4 months of treatment with SLIT, and before the pollen season, this effect can be directly attributed to the treatment with Oralair®. Our observation that frequencies of IgG₄⁺ memory B cells remain increased for at least 3 years can be an explanation for the long-lasting effects attributed to immunotherapy.⁵⁻⁷

Allergen-specific IgG₂ and IgG₄ appear to be robust markers of repeated allergen exposure. Increases in allergen-specific IgG₂ and IgG₄ alongside increased frequency of IgG₂⁺ and IgG₄⁺ memory B cells may arise from class switching of allergen-specific IgG₁⁺ memory B cells upon repeated exposure to allergen. Sequential Ig class switching can only occur 5'–3' along the *IGH* locus, which is arranged in the following order: IgG₃ > IgG₁ > IgG₂ > IgG₄ > IgE > IgA (Figure 5A). As such, IgG₁⁺ B cells may switch to IgG₂ or IgG₄ but not vice versa. In a study of allergen-specific antibodies in children from birth to 10 years old, IgE responses to aeroallergens were typically preceded by IgG.⁶³ Allergen-specific IgG⁺ memory B cells may therefore provide a reservoir for switching to IgE that can be induced by repeated exposure to the allergen and subsequently cause allergic sensitization. A similar pathway may give rise to allergen-specific IgG₂ and IgG₄, whereby allergen exposure promotes class switching of allergen-specific IgG₁⁺ memory B cells to IgG₂ and IgG₄. Switching to IgG₂ and IgG₄ may help explain why many allergies subside with age, perhaps due to repeated allergen exposure throughout childhood. However, it remains to be determined whether allergen-specific IgG₂ and IgG₄ are indispensable for generating clinical benefit by SLIT in lieu of T cell-mediated tolerance.

In line with previous reports, we observed that proliferation of Treg from patients after SLIT was increased in response to in vitro stimulation with RGP. Previous studies have observed new generation of allergen-specific Treg, as well as clonal expansion of allergen-specific Treg in response to AIT.^{56,64} Furthermore, we observed that SLIT increased IL-10 production from RGP-stimulated PBMC, whereas IL-13 was diminished. The cytokines IL-4 and IL-13 promote Ig class switch recombination (CSR) to IgE, and these are predominantly produced by Th2 cells. In addition to Th2 cytokines, CSR to IgG₄ is also regulated by IL-10 which is predominantly secreted by Treg. Our data are consistent with SLIT-induced proliferation of allergen-specific Treg, whereby increased IL-10 production induces Ig class switching of allergen-specific B cells to IgG₄. B regulatory cells (Breg) and monocyte-derived macrophages may also be an alternative source of IL-10 in our in vitro assay, further enhancing IgG₄ class switching in response to RGP.^{65,66}

In conclusion, our data provide evidence for long-lasting effects of allergen SLIT on the memory compartment of the immune system. Increased Treg frequencies and increased IL-10 production were associated with increased frequency of IgG₄⁺ memory B cells and a beneficial shift in the IgG₄⁺/IgE⁺ memory B-cell ratio, reflecting the increased IgG₄/IgE antibody fraction in serum and resultant clinically favorable outcome. Moreover, to our knowledge, our study is the first to demonstrate increases in memory B cells expressing IgG₂

or IgG₄ following allergen immunotherapy. As IgG₂ and IgG₄ have anti-inflammatory properties and are induced following repeated antigen-exposure,⁴¹ this B-cell memory compartment is a potential mechanism by which allergen immunotherapy modifies the natural course of disease. In future studies, it would therefore be of interest to examine the functional properties of allergen-specific B cells, as well as the effector functions of allergen-specific IgG subclasses.⁶⁷

ACKNOWLEDGMENTS

The authors gratefully acknowledge Ms Kirsten Deckert for collection of clinical data and blood samples from patients, and Ms Tracy Phan for technical assistance. We are also grateful to the AMREP flow cytometry team for technical assistance. The studies were financially supported by a grant from the Num Pon Soon Charitable Trust, grant S698 from the Sophia Children's Hospital Fund (SKF) and an NHMRC Senior Research Fellowship GNT1117687 to MCvZ.

CONFLICT OF INTEREST

All authors declare that no conflict of interest exists.

ORCID

Jorn J. Heeringa  <https://orcid.org/0000-0003-0304-8977>

Craig I. McKenzie  <https://orcid.org/0000-0001-7070-620X>

Nirupama Varese  <https://orcid.org/0000-0001-9074-3710>

Mark Hew  <https://orcid.org/0000-0002-7498-0000>

Pei M. Aui  <https://orcid.org/0000-0002-2314-9989>

Jennifer M. Rolland  <https://orcid.org/0000-0002-7891-983X>

Robyn E. O'Hehir  <https://orcid.org/0000-0002-3489-7595>

Menno C. van Zelm  <https://orcid.org/0000-0003-4161-1919>

REFERENCES

- Björkstén B, Clayton T, Ellwood P, Stewart A, Strachan D, Group IPIS. Worldwide time trends for symptoms of rhinitis and conjunctivitis: phase III of the international study of asthma and allergies in childhood. *Pediatr Allergy Immunol*. 2008;19(2):110-124.
- Di Bona D, Plaia A, Leto-Barone MS, La Piana S, Di Lorenzo G. Efficacy of grass pollen allergen sublingual immunotherapy tablets for seasonal allergic rhinoconjunctivitis: a systematic review and meta-analysis. *JAMA Intern Med*. 2015;175(8):1301-1309.
- Di Bona D, Plaia A, Scafidi V, Leto-Barone MS, Di Lorenzo G. Efficacy of sublingual immunotherapy with grass allergens for seasonal allergic rhinitis: a systematic review and meta-analysis. *J Allergy Clin Immunol*. 2010;126(3):558-566.
- Devillier P, Molimard M, Ansolabehere X, et al. Immunotherapy with grass pollen tablets reduces medication dispensing for allergic rhinitis and asthma: A retrospective database study in France. *Allergy*. 2019;74(7):1317-1326.
- Durham SR, Walker SM, Varga EM, et al. Long-term clinical efficacy of grass-pollen immunotherapy. *N Engl J Med*. 1999;341(7):468-475.
- Durham SR, Emminger W, Kapp A, et al. Long-term clinical efficacy in grass pollen-induced rhinoconjunctivitis after treatment with SQ-standardized grass allergy immunotherapy tablet. *J Allergy Clin Immunol*. 2010;125(1):131-138.
- Durham SR, Emminger W, Kapp A, et al. SQ-standardized sublingual grass immunotherapy: confirmation of disease modification 2 years after 3 years of treatment in a randomized trial. *J Allergy Clin Immunol*. 2012;129(3):717-725.
- Pajno GB, Barberio G, De Luca F, Morabito L, Parmiani S. Prevention of new sensitizations in asthmatic children monosensitized to house dust mite by specific immunotherapy. A six-year follow-up study. *Clin Exp Allergy*. 2001;31(9):1392-1397.
- Moller C, Dreborg S, Ferdousi HA, et al. Pollen immunotherapy reduces the development of asthma in children with seasonal rhinoconjunctivitis (the PAT-study). *J Allergy Clin Immunol*. 2002;109(2):251-256.
- Larche M, Akdis CA, Valenta R. Immunological mechanisms of allergen-specific immunotherapy. *Nat Rev Immunol*. 2006;6(10):761-771.
- Akdis CA, Akdis M. Advances in allergen immunotherapy: aiming for complete tolerance to allergens. *Sci Transl Med*. 2015;7(280):280ps6-280ps6.
- Stone KD, Prussin C, Metcalfe DD. IgE, mast cells, basophils, and eosinophils. *J Allergy Clin Immunol*. 2010;125(2 Suppl 2):S73-S80.
- Geha RS, Jabara HH, Brodeur SR. The regulation of immunoglobulin E class-switch recombination. *Nat Rev Immunol*. 2003;3(9):721-732.
- Takatsu K, Kouro T, Nagai Y. Interleukin 5 in the link between the innate and acquired immune response. *Adv Immunol*. 2009;101:191-236.
- Romagnani S. Immunologic influences on allergy and the TH1/TH2 balance. *J Allergy Clin Immunol*. 2004;113(3):395-400.
- Ebner C, Siemann U, Bohle B, et al. Immunological changes during specific immunotherapy of grass pollen allergy: reduced lymphoproliferative responses to allergen and shift from TH2 to TH1 in T-cell clones specific for Phl p 1, a major grass pollen allergen. *Clin Exp Allergy*. 1997;27(9):1007-1015.
- Gardner LM, O'Hehir RE, Rolland JM. High dose allergen stimulation of T cells from house dust mite-allergic subjects induces expansion of IFN-gamma+ T Cells, apoptosis of CD4+IL-4+ T cells and T cell anergy. *Int Arch Allergy Immunol*. 2004;133(1):1-13.
- Francis JN, Till SJ, Durham SR. Induction of IL-10+CD4+CD25+ T cells by grass pollen immunotherapy. *J Allergy Clin Immunol*. 2003;111(6):1255-1261.
- Mobs C, Slotoch C, Löffler H, Jakob T, Hertl M, Pfützner W. Birch pollen immunotherapy leads to differential induction of regulatory T cells and delayed helper T cell immune deviation. *J Immunol*. 2010;184(4):2194-2203.
- Varona R, Ramos T, Escribese MM, et al. Persistent regulatory T-cell response 2 years after 3 years of grass tablet SLIT: Links to reduced eosinophil counts, sIgE levels, and clinical benefit. *Allergy*. 2019;74(2):349-360.
- Gardner LM, Thien FC, Douglass JA, Rolland JM, O'Hehir RE. Induction of T 'regulatory' cells by standardized house dust mite immunotherapy: an increase in CD4+ CD25+ interleukin-10+ T cells expressing peripheral tissue trafficking markers. *Clin Exp Allergy*. 2004;34(8):1209-1219.
- Wachholz PA, Soni NK, Till SJ, Durham SR. Inhibition of allergen-IgE binding to B cells by IgG antibodies after grass pollen immunotherapy. *J Allergy Clin Immunol*. 2003;112(5):915-922.
- Scadding GW, Shamji MH, Jacobson MR, et al. Sublingual grass pollen immunotherapy is associated with increases in sublingual Foxp3-expressing cells and elevated allergen-specific immunoglobulin G4, immunoglobulin A and serum inhibitory activity for immunoglobulin E-facilitated allergen binding to B cells. *Clin Exp Allergy*. 2010;40(4):598-606.
- O'Hehir RE, Gardner LM, de Leon MP, et al. House dust mite sublingual immunotherapy: the role for transforming growth

- factor-beta and functional regulatory T cells. *Am J Respir Crit Care Med*. 2009;180(10):936-947.
25. Akdis CA, Akdis M. Mechanisms of allergen-specific immunotherapy and immune tolerance to allergens. *World Allergy Organ J*. 2015;8(1):17.
 26. Royer B, Varadaradjalou S, Saas P, Guillosson JJ, Kantelip JP, Arock M. Inhibition of IgE-induced activation of human mast cells by IL-10. *Clin Exp Allergy*. 2001;31(5):694-704.
 27. Schandene L, Alonso-Vega C, Willems F, et al. B7/CD28-dependent IL-5 production by human resting T cells is inhibited by IL-10. *J Immunol*. 1994;152(9):4368-4374.
 28. Ohkawara Y, Lim KG, Xing Z, et al. CD40 expression by human peripheral blood eosinophils. *J Clin Invest*. 1996;97(7):1761-1766.
 29. Jeannin P, Lecoanet S, Delneste Y, Gauchat JF, Bonnefoy JY. IgE versus IgG4 production can be differentially regulated by IL-10. *J Immunol*. 1998;160(7):3555-3561.
 30. Santos AF, James LK, Bahnson HT, et al. IgG4 inhibits peanut-induced basophil and mast cell activation in peanut-tolerant children sensitized to peanut major allergens. *J Allergy Clin Immunol*. 2015;135(5):1249-1256.
 31. Di Bona D, Plaia A, Leto-Barone MS, La Piana S, Di Lorenzo G. Efficacy of subcutaneous and sublingual immunotherapy with grass allergens for seasonal allergic rhinitis: a meta-analysis-based comparison. *J Allergy Clin Immunol*. 2012;130(5):1097-1107.
 32. Bohle B, Kinaciyan T, Gerstmayr M, Radakovics A, Jahn-Schmid B, Ebner C. Sublingual immunotherapy induces IL-10-producing T regulatory cells, allergen-specific T-cell tolerance, and immune deviation. *J Allergy Clin Immunol*. 2007;120(3):707-713.
 33. Suarez-Fueyo A, Ramos T, Galan A, et al. Grass tablet sublingual immunotherapy downregulates the TH2 cytokine response followed by regulatory T-cell generation. *J Allergy Clin Immunol*. 2014;133(1):130-138.
 34. Bahceciler NN, Arikian C, Taylor A, et al. Impact of sublingual immunotherapy on specific antibody levels in asthmatic children allergic to house dust mites. *Int Arch Allergy Immunol*. 2005;136(3):287-294.
 35. Sugimoto M, Kamemura N, Nagao M, et al. Differential response in allergen-specific IgE, IgGs, and IgA levels for predicting outcome of oral immunotherapy. *Pediatr Allergy Immunol*. 2016;27(3):276-282.
 36. Rolinck-Werninghaus C, Kopp M, Liebke C, Lange J, Wahn U, Niggemann B. Lack of detectable alterations in immune responses during sublingual immunotherapy in children with seasonal allergic rhinoconjunctivitis to grass pollen. *Int Arch Allergy Immunol*. 2005;136(2):134-141.
 37. Dehlink E, Eiwegger T, Gerstmayr M, et al. Absence of systemic immunologic changes during dose build-up phase and early maintenance period in effective specific sublingual immunotherapy in children. *Clin Exp Allergy*. 2006;36(1):32-39.
 38. Berkowska MA, Driessen G, Bikos V, et al. Human memory B cells originate from three distinct germinal center-dependent and -independent maturation pathways. *Blood*. 2011;118(8):2150-2158.
 39. Jackson K, Wang Y, Collins AM. Human immunoglobulin classes and subclasses show variability in VDJ gene mutation levels. *Immunol Cell Biol*. 2014;92(8):729-733.
 40. van Zelm MC. B cells take their time: sequential IgG class switching over the course of an immune response? *Immunol Cell Biol*. 2014;92(8):645-646.
 41. de Jong BG, IJsepeert H, Marques L, et al. Human IgG2- and IgG4-expressing memory B cells display enhanced molecular and phenotypic signs of maturity and accumulate with age. *Immunol Cell Biol*. 2017;95(9):744-752.
 42. O'Hehir RE, Varese NP, Deckert K, et al. Epidemic thunderstorm asthma protection with five-grass pollen tablet sublingual immunotherapy: a clinical trial. *Am J Respir Crit Care Med*. 2018;198(1):126-128.
 43. O'Hehir R, Heeringa JJ, Deckert K, Rolland JM, van Zelm MC, Hew M. Preseasonal grass pollen SLIT in at risk individuals confers protection from epidemic thunderstorm asthma [abstract]. *Allergy*. 2017;72(Suppl S103):759.
 44. Davies JM, Bright ML, Rolland JM, O'Hehir RE. Bahia grass pollen specific IgE is common in seasonal rhinitis patients but has limited cross-reactivity with Ryegrass. *Allergy*. 2005;60(2):251-255.
 45. Berkowska MA, Heeringa JJ, Hajdarbegovic E, et al. Human IgE+ B cells are derived from T cell-dependent and T cell-independent pathways. *J Allergy Clin Immunol*. 2014;134(3):688-697.
 46. Heeringa JJ, Karim AF, van Laar J, et al. Expansion of blood IgG4(+) B, TH2, and regulatory T cells in patients with IgG4-related disease. *J Allergy Clin Immunol*. 2018;141(5): 1831-1843.
 47. Tiller T, Meffre E, Yurasov S, Tsuiji M, Nussenzweig MC, Wardemann H. Efficient generation of monoclonal antibodies from single human B cells by single cell RT-PCR and expression vector cloning. *J Immunol Methods*. 2008;329(1-2):112-124.
 48. Haahtela T, Burbach GJ, Bachert C, et al. Clinical relevance is associated with allergen-specific wheal size in skin prick testing. *Clin Exp Allergy*. 2013;44(3):407-416.
 49. Ciprandi G, Tosca MA, Capasso M. Exhaled nitric oxide in children with allergic rhinitis and/or asthma: a relationship with bronchial hyperreactivity. *J Asthma*. 2010;47(10):1142-1147.
 50. Verini M, Consilvio NP, Di Pillio S, et al. FeNO as a marker of airways inflammation: the possible implications in childhood asthma management. *J Allergy*. 2010;2010:1-7.
 51. Lin SY, Ereksom N, Kim JM, et al. Sublingual immunotherapy for the treatment of allergic rhinoconjunctivitis and asthma: a systematic review. *JAMA*. 2013;309(12):1278-1288.
 52. Wilson DR, Lima MT, Durham SR. Sublingual immunotherapy for allergic rhinitis: systematic review and meta-analysis. *Allergy*. 2005;60(1):4-12.
 53. Jutel M, Jaeger L, Suck R, Meyer H, Fiebig H, Cromwell O. Allergen-specific immunotherapy with recombinant grass pollen allergens. *J Allergy Clin Immunol*. 2005;116(3):608-613.
 54. Muller U, Helbling A, Bischof M. Predictive value of venom-specific IgE, IgG and IgG subclass antibodies in patients on immunotherapy with honey bee venom. *Allergy*. 1989;44(6):412-418.
 55. Aalberse RC, van der Gaag R, van Leeuwen J. Serologic aspects of IgG4 antibodies. I. Prolonged immunization results in an IgG4-restricted response. *J Immunol*. 1983;130(2):722-726.
 56. Wachholz PA, Durham SR. Mechanisms of immunotherapy: IgG revisited. *Curr Opin Allergy Clin Immunol*. 2004;4(4):313-318.
 57. Anderson CL, Abraham GN. Characterization of the Fc receptor for IgG on a human macrophage cell line, U937. *J Immunol*. 1980;125(6):2735-2741.
 58. Lienthal G-M, Rahmöller J, Petry J, Bartsch YC, Leliavski A, Ehlers M. Potential of murine IgG1 and human IgG4 to inhibit the classical complement and Fcγ receptor activation pathways. *Front Immunol*. 2018;9:958-958.
 59. MacGlashan D Jr, Hamilton RG. Parameters determining the efficacy of CD32 to inhibit activation of FcεRI in human basophils. *J Allergy Clin Immunol*. 2016;137(4):1256-1258.
 60. Tarlinton D, Good-Jacobson K. Diversity among memory B cells: origin, consequences, and utility. *Science*. 2013;341(6151): 1205-1211.
 61. Vourdas D, Syrigou E, Potamianou P, et al. Double-blind, placebo-controlled evaluation of sublingual immunotherapy with standardized olive pollen extract in pediatric patients with allergic rhinoconjunctivitis and mild asthma due to olive pollen sensitization. *Allergy*. 1998;53(7):662-672.
 62. Lima MT, Wilson D, Pitkin L, et al. Grass pollen sublingual immunotherapy for seasonal rhinoconjunctivitis: a randomized controlled trial. *Clin Exp Allergy*. 2002;32(4):507-514.

63. Huang X, Tsilochristou O, Perna S, et al. Evolution of the IgE and IgG repertoire to a comprehensive array of allergen molecules in the first decade of life. *Allergy*. 2018;73(2):421-430.
64. Till SJ, Francis JN, Nouri-Aria K, Durham SR. Mechanisms of immunotherapy. *J Allergy Clin Immunol*. 2004;113(6):1025-1034.
65. van de Veen W. The role of regulatory B cells in allergen immunotherapy. *Curr Opin Allergy Clin Immunol*. 2017;17(6):447-452.
66. Bianchini R, Roth-Walter F, Ohradanova-Repic A, et al. IgG4 drives M2a macrophages to a regulatory M2b-like phenotype: potential implication in immune tolerance. *Allergy*. 2019;74(3):483-494.
67. van Zelm MC, McKenzie CI, Varese N, Rolland JM, O'Hehir RE. Recent developments and highlights in immune monitoring of allergen immunotherapy. *Allergy*. 2019. <https://doi.org/10.1111/all.14078>. [Epub ahead of print].

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section.

How to cite this article: Heeringa JJ, McKenzie CI, Varese N, et al. Induction of IgG₂ and IgG₄ B-cell memory following sublingual immunotherapy for ryegrass pollen allergy. *Allergy*. 2019;00:1-12. <https://doi.org/10.1111/all.14073>