



ORIGINAL ARTICLE



WILEY

High-dose posaconazole for azole-resistant aspergillosis and other difficult-to-treat mould infections

Alexander F. A. D. Schauwvlieghe^{1,2} | Jochem B. Buil^{3,4} | Paul E. Verweij^{3,4} | Rogier A. S. Hoek⁵ | Jan J. Cornelissen² | Nicole M. A. Blijlevens⁶ | Stefanie S. V. Henriët^{4,7} | Bart J. A. Rijnders¹ | Roger J. M. Brüggemann^{4,8}

¹Department of Internal Medicine, Section of Infectious Diseases, Erasmus MC University Medical Centre, Rotterdam, The Netherlands

²Department of Haematology, Erasmus MC Cancer Institute, Rotterdam, The Netherlands

³Department of Medical Microbiology, Radboud University Medical Centre, Nijmegen, The Netherlands

⁴Center of Expertise in Mycology Radboudumc/CWZ, Nijmegen, The Netherlands

⁵Department of Pulmonary Medicine, Erasmus MC, University Medical Centre Rotterdam, Rotterdam, The Netherlands

⁶Department of Haematology, Radboud University Medical Centre, Nijmegen, Netherlands

⁷Department of Paediatric Infectious Diseases and Immunology, Radboud Institute for Molecular Life Sciences, Amalia Children's Hospital, Radboud University Medical Centre, Nijmegen, The Netherlands

⁸Department of Pharmacy, Radboud University Medical Centre, Nijmegen, The Netherlands

Correspondence

Alexander F.A.D. Schauwvlieghe, Department of Internal Medicine, Section of Infectious Diseases, Erasmus MC University Medical Centre, Rotterdam, The Netherlands
Email: a.schauwvlieghe@erasmusmc.nl

Funding information

This study was performed as part of our routine work.

Summary

Background: Oral follow-up therapy is problematic in moulds with reduced azole-susceptibility, such as azole-resistant *Aspergillus fumigatus* infection. Currently, only intravenous liposomal amphotericin B (L-AmB) is advocated by guidelines for the treatment of azole-resistant aspergillosis infections. Preclinical research indicates that high-dose posaconazole (HD-POS) might be a feasible option provided that high drug exposure (ie POS serum through levels >3 mg/L) can be achieved and is safe.

Objectives: To describe our experience with the use of oral HD-POS as treatment strategies for patients infected with pathogens with a POS MIC close to the clinical breakpoint.

Patients/Methods: We review evidence supporting the use of HD-POS and describe our experience on safety and efficacy in 16 patients. In addition, we describe the adverse events (AE) observed in 25 patients with POS concentrations at the higher end of the population distribution during treatment with the licensed dose.

Results: Sixteen patients were treated intentionally with HD-POS for voriconazole-resistant invasive aspergillosis (7/16), mucormycosis (4/16), salvage therapy for IA (4/16) and IA at a sanctuary site (spondylodiscitis) in 1. Grade 3-4 AEs were observed in 6, and all of them were considered at least possibly related. Grade 3-4 AEs were observed in 5 of the 25 patients with spontaneous high POS serum through levels considered at least possibly related using Naranjo scale.

Conclusions: High-dose posaconazole is a treatment option if strict monitoring for both exposure and for AE is possible.

KEYWORDS

antifungal susceptibility, antimycotic chemotherapy, aspergillosis

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2019 The Authors. *Mycoses* published by Blackwell Verlag GmbH

Schauwvlieghe, Buil, Rijnders and Brüggemann contributed equally as first and last author.

1 | INTRODUCTION

Invasive aspergillosis (IA) in patients with haematological malignancies is associated with a mortality of 20%-30%.^{1,2} Triazole resistance is increasingly reported in different countries through culture-based surveillance studies³ and is associated with a much higher mortality of 50%-88%.^{4,5} In 2015, a consensus meeting on the management of azole-resistant IA was organised⁶ and liposomal amphotericin B (L-AmB) was advocated as the preferred therapy but has obvious toxicity limitations and can only be administered intravenously. Treatment of IA has to be continued for a minimum of 6-12 weeks but occasionally much longer.⁷ Other treatment options are therefore urgently needed. Phase II studies on new antifungals are just about to start and subsequent phase III studies typically take 3 or 4 years to complete. Therefore, these drugs will not provide a short term solution. Targeting high-exposure posaconazole (POS) may be a potential oral step-down treatment option for azole-resistant IA and other difficult-to-treat mould infections.

Posaconazole is approved in patients with haematological malignancies both for prophylaxis and treatment of refractory IA or when intolerance to first-line agents occurs.^{8,9} The agent is available as oral suspension, a delayed-release tablet and an intravenous formulation. Oral absorption of POS oral suspension is affected by food and gastric pH. In contrast, POS tablets contain the active drug mixed with a pH-sensitive polymer¹⁰ and this polymer releases the drug in the intestines, causing threefold increased exposures compared to POS oral suspension.¹¹

Therapeutic drug monitoring (TDM) has been widely implemented to assess therapeutic efficacy of POS oral suspension but its usefulness is in a state of flux following the introduction

of the new POS formulations specifically in the setting of prophylaxis.¹²⁻¹⁴ Current guidelines recommend a Ctrough concentration of ≥ 0.7 mg/L for prophylaxis and >1.0 mg/L for primary and >1.25 mg/L for salvage therapy,¹⁵ although these concentrations were determined independent of the susceptibility of the infecting pathogen.¹³

These targets have been derived for susceptible pathogens and are not valid for pathogens with attenuated susceptibilities. A different approach is needed to optimise treatment in case of reduced susceptibility.

Preclinical research indicates that high-dose posaconazole (HD-POS) might be a feasible option provided that high drug exposure (ie POS serum through levels >3 mg/L) can be achieved and is safe. Hence, we argued that oral high-dose treatment strategies might be feasible to treat pathogens with relatively low MICs/MICs just above the clinical breakpoint (low-resistant). Human data on the treatment of pathogens with reduced susceptibility as well as safety of POS Ctrough concentrations of >3 mg/L are sparse.

Here, we review the evidence supporting the use of HD-POS and describe our experience on safety and efficacy in 16 patients. In addition, we describe the adverse events (AE) observed in 25 patients with POS concentrations at the higher end of the population distribution during treatment with the licensed dose.

2 | PATIENTS/METHODS

We set out to explore safety of HD-POS and retrospectively collected clinical and laboratory data of patients from 2 Dutch academic medical centres (Erasmus University Medical Centre,

Adverse event	Grade 1	Grade 2	Grade 3	Grade 4	Grade 5
Diarrhoea	1				
Nausea		1			
Vomiting	3				
Increased hepatic enzymes	4	1	1 ³	2 ^{5,7}	
Cardiac troponin T increased			1 ⁶		
Electrocardiogram QTc corrected interval prolonged	1	1	1 ⁶		
Leukopenia				1 ⁴	
Hypokalaemia	3	3			
Hyperkalaemia	1				
Headache		1			
Delirium	1		1 ²		
Alopecia	1				
Hypertension		2			
Heart failure			1 ⁵		
Rash	1				

TABLE 1 Adverse events of 16 patients receiving intentionally HD-POS graded accordingly to the Common Terminology Criteria for Adverse Events (version 4.03)

Note: Digits refer to the number of patients in whom these AEs have been documented.

Prolongation in the QTc interval was assessed by comparing electrocardiograms obtained at baseline and during HD-POS treatment, if available.

Naranjo¹⁶ adverse drug reaction probability scale: >9 : definite, 5 to 8: probable, 1-4: possible. -3 to 0: doubtful.

TABLE 2 Underlying condition, IFD, *A fumigatus* genotype and phenotype, and outcome in 16 patients treated with high-dose posaconazole (HD-POS)

Patient	Age (years)	Underlying disease	IFD, classification	Reason HD-POS	Sample with culture
1	69	Mixed dust pneumoconiosis	CPA	Resistant strain	Sputum: <i>A fumigatus</i>
2	51	AML, AlloTx	IPA, probable	Resistant strain	No positive culture
3	18	ALL	IPA, proven (cerebral)	Resistant strain	Sputum: <i>A fumigatus</i>
4	46	SOT (kidney), PTLD	IPA, probable	Resistant strain	BAL: <i>A fumigatus</i>
5	69	AML	IPA, probable	Resistant strain	No positive culture
6	61	No relevant	CPA	Resistant strain	BAL: <i>A fumigatus</i>
7	32	SOT (lung)	Pulmonary mucormycosis, proven	Mucormycosis	Lung: <i>Rhizopus</i> species
8	17	ALL	IPA, probable	Mixed infection (R/S)	BAL: <i>A fumigatus</i> R and S
9	50	AML, AlloTx	Mucormycosis, probable	Mucormycosis	No positive culture
10	58	SLE with pancytopenia	Mucormycosis, proven	Mucormycosis	Liver biopsy: microscopy: hyphy. No positive culture. Spleen biopsy PCR positive
11	67	DM type II	Mucormycosis, probable (skin)	Mucormycosis	Tissue sample wound: <i>Rhizopus oryzae</i>
12	2	ALL	Mucormycosis, proven	Mucormycosis	Multiple skin biopsies: <i>Lichtheimia corymbifera</i>
13	50	No relevant	IA, proven	Sanctuary sites infection	Spinal biopsy: <i>A fumigatus</i>
14	68	AML	IPA, probable	Salvage	No positive culture
15	65	AML	IPA, probable	Salvage	Sputum: <i>A nidulans</i>
16	8	ALL	IPA, proven	Salvage <i>A terreus</i>	Lobectomy, lung tissue: <i>A terreus</i>

Note: Calculated target Ctrough based on the MIC is taken from Seyedmousavi et al²⁸

Abbreviations: AlloTx, allogeneic stem cell transplant; C, concentration; CPA, chronic pulmonary aspergillosis; HD-POS, high-dose posaconazole; IA, invasive aspergillosis; IFD, invasive fungal diseases; IPA, Invasive pulmonary aspergillosis; ISA, isavuconazole; ITZ, Itraconazole; POS, posaconazole; PTLD, post-transplant lymphoproliferative disease; R, resistant; S, Susceptible; SOT, solid organ transplantation; VCZ, voriconazole.

^aMIC was determined according to the EUCAST method for susceptibility testing of moulds (version 9.2). Patients were classified following the revised definitions of the European Organization for Research and Treatment of Cancer/Mycosis Study Group (EORTC/MSG).⁴²

^bMIC was determined according to the CLSI method for susceptibility testing of moulds(M38-A2)

^cthis patient was included because the patient was treated with POS 400 mg BID despite the low Ctrough level.

Rotterdam and Radboud University Medical Centre, Nijmegen) in which POS Ctrough concentrations >3 mg/L had been documented in two different populations. All patients were in care by one of the authors of this paper. Data were extracted and reviewed by JB and AS Group 1 consisted of patients intentionally treated with HD-POS targeting POS Ctrough concentrations >3 mg/L and Group 2 were patients that reached POS Ctrough concentrations >3 mg/L with the licensed dose. We focused on AEs (related or unrelated to POS) described in the patient files and laboratory data. Data from these patients were reviewed for toxicities according to the Common Terminology Criteria for Adverse Events (CTCAE) version 4.03. An AE was defined as unfavourable or unintended sign or symptom while the patient was treated with POS, whether or not the sign or symptom was related to POS. The Naranjo scale was used to determine for the assessment of causality of potential AE with POS. This is a questionnaire for determining a potential

likelihood that an adverse drug reaction is actually linked to a drug. Probability is assigned using a scoring system with the following possible results: definite, probable, possible of doubtful.¹⁶ Medians and 25th to 75th inter-quartile ranges were used for statistic descriptions. This type of research does not fall under the Dutch law of research on human subjects. However, to safeguard the privacy of the patients, the data were stored anonymously after data extraction.

3 | RESULTS

3.1 | Group 1

Sixteen patients were treated intentionally with HD-POS for voriconazole-resistant IA (7/16), mucormycosis (4/16), salvage

Aspergillus PCR result	MIC(mg/L) ^a				POS concentration: calculated target		Outcome
	ITZ	VCZ	POS	ISA	Highest C	Calculated Target	
TR ₄₆ /Y121F/T289A	>16	8	1	4	3.8	6.18-6.66	Alive
Y121F/T289A in BAL					6.1		Dead
TR ₃₄ /L98H	16	8	2	8	6	>10	Alive
TR ₃₄ /L98H	>16	4	0.5	8	0.2 ^c	3.09-3.33	Dead
TR ₃₄ /L98H in BAL					4		Dead
	>16	8	1	8	6.6	6.18-6.66	Alive
	1	8	0.25	1	3.8	1.44-1.55	Alive
	>16	4	0.5	8	5.6	3.09-3.33	Alive
					5.2		Alive
PCR: <i>Rizomucor pusillus</i>					5.0		Alive
	0.25	8	0.25	1	3.5	1.44-1.55	Alive
	0.5	16	0.5	>16	6.6	3.09-3.33	Alive
	0.25 ^b	0.25 ^b	0.063 ^b	0.5 ^b	3.6		Alive
					3.8		Dead
Wild-type <i>A. fumigatus</i> in BAL	0.25	0.25	0.25	0.5	3.1	1.44-1.55	Dead
Lung biopsy: <i>Aspergillus</i> Species	0.125	1	0.031	1	4.7		Alive

therapy for IA (4/16) and IA at a sanctuary site (spondylodiscitis) in 1. The median POS dose given was 600 (IQR 400-750) mg daily when the POS Ctrough concentrations of > 3 mg/L were reached after a median of 8 (IQR 6.40) days. Ten patients had significantly higher Ctrough concentration (above 4 mg/L) and 6 patients had Ctrough concentrations between 3.0 and 4.0 mg/L and on average patients had these concentrations for a median 76 days (IQR 20-162). Thirteen patients received POS tablet, 1 patient posaconazole-oral suspension (POS-OS) and 2 patients a combination of formulations. AEs are described in Table 1. Grade 3-4 AEs were observed in 6 patients, and all of them were considered at least possibly related using Naranjo scale. In 3 out of 16 patients, the treatment was stopped following an AE: arterial hypertension (grade 2), QTc prolongation, cardiac troponin T increased and left ventricular failure (grade 3) and leukocytopenia (grade 4).

3.2 | Efficacy

Of the 7 patients with azole-resistant IA treated with HD-POS, 4 survived while 3 died from their underlying disease but unrelated to the IA. In two patients, HD-POS was used as salvage therapy. One patient with IA caused by *Aspergillus terreus* was treated with HD-POS because serum galactomannan levels increased under conventional dosage which is a predictor of poor outcome (Table 2). All patients with mucormycosis survived.

3.3 | Group 2

This group consisted of 25 patients. The median POS Ctrough concentration was 4.3 mg/L (IQR 3.5-6.0). 19, 5 and 1 patient received POS tablet, POS-OS and the IV formulation, respectively.

	Grade 1	Grade 2	Grade 3	Grade 4	Grade 5
Adverse event					
Diarrhoea	4				
Nausea	4				
Vomiting	2				
Increased hepatic enzymes	2	3	1 ⁴		
Electrocardiogram QT corrected interval prolonged	2	1			
GGT increased		1			
Anorexia	5	1			
Hyponatremia	2		1 ¹		
Hypokalaemia	7	1			
Headache	5				
Seizure		1			
Alopecia	2				
Hypertension		2	1 ⁷	1 ⁴	
Hypotension					1 ⁷
Rash	3				

Note: Digits refer to the number of patients in whom these AEs have been documented.

Refractory shock, rapidly fatal. Distributive shock most likely according to treating physician.

These grade 3 or 4 AE were considered at least possible related to POS.

Naranjo¹⁶ adverse drug reaction probability scale: >9: definite, 5 to 8: probable, 1-4: possible. -3 to 0: doubtful.

TABLE 3 Adverse events of 25 patients receiving POS with high spontaneous concentration graded accordingly to the Common Terminology Criteria for Adverse Events(version 4.03)

Posaconazole was given to 18 and 7 patients for prophylaxis and treatment, respectively. All observed AEs are described in Table 3. The most frequently observed AE were hypokalaemia in eight patients and neurological in six patients (headache, convulsions). Grade 3-4 AEs were observed in 5, and all of them were considered at least possibly related using Naranjo scale. In 8 of the 25 patients, the dosage was reduced. Follow-up Ctrough concentrations were between 1.1 and 4.3 mg/L after dosage reduction.


4 | DISCUSSION

Little is known about the toxicity of patients attaining high POS Ctrough of >3 mg/L. The upper boundary level of average POS serum concentrations of 3.75 mg/L is set by the European Medicines Agency based on experience with the POS-OS and preclinical toxicology findings.¹⁷ In this study, we reviewed the safety and tolerability of HD-POS. In both group 1 and group 2, three patients were seen with a combination of hypertension and hypokalaemia that required antihypertensive therapy and potassium supplementation. The most striking case was a child treated with POS, L-AmB and micafungin for a proven aspergillosis following surgical removal of *Aspergillus* lesions in the spleen, left lung and right kidney. This patient developed several hypertensive crises and developed hypokalaemia for which oral supplementation was needed. 8 months after POS treatment, the

patient died due to a vasopressor refractory shock. During these hypertensive crises, aldosterone could not be measured (<50 pmol/L) and renine was within normal range. In retrospect, POS may have caused the hypertension and hypokalaemia. Recently, a case of POS-induced heart failure, hypertension and hypokalaemia was described with low renin and aldosterone levels. The inhibition of the enzyme 11-beta-hydroxysteroid dehydrogenase 2 is suggested as the potential mechanism causing apparent mineralocorticoid excess.¹⁸⁻²⁰ This enzyme is homeostatic regulator and damps mineralocorticoid activity by converting cortisol to cortisone.

The AE of HD-POS observed in this study are in line with previous reports of AE due to POS. A phase III study assessing PK and safety of POS tablet demonstrated that nausea and diarrhoea were the most common treatment-related AEs leading to POS discontinuation in 2% and 1%, respectively.²¹ Only 9 patients (10%) in this study attained an average Ctrough concentration between 2.5 and 3.75 mg/L, and six patients (3%) reached Ctrough concentrations ≥3.75 mg/L. No increase of AEs in patients with higher POS serum concentrations was observed but the study was not powered to detect such a relation. Very recently, PK and safety results from a phase 3 study of IV POS in patients at risk for invasive fungal disease were published. Six per cent of the patients had a steady-state concentration between >2.5 and ≤3.65 mg/L without signs of toxicity.²² In a retrospective analysis of 64 patients receiving POS tablet as prophylaxis, median POS steady-state concentrations of 1.67 mg/L (0.52-3.83 mg/L) were documented. In 21% of the patients, a QT_c

FIGURE 1 Posaconazole minimal inhibitory concentrations (MICs) distributions of most common Mucorales species: *Rhizopus oryzae*, *Mucor circinelloides*, *Rhizopus microspores* and *Lichtheimia corymbifera*. MICs were extracted from Espinel-Ingroff et al³⁶ MICs were determined according to the CLSI method for susceptibility testing of moulds (M38-A2)



prolongation was observed and the median steady-state concentration was 2.04 mg/L.²³ In a single-centre study, 343 courses of POS prophylaxis (IV or tablet) were assessed for safety and effectiveness. 20% of these patients developed liver injury, mostly hyperbilirubinemia but this is often multifactorial. More importantly, grade 3-4 elevations in hepatic enzymes were only observed in 2% of the patients without pre-existing liver injury with mostly spontaneous resolution despite treatment continuation.²⁴ Thus, in the current literature, information about the toxicity of high POS serum concentrations is limited but no increase in the number of AEs was observed in patients with higher than average serum concentrations.

4.1 | Azole-resistant IA

The large majority of azole-resistant *A. fumigatus* isolates harbour TR₃₄/L98H or TR₄₆/Y121F/T289A mutations in the *cyp51A*

gene,^{25,26} encoding the cytochrome p450 sterol 14 α -demethylase, the target of azoles. *A. fumigatus* isolates carrying resistance-associated mutations have high minimal inhibitory concentrations (MICs) for itraconazole and/or voriconazole as well as isavuconazole.²⁷ The MIC of POS often remains close to the susceptible population (ie MIC ≤ 0.5 to 1 mg/L).²⁸ MIC levels of POS >0.25 mg/L are considered resistant according to the EUCAST breakpoint, but this is based on population susceptibility and on concentrations achieved with the POS-OS at licensed dose. Indeed, drug exposure with POS-OS will marginally cover the *A. fumigatus* wild-type population, let alone low-level POS-resistant isolates. Higher exposures can be achieved with the newer formulations.¹³ The pharmacodynamic-pharmacokinetic (PK-PD) relationships of POS have been studied in vivo. A murine model of IA indicated that low-level POS-resistant isolates can be treated when the POS exposure is increased. Two in vivo studies demonstrated that POS retains efficacy against *A. fumigatus* isolates with POS-MIC of 0.5 mg/L as long

as POS exposure is sufficiently high.^{29,30} Based on these experiments, the required POS exposure (area under the concentration time curve (AUC)) in patients can be calculated for isolates with an increased MIC. The probability of target attainment for treatment of IA using standard dosing of POS tablet is estimated to be ~80% for isolates with POS-MIC of 0.25 mg/L and >90% for isolates with a POS-MIC of 0.125 mg/L.²⁸ The probability of target attainment for a POS-MIC of 0.5 mg/L was 24% and for a POS-MIC ≥ 0.5 mg/L it was 0%.

As determination of the AUC requires multiple sampling moments, and this AUC is linear correlated to Ctrough concentrations, quite often the Ctrough concentrations are used in daily practice as surrogate markers.^{13,28} Monte Carlo simulations estimated that the POS Ctrough concentrations needed to be 1.44–1.55 mg/L for isolates with a POS-MIC of 0.25 mg/L and 3.09–3.33 mg/L for isolates with a POS-MIC of 0.5 mg/L.²⁸

As the aforementioned in vivo experiments indicated that *A. fumigatus* with a POS-MIC of 0.5 mg/L can be treated with elevated POS dosing, we hypothesised that targeting high exposure with HD-POS is an oral step-down treatment option for azole-resistant IA. Although clinical evidence supporting HD-POS has not been described, preclinical animal studies and experience in veterinary medicine provided proof-of-principle for its efficiency.^{28,29}

4.2 | Mucormycosis

Limited in vivo models are available that assess POS for the treatment of mucormycosis. A neutropenic mouse model indicated similar pharmacodynamics for mucormycosis compared to *A. fumigatus* infections. An AUC/MIC of 87 was needed to treat *Rhizopus oryzae* infection, which was comparable to the target needed for IA (AUC/MIC of 76).³¹ Efficacy of POS showed a dose-response relationship in another in vivo model of experimental mucormycosis in which a dose of 100 mg/kg/day showed significant reduction of mortality of *Lichtheimia corymbifera* infection.³² Similar dose-response relationships were seen for *Mucor* species and *R. oryzae*.^{33,34} Compared to *A. fumigatus* isolates, the MICs of Mucorales are often higher with a geometric mean CLSI MIC of 0.39 mg/L³⁵ and an epidemiological cut-off value of 1 mg/L for *L. corymbifera*, *R. oryzae* and *R. microspores* and 4 mg/L for *M. circinelloides* (Figure 1).³⁶ Furthermore, the EUCAST MICs for Mucorales are higher than CLSI MICs for most species.³⁷ Taken into account, the similar target AUC/MIC for Mucorales as *A. fumigatus*, but higher MICs for Mucorales isolates compared to *A. fumigatus*, it seems reasonable to pursue higher than normal POS serum concentrations for the treatment of mucormycosis as long as this is not associated with toxicity.¹³

Posaconazole-oral suspension has been used as salvage therapy for mucormycosis with a success rate of approximately 60%–80%.³⁸ A recently published matched-paired analysis assessed the clinical effectiveness and safety of POS tablets and intravenous formulation in comparison with amphotericin B as first-line treatment and

with POS-OS as salvage treatment for invasive mucormycosis. POS tablets and intravenous formulation were effective in terms of treatment response and associated mortality. However, these observations should be interpreted with caution given the small sample size in this study.⁴³ Clinical data on PK/PD are lacking due to limited susceptibility data from clinical studies.³⁸

4.3 | Dosing and TDM

The pharmacokinetics of posaconazole tablets are best described by a one-compartment pharmacokinetic model with sequential zero-order and first-order absorption and a first-order disposition from the central compartment. Recently, several covariates were identified influencing bioavailability (like disease state, body weight, formulation), adsorption rate (food status) and clearance (dosing regimen) of POS tablets. Only body weight was considered clinically relevant.³⁹ Knowledge on the PK of POS helps to identify the optimal dose when targeting high exposure. Subsequently, an infrastructure is needed where one can quickly assess drug concentrations to deploy an adaptive approach in terms of dosing. With the new formulations of POS, a loading dose is given, which enables early assessment, typically by day 3, of POS concentrations. Follow-up samples are measured again before the 5th dose of every changed dosage.

The pharmacokinetics described above translates into an expected doubling of the Ctrough concentration when the dose of POS tablet or IV formulation is doubled. For example, when the Ctrough concentration is 1.5 mg/L, increasing the dose from 300 mg once daily to 300 mg twice daily can be expected to lead to a serum concentration of 3 mg/L. For safety reasons, we advise to increase the dose with no more than 200 mg per step.

4.4 | Inhibitory potential of HD-POS

Posaconazole is a strong CYP3A4 inhibitor, and the clinician should therefore also remain vigilant for drug interactions. In our case series, we had two patients with significant interactions. Toxicity of HD-POS in combination with vincristine was seen in a child with ALL, resulting in hepatotoxicity, convulsions and hypertension which might be attributed to the inhibition of CYP3A4 as well as P-gp resulting in increased levels of vincristine.⁴⁰ Another allogeneic stem cell transplant patient developed IA despite prophylaxis with voriconazole. Treatment with L-AmB was started but switched to HD-POS for progressive renal impairment. POS Ctrough concentration was 5.2 mg/L. After the patient was treated with panobinostat, a histone deacetylase (HDAC) inhibitor, grade 4 leukopenia developed. After 4 weeks of persisting grade 4 leukopenia, POS treatment was stopped as presumed culprit and leukopenia improved. This interaction could have been predicted based on the interaction of panobinostat with ketoconazole where panobinostat maximum serum concentrations were increased by an average of 1.6-fold.⁴¹

4.5 | Safety monitoring for HD-POS

We propose that the following safety measures are taken if HD-POS is used as a treatment strategy. At least the following laboratory tests should be performed twice weekly during the first 2 weeks and as long as the POS dosage is being increased: electrolytes, renal clearance, haemoglobin, leucocyte differentiation, thrombocytes and liver enzymes. Posaconazole may cause QT prolongation. Therefore, an ECG should be recorded before the start of HD-POS as well as during treatment. If no laboratory abnormalities possibly related to POS are observed the monitoring interval can be increased.

In conclusion, registration of new antifungals with efficacy against azole-resistant *A. fumigatus* is expected to take several more years. Therefore, targeting high serum concentrations of POS using the tablet or IV formulation is, in our point of view, a possible step-down option in patients with azole-resistant IA as long as the POS-MIC is <1 mg/L and for patients treated for mucormycosis with L-AmB. It should only be used when close monitoring for AE is implemented as described above in conjunction with TDM and when the benefits are likely to outweigh the risks.

ACKNOWLEDGMENTS

Part of this work was orally presented at the following conference: 28th European Congress of Clinical Microbiology and Infectious Diseases, Madrid Spain, 21–24 April 2018 (Title: Clinical experience with high-dose posaconazole for the treatment of azole-resistant invasive aspergillosis: O1059).

AUTHOR CONTRIBUTIONS

AFADS and JBB collected the clinical data. AFADS, JBB, RJB and BJAR analysed the data. AFADS, JBB, RJB and BJAR wrote the initial draft. All authors critically revised the initial draft and final manuscript.

ETHICAL APPROVAL

The authors confirm that the ethical policies of the journal, as noted on the journal's author guidelines page, have been adhered to. No ethical approval was required as this type of research does not fall under the Dutch law of research on human subjects.

TRANSPARENCY DECLARATIONS

AFADS has reports non-financial travel support from Abvie, Gilead, Amgen, Roche, and Pfizer to attend international conferences, outside the submitted work. P. E. V. reports grants from Gilead Sciences, grants from MSD, grants from Pfizer, grants from F2G, non-financial support from OLM, non-financial support from IMMY, outside the submitted work. RJB has served as a consultant to and has received unrestricted and research grants from Astellas Pharma, Inc, F2G, Gilead Sciences, Merck Sharpe and Dohme Corp., and Pfizer, Inc All contracts were through Radboudumc, and payments were invoiced by Radboudumc. None of the work is related to this manuscript. All other authors: none to declare.

ORCID

Jochem B. Buil  <https://orcid.org/0000-0003-4031-0778>

REFERENCES

- Maertens JA, Raad II, Marr KA, et al. Isavuconazole versus voriconazole for primary treatment of invasive mould disease caused by *Aspergillus* and other filamentous fungi (SECURE): a phase 3, randomised-controlled, non-inferiority trial. *Lancet*. 2016;387:760–769.
- Slobbe L, Polinder S, Doorduyn JK, et al. Outcome and medical costs of patients with invasive aspergillosis and acute myelogenous leukemia-myelodysplastic syndrome treated with intensive chemotherapy: an observational study. *Clin Infect Dis*. 2008;47:1507–1512.
- Perlin DS, Rautemaa-Richardson R, Alastruey-Izquierdo A. The global problem of antifungal resistance: prevalence, mechanisms, and management. *Lancet Infect Dis*. 2017;17:e383–e392.
- Chong GM, van der Beek MT, von dem Borne PA, et al. PCR-based detection of *Aspergillus fumigatus* Cyp51A mutations on bronchoalveolar lavage: a multicentre validation of the AsperGenius assay^(R) in 201 patients with haematological disease suspected for invasive aspergillosis. *J Antimicrob Chemother*. 2016;71:3528–3535.
- van der Linden JW, Snelders E, Kampinga GA, et al. Clinical implications of azole resistance in *Aspergillus fumigatus*, The Netherlands, 2007–2009. *Emerg Infect Dis*. 2011;17:1846–1854.
- Verweij PE, Ananda-Rajah M, Andes D, et al. International expert opinion on the management of infection caused by azole-resistant *Aspergillus fumigatus*. *Drug Resist Updat*. 2015;22:30–40.
- Patterson TF, Thompson GR 3rd, Denning DW, et al. Practice guidelines for the diagnosis and management of aspergillosis: 2016 Update by the Infectious Diseases Society of America. *Clin Infect Dis*. 2016;63:e1–e60.
- Cornely OA, Maertens J, Winston DJ, et al. Posaconazole vs. fluconazole or itraconazole prophylaxis in patients with neutropenia. *N Engl J Med*. 2007;356:348–359.
- Ullmann AJ, Lipton JH, Vesole DH, et al. Posaconazole or fluconazole for prophylaxis in severe graft-versus-host disease. *N Engl J Med*. 2007;356:335–347.
- Krishna G, Ma L, Martinho M, Preston RA, O'Mara E. A new solid oral tablet formulation of posaconazole: a randomized clinical trial to investigate rising single- and multiple-dose pharmacokinetics and safety in healthy volunteers. *J Antimicrob Chemother*. 2012;67:2725–2730.
- Liebenstein TK, Widmer KM, Fallon MJ. Retrospective analysis of goal drug level attainment of posaconazole for invasive fungal infection prophylaxis in patients with acute myeloid leukemia pre- and post-switch to tablet formulation. *J Oncol Pharm Pract*. 2018;24(8):599–603.
- Jang SH, Colangelo PM, Gobburu JV. Exposure-response of posaconazole used for prophylaxis against invasive fungal infections: evaluating the need to adjust doses based on drug concentrations in plasma. *Clin Pharmacol Ther*. 2010;88:115–119.
- Dekkers BGJ, Bakker M, van der Elst KCM, et al. Therapeutic drug monitoring of posaconazole: an update. *Curr Fungal Infect Rep*. 2016;10:51–61.
- Stott KE, Hope WW. Therapeutic drug monitoring for invasive mould infections and disease: pharmacokinetic and pharmacodynamic considerations. *J Antimicrob Chemother*. 2017;72:i12–i18.
- Ashbee HR, Barnes RA, Johnson EM, Richardson MD, Gorton R, Hope WW. Therapeutic drug monitoring (TDM) of antifungal agents: guidelines from the British Society for Medical Mycology. *J Antimicrob Chemother*. 2014;69:1162–1176.
- Naranjo CA, Busto U, Sellers EM, et al. A method for estimating the probability of adverse drug reactions. *Clin Pharmacol Ther*. 1981;30:239–245.

17. European Medicines Agency, Committee for Medicinal Products for Human Use. Assessment report Noxafil Procedure No. EMEA/H/C/000610/X/0028. http://www.ema.europa.eu/docs/en_GB/document_library/EPAR_-_Assessment_Report_-_Variation/human/000610/WC500168187.pdf. Accessed June 28, 2018.
18. Kuriakose K, Nesbitt WJ, Greene M, Harris B. Posaconazole-induced pseudohyperaldosteronism. *Antimicrob Agents Chemother.* 2018;62:e02130-17.
19. Barton K, Davis TK, Marshall B, Elward A, White NH. Posaconazole-induced hypertension and hypokalemia due to inhibition of the 11 β -hydroxylase enzyme. *Clin Kidney J.* 2018;11:691-693.
20. Boughton C, Taylor D, Ghataore L, Taylor N, Whitelaw BC. Mineralocorticoid hypertension and hypokalaemia induced by posaconazole. *Endocrinol Diabetes Metab Case Rep.* 2018;2018:17-0157.
21. Cornely OA, Duarte RF, Haider S, et al. Phase 3 pharmacokinetics and safety study of a posaconazole tablet formulation in patients at risk for invasive fungal disease. *J Antimicrob Chemother.* 2016;71:718-726.
22. Cornely OA, Robertson MN, Haider S, et al. Pharmacokinetics and safety results from the Phase 3 randomized, open-label, study of intravenous posaconazole in patients at risk of invasive fungal disease. *J Antimicrob Chemother.* 2017;72:3406-3413.
23. Belling M, Kanate AS, Shillingsburg A, et al. Evaluation of serum posaconazole concentrations in patients with hematological malignancies receiving posaconazole suspension compared to the delayed-release tablet formulation. *Leuk Res Treatment.* 2017;2017:1-6.
24. Tverdek FP, Heo ST, Aitken SL, et al. Real-life assessment of the safety and effectiveness of the new tablet and intravenous formulations of posaconazole in the prophylaxis of invasive fungal infections via analysis of 343 courses. *Antimicrob Agents Chemother.* 2017;61(8):e00188-17.
25. de Greeff SC, NethMap MJW. Consumption of antimicrobial agents and antimicrobial resistance among medically important bacteria in the Netherlands/MARAN 2017: Monitoring of antimicrobial resistance and antibiotic usage in animals in the Netherlands in 2016; 2017. Rijksinstituut voor Volksgezondheid en Milieu RIVM.
26. Snelders E, Camps SM, Karawajczyk A, et al. Triazole fungicides can induce cross-resistance to medical triazoles in *Aspergillus fumigatus*. *PLoS ONE.* 2012;7:e31801.
27. Buil JB, Bruggemann RJM, Wasmann RE, et al. Isavuconazole susceptibility of clinical *Aspergillus fumigatus* isolates and feasibility of isavuconazole dose escalation to treat isolates with elevated MICs. *J Antimicrob Chemother.* 2018;73:134-142.
28. Seyedmousavi S, Mouton JW, Melchers WJ, Bruggemann RJ, Verweij PE. The role of azoles in the management of azole-resistant aspergillosis: from the bench to the bedside. *Drug Resist Updat.* 2014;17:37-50.
29. Mavridou E, Bruggemann RJ, Melchers WJ, Mouton JW, Verweij PE. Efficacy of posaconazole against three clinical *Aspergillus fumigatus* isolates with mutations in the *cyp51A* gene. *Antimicrob Agents Chemother.* 2010;54:860-865.
30. Lepak AJ, Marchillo K, Vanhecker J, Andes DR. Posaconazole pharmacodynamic target determination against wild-type and Cyp51 mutant isolates of *Aspergillus fumigatus* in an *in vivo* model of invasive pulmonary aspergillosis. *Antimicrob Agents Chemother.* 2013;57:579-585.
31. Lewis RE, Albert ND, Kontoyiannis DP. Comparative pharmacodynamics of posaconazole in neutropenic murine models of invasive pulmonary aspergillosis and mucormycosis. *Antimicrob Agents Chemother.* 2014;58:6767-6772.
32. Dannaoui E, Meis JF, Loebenberg D, Verweij PE. Activity of posaconazole in treatment of experimental disseminated zygomycosis. *Antimicrob Agents Chemother.* 2003;47:3647-3650.
33. Sun QN, Najvar LK, Bocanegra R, Loebenberg D, Graybill JR. In vivo activity of posaconazole against *Mucor spp.* in an immunosuppressed-mouse model. *Antimicrob Agents Chemother.* 2002;46:2310-2312.
34. Spreghini E, Orlando F, Giannini D, Barchiesi F. In vitro and in vivo activities of posaconazole against zygomycetes with various degrees of susceptibility. *J Antimicrob Chemother.* 2010;65:2158-2163.
35. Chowdhary A, Kathuria S, Singh PK, et al. Molecular characterization and in vitro antifungal susceptibility of 80 clinical isolates of mucormycetes in Delhi, India. *Mycoses.* 2014;57(Suppl 3):97-107.
36. Espinel-Ingroff A, Chakrabarti A, Chowdhary A, et al. Multicenter evaluation of MIC distributions for epidemiologic cutoff value definition to detect amphotericin B, posaconazole, and itraconazole resistance among the most clinically relevant species of Mucorales. *Antimicrob Agents Chemother.* 2015;59:1745-1750.
37. Chowdhary A, Singh PK, Kathuria S, Hagen F, Meis JF. Comparison of the EUCAST and CLSI broth microdilution methods for testing isavuconazole, posaconazole, and amphotericin B against molecularly identified Mucorales species. *Antimicrob Agents Chemother.* 2015;59:7882-7887.
38. Greenberg RN, Mullane K, van Burik JA, et al. Posaconazole as salvage therapy for zygomycosis. *Antimicrob Agents Chemother.* 2006;50:126-133.
39. van Iersel M, Rossenu S, de Greeff R, Waskin H. A population pharmacokinetic model for a solid oral tablet formulation of posaconazole. *Antimicrob Agents Chemother.* 2018;62(7):e02465-17.
40. Pekpak E, Ileri T, Ince E, Ertem M, Uysal Z. Toxicity of vincristine combined with posaconazole in children with acute lymphoblastic leukemia. *J Pediatr Hematol Oncol.* 2018;40(5):e309-e310.
41. Hamberg P, Woo MM, Chen LC, et al. Effect of ketoconazole-mediated CYP3A4 inhibition on clinical pharmacokinetics of panobinostat (LBH589), an orally active histone deacetylase inhibitor. *Cancer Chemother Pharmacol.* 2011;68:805-813.
42. De Pauw B, Walsh TJ, Donnelly JP, et al. Revised definitions of invasive fungal disease from the European Organization for Research and Treatment of Cancer/Invasive Fungal Infections Cooperative Group and the National Institute of Allergy and Infectious Diseases Mycoses Study Group (EORTC/MSG) Consensus Group. *Clin Infect Dis.* 2008;46(12):1813-1821.
43. Salmanton-García J, Seidel D, Koehler P, et al. Matched-paired analysis of patients treated for invasive mucormycosis: standard treatment versus posaconazole new formulations (MoveOn). *J Antimicrob Chemother.* 2019;74(11):3315-3327.

How to cite this article: Schauwvlieghe AFAD, Buil JB, Verweij PE, et al. High-dose posaconazole for azole-resistant aspergillosis and other difficult-to-treat mould infections. *Mycoses.* 2020;63:122-130. <https://doi.org/10.1111/myc.13028>