

Collaborative International Research in Clinical and Longitudinal Experience Study in NMOSD

Lawrence J. Cook, PhD, John W. Rose, MD, Jessica S. Alvey, MS, Anna Marie Jolley, BS, Renee Kuhn, BS, Brie Marron, BS, Melissa Pederson, MEd, Rene Enriquez, BS, Jeff Yearley, BA, Stephen McKechnie, MS, May H. Han, MD, Anna J. Tomczak, MS, Michael Levy, MD, PhD, Maureen A. Mealy, MScN, Jessica Coleman, BA, Jeffrey L. Bennett, MD, PhD, Ruth Johnson, BS, Myka Barnes-Garcia, BA, Anthony L. Traboulsee, MD, Robert L. Carruthers, MD, Lisa Eunyong Lee, MSc, Julia J. Schubert, BS, Katrina McMullen, PhD, Ilya Kister, MD, Zoe Rimler, BS, Allyson Reid, BA, Nancy L. Sicotte, MD, Sarah M. Planchon, PhD, Jeffrey A. Cohen, MD, Diane Ivancic, CCRP, Jennifer L. Sedlak, BSN, Ilana Katz Sand, MD, Pavle Repovic, MD, PhD, Lilyana Amezcua, MD, MS, Ana Pruitt, BS, Erika Amundson, BA, Tanuja Chitnis, MD, Devin S. Mullin, BS, Eric C. Klawiter, MD, Andrew W. Russo, BS, Claire S. Riley, MD, Kaho B. Onomichi, MS, Libby Levine, RN, ANP-BC, Katherine E. Nelson, BA, Nancy M. Nealon, MD, Casey Engel, BA, Mason Kruse-Hoyer, MD, MA, Melanie Marcille, BA, Leticia Tornes, MD, Anne Rumpf, BS, Angela Greer, BS, Megan Kenneally Behne, AS, Renee R. Rodriguez, AS, Daniel W. Behne, MArch, Derek W. Blackway, BA, Brian Coords, MA, Terrence F. Blaschke, MD, Judy Sheard, MPH, MA, Terry J. Smith, MD, Jacinta M. Behne, MA, and Michael R. Yeaman, PhD, on behalf of The Guthy-Jackson Charitable Foundation International Clinical Consortium (GJCF-ICC)

Correspondence

Dr. Cook
larry.cook@hsc.utah.edu

Neurol Neuroimmunol Neuroinflamm 2019;6:e583. doi:10.1212/NXI.0000000000000583

Abstract

Objective

To develop a resource of systematically collected, longitudinal clinical data and biospecimens for assisting in the investigation into neuromyelitis optica spectrum disorder (NMOSD) epidemiology, pathogenesis, and treatment.

Methods

To illustrate its research-enabling purpose, epidemiologic patterns and disease phenotypes were assessed among enrolled subjects, including age at disease onset, annualized relapse rate (ARR), and time between the first and second attacks.

Results

As of December 2017, the Collaborative International Research in Clinical and Longitudinal Experience Study (CIRCLES) had enrolled more than 1,000 participants, of whom 77.5% of the NMOSD cases and 71.7% of the controls continue in active follow-up. Consanguineous relatives of patients with NMOSD represented 43.6% of the control cohort. Of the 599 active cases with complete data, 84% were female, and 76% were anti-AQP4 seropositive. The majority were white/Caucasian (52.6%), whereas blacks/African Americans accounted for 23.5%, Hispanics/Latinos 12.4%, and Asians accounted for 9.0%. The median age at disease onset was 38.4 years, with a median ARR of 0.5. Seropositive cases were older at disease onset, more likely to be black/African American or Hispanic/Latino, and more likely to be female.

Conclusions

Collectively, the CIRCLES experience to date demonstrates this study to be a useful and readily accessible resource to facilitate accelerating solutions for patients with NMOSD.

From the University of Utah School of Medicine (L.J.C., J.W.R., J.S.A., A.M.J., R.K., B.M., M.P., R.E., J.Y., S.M.), Salt Lake City; Division of Neuroimmunology and Multiple Sclerosis Center (M.H.H.), Department of Neurology and Neurological Sciences, Stanford University; Department of Neurology and Neurological Sciences (A.J.T.), Stanford School of Medicine, CA; Department of Neurology, Johns Hopkins University School of Medicine (M.L., M.A.M., J.C.), Baltimore, MD; Departments of Neurology and Ophthalmology (J.L.B., R.J., M.B.-G.), University of Colorado School of Medicine, Aurora; Department of Medicine & Neurology (A.L.T., R.L.C., L.E.L., J.J.S., K.M.), University of British Columbia, Vancouver, Canada; NYU Langone Health (I.K., Z.R., A.R.), New York; Department of Neurology, Cedars-Sinai Medical Center (N.L.S.), Los Angeles, CA; Mellen Center for MS Treatment and Research (S.M.P., J.A.C., D.I., J.L.S.), Neurological Institute, Cleveland Clinic, OH; Icahn School of Medicine at Mount Sinai (I.K.S.), New York; Multiple Sclerosis Center (P.R.), Swedish Neuroscience Institute, Seattle, WA; Department of Neurology (L.A., A.P., E.A.), Keck School of Medicine, University of Southern California, Los Angeles; Department of Neurology (T.C., D.S.M.), Brigham and Women's Hospital, Harvard Medical School, Boston, MA; Department of Neurology (E.C.K., A.W.R.), Massachusetts General Hospital, Harvard Medical School, Boston; Department of Neurology (C.S.R., K.B.O., L.L., K.E.N.), Columbia University Medical Center; Weill Cornell Medicine (M.M.N., C.E., M.K.-H., M.M.), New York; Department of Neurology (L.T.), Division of Multiple Sclerosis, University of Miami Miller School of Medicine, FL; PPD (A.R., A.G.), Wilmington, NC; The Guthy-Jackson Charitable Foundation (M.K.B., R.R.R., D.W. Behne., D.W. Blackway, B.C., J.S., J.M.B.), Beverly Hills; Departments of Medicine and of Molecular Pharmacology (T.F.B.), Stanford University School of Medicine, CA; Kellogg Eye Center (T.J.S.), University of Michigan Medical School, Ann Arbor; Department of Medicine, University of California, Los Angeles (M.R.Y.); and Harbor-UCLA Medical Center/LABioMed (M.R.Y.), Torrance, CA.

Presented in part at the European Committee for Treatment and Research in Multiple Sclerosis (ECTRIMS), London, England, August 2016, Abstract Nos. 874 and 1615.

Go to Neurology.org/NN for full disclosures. Funding information is provided at the end of the article.

The Article Processing Charge was funded by The Guthy-Jackson Charitable Foundation.

GJCF-ICC Coinvestigators are listed in Appendix 2.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 (CC BY-NC-ND), which permits downloading and sharing the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

Glossary

AQP4 = aquaporin-4; **ARR** = annual relapse rate; **CRC** = Clinical Research Coordinator; **EDSS** = Expanded Disability Status Scale; **GJCF** = The Guthy-Jackson Charitable Foundation; **ICC** = International Clinical Consortium; **IQR** = interquartile range; **MOG** = myelin oligodendrocyte glycoprotein; **NMOSD** = neuromyelitis optica spectrum disorder; **PSF** = patient study file.

Neuromyelitis optica spectrum disorder (NMOSD) represents a chronic, potentially debilitating and life-threatening neuroinflammatory process primarily targeting the optic nerves, spinal cord, and brain.^{1–4} The typical clinical course of NMOSD is marked by multiple relapses resulting in cumulative neurologic disabilities. These events are interspersed with remissions from disease activity of variable duration. Heightened awareness of the disease among health care providers and the public and recent advances in diagnostic precision have increased estimates of worldwide NMOSD prevalence, reaching as high as 10 per 100,000 in some populations.^{5–8} This projection translates to more than 15,000 patients with NMOSD in the United States, suggesting that hundreds of thousands of cases exist worldwide. NMOSD disproportionately affects females (up to 7:1 female-to-male ratio), with anti-aquaporin-4 (AQP4) antibody (hereafter referred to as anti-AQP4) positive disease having even greater propensity for women.^{9,10} Yet, many details regarding etiology, pathogenesis, risk factors, and demography of NMOSD are in need of greater understanding.

Although case series and observational studies suggest benefit from immunotherapy, to date no treatment of NMOSD has been proven safe and effective in prospective, double-masked and adequately powered clinical trials.^{11–13} Because of its rarity, insufficient access to well-characterized patient cohorts has historically hindered studies as has an absence of high-fidelity preclinical models of human disease. Limitations in carefully standardized, longitudinal clinical research tools have also impeded investigation of NMOSD immunopathogenesis. However, 4 separate clinical trials have now reported positive results evaluating 3 compounds (eculizumab, satralizumab, and inebilizumab) in studies assessing efficacy in delaying or preventing relapses in NMOSD.

The Guthy-Jackson Charitable Foundation (GJCF) initiated an observational study of NMOSD in which patients and comparative controls are enrolled and evaluated longitudinally in a standardized manner. This study, known as Collaborative International Research in Clinical and Longitudinal Experience Studies (CIRCLES) for NMOSD, was launched in November 2013. In CIRCLES, participant clinical data, demographic profiles, and biospecimens were collected at geographically dispersed academic medical centers located throughout North America (figure 1).

The design and performance of the CIRCLES study are described here, along with initial data analysis illustrating

the utility of its database and biorepository to advance scientific knowledge and clinical care in NMOSD. It is anticipated that this study will accelerate greater understanding of NMOSD and in turn the development of safe and effective therapies to benefit patients with NMOSD and perhaps those diagnosed with other autoimmune diseases.

Methods

Clinical research standards

Human subjects protection

Participant enrollment is conducted in accordance with the guidelines specified by the Office of Human Research Protections of the US Food and Drug Administration. A standardized protocol, manual of operations, patient study file (PSF), and informed consent or assent documents were approved by the institutional review board of each participating institution. Written and verbal consent or assent was obtained before beginning study procedures. The protocol and PSF were updated periodically.

Study goals and design

Design

CIRCLES is a prospective, multicenter, cross-sectional, and longitudinal study enabling comparisons of NMOSD cases and controls from which clinical data and biospecimens were collected using standardized methods. These were collected from cases at 6-month intervals and at least annually from control participants. When possible, clinical data and biospecimens were obtained from cases during or within 10 days following clinically confirmed relapses.


Goals

Two primary goals of CIRCLES include the following: (1) establish a cohort of patients with NMOSD and comparative controls who are longitudinally assessed at standardized intervals and (2) analyze acquired clinical data and biospecimens, thus improving knowledge of NMOSD and the patient experience.

Sites

Multiple study sites were established at academic institutions throughout North America (figure 1), each led by a clinical investigator/neurologist with expertise in NMOSD. Study sites were selected based on the size/activity of their NMOSD

Figure 1 Geographic location of CIRCLES clinical sites



patient cohort and capability to collect data and biospecimens in the protocol-defined manner. Biospecimens, predominantly peripheral blood constituents, are rapidly transferred to a centralized commercial laboratory for processing and archiving.

Cohorts

The study comprises 3 participant cohorts based on the following inclusion criteria: (1) cases with clinically diagnosed NMOSD according to either the Wingerchuk 2006¹⁴ or International Panel for NMOSD Diagnosis 2015¹⁴ criteria and classified with respect to anti-AQP4 serostatus; (2) comparative disease controls (including CNS autoimmune diseases [e.g., MS]; other autoimmune diseases [e.g., systemic lupus erythematosus, Sjögren syndrome, and type I diabetes mellitus]; chronic non-autoimmune inflammatory or systemic conditions [e.g., cardiovascular disease and type 2 diabetes]); and (3) healthy controls (i.e., those not carrying a chronic disease diagnosis at enrollment). Controls included consanguineous relatives and unrelated individuals (tables 1 and 2). Enrollment is targeted to a 2:1 ratio of cases to controls and is monitored centrally. Individuals (both cases and controls) are excluded if the treating physician feels that they are not appropriate for the study. Control participants are not sex or age matched. Some comparative disease controls are recruited from referral cohorts (e.g., MS). Others are recruited through opportunity or convenience.

Sites are instructed to enroll control participants at a rate of 50% MS and 50% from the other categories.

Intervals

After enrollment, NMOSD cases are evaluated clinically at 6-month intervals to provide an updated clinical history and complete set of biospecimens. Control participants undergo these same assessments at least annually. The panel of biospecimens routinely collected is listed in supplemental table 1 (links.lww.com/NXI/A121). The protocol allows collection of CSF and additional tissues (e.g., placenta) as available from medically indicated care. Relapses are evaluated regardless of interval and adjudicated by site neurologists.

Participating cohorts

Eligibility

Individuals fulfilling inclusion criteria and absent exclusion criteria are eligible for enrollment. Individuals weighing <17 kg are excluded from blood collections but may otherwise participate.

Enrollment

Clinical research coordinators (CRCs) screen information pertaining to inclusion and exclusion criteria. Individuals receive study information through mail and/or social media and, where institutional review board-approved, have the

Table 1 Summary of case participant characteristics by serostatus

	Overall ^a (N = 599)	NMO-IgG status		p Value
		Negative (N = 139)	Positive (N = 449)	
Female	504 (84.1%)	102 (73.4%)	393 (87.5%)	<0.001 ^g
Participant primary ethnicity/race designation				<0.001 ^g
Asian	54 (9.0%)	17 (12.2%)	37 (8.2%)	
Black or African American	141 (23.5%)	16 (11.5%)	125 (27.8%)	
Hispanic or Latino	74 (12.4%)	13 (9.4%)	60 (13.4%)	
White	315 (52.6%)	87 (62.6%)	218 (48.6%)	
Other	15 (2.5%)	6 (4.3%)	9 (2.0%)	
Not reported	0 (0%)	0 (0%)	0 (0%)	
Age at consent	47.1 (36.0–57.2)	43.3 (29.8–52.3)	48.6 (37.7–59.1)	<0.001 ^h
Age at first episode onset^b	38.4 (28.9–50.6)	35.1 (25.8–46.4)	39.4 (29.9–52.5)	0.002 ^h
Relapse/year from disease onset to most recent visit^c	0.5 (0.3–0.8)	0.5 (0.3–0.9)	0.4 (0.3–0.8)	0.031 ^h
Time (y) from first episode onset to enrollment^b	4.6 (1.5–10.1)	3.8 (1.4–7.0)	4.9 (1.5–11.0)	0.008 ^h
Longitudinally extensive transverse myelitis^d	396 (66.1%)	83 (59.7%)	305 (67.9%)	0.064 ^g
Optic neuritis^f	395 (65.9%)	102 (73.4%)	286 (63.7%)	0.038 ^g
Brainstem syndrome^f	155 (25.9%)	40 (28.8%)	111 (24.7%)	0.346 ^g
Focal transverse myelitis^e	143 (23.9%)	46 (33.1%)	94 (20.9%)	0.004 ^g
Area postrema syndrome^f	88 (14.7%)	20 (14.4%)	66 (14.7%)	0.920 ^g
Cerebral syndrome^f	74 (12.4%)	20 (14.4%)	53 (11.8%)	0.425 ^g
Diencephalic syndrome^f	23 (3.8%)	7 (5.0%)	15 (3.3%)	0.360 ^g

Abbreviation: NMO = neuromyelitis optica.

^a Eleven case participants have undetermined serostatus.

^b Age at first episode onset and time from first episode onset to enrollment not recorded on 6 participants.

^c Relapses per year from disease onset to most recent visit not recorded on 32 participants because of insufficient follow-up or missing data.

^d Longitudinally extensive transverse myelitis not recorded on 3 participants.

^e Focal transverse myelitis not recorded on 4 participants.

^f Optic neuritis, brainstem syndrome, area postrema syndrome, cerebral syndrome, and diencephalic syndrome not recorded on 2 participants.

^g χ^2 test of association.

^h Wilcoxon rank-sum test.

option to provide preliminary information telephonically with consent obtained in advance of study participation. At enrollment, a thorough review of medical records and clinical examination is performed by the study neurologist. A complete disease history and additional relevant study data are collected during the initial interview (table e-3, links.lww.com/NXI/A121). The majority of participants to date have been enrolled coinciding with medically indicated appointments; however, in some cases, enrollment was conducted per protocol at patient-oriented educational events.

Clinical database

Clinical metadata

The CIRCLES PSF (table e-2, links.lww.com/NXI/A121) is completed for each participant at enrollment and updated at each follow-up study visit. The PSF data include

demographics, disease phenotype, treatment history, and other relevant characteristics. Other than NMOSD disease history, identical clinical data are collected as appropriate from case and healthy control participants.

Data security

Data are entered into a web-accessible and password-secured electronic data capture system. The CIRCLES study incorporates a query management system that executes nightly. An email notification is generated for each site's CRC identifying any new discrepant data. A weekly reminder email is also provided for remaining discrepant data. The system tracks queries from generation to resolution. Data are curated for quality, consistency, and completeness by the Data Coordinating Center biostatistics group before archiving. Study data are backed up hourly, and a full snapshot of the study is archived nightly.

Table 2 Summary of control participant characteristics by relatedness to NMOSD cases

	Case blood relative (N = 123)	Unrelated (N = 95)	Total (N = 218)
Female	73 (59.3%)	71 (74.7%)	144 (66.1%)
Race^a			
Asian	9 (7.3%)	8 (8.4%)	17 (7.8%)
Black or African American	8 (6.5%)	13 (13.7%)	21 (9.6%)
Hispanic or Latino	20 (16.3%)	16 (16.8%)	36 (16.5%)
White	81 (65.9%)	56 (58.9%)	137 (62.8%)
Other	4 (3.3%)	2 (2.1%)	6 (2.8%)
Age at consent (y)	44.9 (34.2–53.9)	50.9 (38.9–58.3)	47.1 (35.9–56.2)
Comparative disease			
MS	37 (30.1%)	0 (0.0%)	37 (17.0%)
CNS autoimmune disease other than MS	10 (8.1%)	0 (0.0%)	10 (4.6%)
Systemic autoimmune disease	5 (4.1%)	14 (14.7%)	19 (8.7%)
CNS disorder unrelated to an inflammatory disease	4 (3.3%)	1 (1.1%)	5 (2.3%)
Systemic chronic condition	3 (2.4%)	4 (4.2%)	7 (3.2%)
None of the above	69 (56.1%)	78 (82.1%)	147 (67.4%)

Abbreviation: NMOSD = neuromyelitis optica spectrum disorder.

^a Race not recorded on 1 participant.

Biospecimen repository

Collection, processing, and storage

Biospecimens are collected according to standard operating procedures at enrollment and follow-up visits. A panel of blood specimens is obtained (table e-1, links.lww.com/NXI/A121) by routine venipuncture by a certified phlebotomist at each scheduled clinical visit. Biospecimens are transported by express courier to a commercial laboratory for processing, systematic labeling, and archiving within 24 hours of collection under certified storage conditions (liquid nitrogen for peripheral blood mononuclear cells and –80°C for sera, plasma, RNA, and DNA).

Quality and serostatus

Biospecimens are routinely assessed for quality postprocessing and before cryopreservation. Autoantibody serostatus is determined by the respective study site based on reference laboratory assay or review of the case record. For analytical purposes, seropositivity is defined as having detected anti-AQP4 at any point during the participant's history.

Statistical analysis

Analytical range

The current report encompasses data sets obtained from 2013 through 2017. The CIRCLES study is ongoing.

Analytical methods

Descriptive statistics (medians or interquartile ranges [IQRs] for numeric variables; counts and percentages for categorical

variables) were used to analyze data. Wilcoxon rank-sum tests were used to assess relationships between age at disease onset, annualized relapse rate (ARR) during enrollment in the study, and time between the first and second attacks in relation to other demographic characteristics in NMOSD cases. Relationships between serostatus and race, serostatus and sex, and race and sex were examined using χ^2 tests. All analyses were performed in SAS 9.4 (Cary, NC).

Data availability

Access to data and biospecimens is provided to qualified scholars in a peer-reviewed process. Applications are adjudicated by a biorepository oversight committee elected from among the members of the GJCF International Clinical Consortium (ICC).

Results

Study enrollment

As of December 2017, CIRCLES had enrolled 849 NMOSD cases and 339 controls, of which 658 (77.5%) and 243 (71.7%), respectively, continue to participate. The percentage of enrollees remaining active has increased over time. Of the 161 participants enrolled in 2013, 49.7% are still active. This compares to 73.7% of those enrolled in 2014, 67.9% in 2015, 74.4% in 2016, and 90.8% in 2017. Inability to contact accounted for most inactivity (74%), followed by withdraw of consent (14%), no longer able to participate (9%), and death (2%). Of all participants, 60.3%, 31.3%, and 8.4% were enrolled

at study sites, national patient day events, and regional patient day events, respectively. In-clinic enrollees, compared with national and regional patient day enrollees, were more likely to remain active (89.7% vs 55.1% and 74.5%, respectively). The size of the enrolled cohort varied among study sites, with the largest site enrolling 286 (24.1%) of all participants and the smallest enrolling 20 (1.7%). Among active NMOSD cases, 495 (75.2%) have undergone one or more follow-up visits with biospecimen collection. Among active controls, 129 (53.1%) had one or more follow-up visits and biospecimen collection. The remainder of the analyses presented are based on these active cases and control participants who have complete data as of the end of the study period.

Cohort demographics and epidemiology

In the CIRCLES cohort, the female-to-male ratio was 5.3:1 among cases (tables 1 and 2). NMOSD cases self-identified as white/Caucasian (52.6%), 23.5% black/African American, 12.4% Hispanic/Latino, 9.0% Asian, and 2.5% from all other races/ethnicities. The overall median age at NMOSD onset was 38.4 years (IQR 28.9–50.6 years), appears to be normally distributed, and spans the range from 2.7 to 79.9 years (figure 2). The median time between disease onset and time to study enrollment was 4.6 years (IQR 1.5–10.1 years). CIRCLES cases experienced an ARR of 0.5 (IQR 0.3–0.9). One hundred thirty-two cases appear to be monophasic. For cases with ≥ 2 clinically documented NMOSD relapses ($n = 207$), the median time between the first and second attacks was 0.8 years (9 months; IQR 3.3–27.9). The most common relapse manifestations were longitudinally extensive transverse myelitis (≥ 2 vertebral segments; 396, 66.1%) and optic neuritis (395, 65.9%). Brainstem syndromes were identified in 155 (25.9%) cases, 143 (23.9%) manifested focally confined transverse myelitis, 88 (14.7%) area postrema syndrome (e.g., prolonged or intractable nausea/vomiting or hiccups), 74 (12.4%) cerebral syndrome (cognitive

and/or sensory impairment, pain, bowel and/or bladder dysfunction, or limb weakness), and 23 (3.8%) diencephalic and/or brainstem syndromes (facial numbness, hearing loss, dysphagia, or dysarthria) (table 1).

Biospecimen repertoire

Of the active NMOSD cases, 193 (32.2%) provided a single set of biospecimens; 2 longitudinal sets were collected from 151 (25.2%) cases, 3 sets from 107 (17.9%), and 4 or more from 131 (21.9%). Of the active controls, 102 (46.8%) provided a single biospecimen set, 65 (29.8%) 2 sets, 26 (11.9%) 3 sets, and 21 individuals (9.6%) have provided 4 or more longitudinal samples.

Correlation analyses

Bioinformatic analyses have revealed several significant correlates in the CIRCLES cohort (table 1). Anti-AQP4 seropositive cases were more likely to be female (87.5%) compared with seronegative cases (73.4%, $p < 0.001$). Significant differences in racial distribution by serostatus ($p < 0.001$) were also detected. This result was driven largely by differences in the black/African American and white race categories. Although black/African American participants accounted for only 11.5% of the seronegative population, they comprise 27.8% of seropositive cases. Similarly, Hispanic/Latino cases represent only 9.4% of seronegative cases, but 13.4% of seropositive cases. Conversely, white/Caucasian cases account for 62.6% of seronegative cases, but less than half (48.6%) of the seropositive cases. Similarly, Asians account for a higher percentage of the seronegative population (12.2%) compared with the seropositive population (8.2%).


Overall, seropositive cases are older at initial NMOSD attack compared with seronegative participants (39.4 vs 35.1; $p = 0.002$; figure 2). White/Caucasian and Asian cases tend to be older at first attack compared with black/African Americans or Hispanics (41.1 and 38.3 vs 36.5 and 36.0, $p < 0.001$; table 3). Seropositive cases tend to have lower ARR than those who are seronegative (0.46 vs 0.55; $p = 0.030$). No significant differences were detected between race and ARR ($p = 0.34$).

Of the 218 control participants, 95 (43.6%) are consanguineous with an enrolled case. One hundred forty-four (66.1%) are female. The median age of controls at enrollment was 47.1 years. Of the related controls, none have MS, 17.9% have another autoimmune disease, and 82.1% have no comparative disease or chronic condition. Of the unrelated controls, 28.5% have MS, 15.4% have another autoimmune disease, and 56.1% have no comparative disease or condition (table 2).

Discussion

The CIRCLES study represents a unique and multicenter longitudinal observational study, which has successfully recruited and retained a large number of patients affected by the rare disease NMOSD. The substantial number of control participants who are consanguineous with enrolled cases also

Figure 2 Age at onset by serostatus and gender



provides important new opportunities to understand disease resilience. In this respect, individuals with familial genotypes and environmental exposures, but who do not manifest NMOSD, can be evaluated in relation to patients with NMOSD. In addition, the CIRCLES project reflects the collaborative input of the GJCF-ICC, a global network of scientific and medical experts in NMOSD.

The demographic characteristics of the CIRCLES cohort to date are comparable to those of previously described NMOSD registries.^{13,15–22} Interesting relationships have emerged from initial demographic, epidemiologic, and correlational analyses of this cohort. Key relationships include identification of correlations between disease attributes and sex, age, and race. NMOSD cases are predominantly female (5.3:1) and anti-AQP4 seropositive (76.4%). Disease onset most commonly occurs in the fourth decade of life. These findings are congruent with recent epidemiologic studies of NMOSD regarding sex predominance, age at onset, and disease clustering in individuals and their first-degree relatives.^{5,23–25}

Analysis of the CIRCLES cohort supports the concept that a sizable proportion of cases satisfying either 2006 or 2015 diagnostic criteria for NMOSD^{14,26} includes individuals in whom anti-AQP4 is not detected. Approximately 20.6% of female and 39.8% of male cases in CIRCLES are anti-AQP4 seronegative, representing a significant difference based on sex ($p < 0.001$). Whether such proportions accurately reflect anti-AQP4 serostatus worldwide, correlate with specific disease phenotypes, or inform regarding response to therapy remains uncertain. Key among the proximate determinants of this serostatus are anti-AQP4 assay sensitivities and specificities. For example, it is possible that therapies inadvertently affect selection bias (e.g., rituximab targeting B cells). However, no consistent evidence published to date has proven these therapies alter antibody detection in Clinical Laboratory Improvement Amendments–approved assays. Seropositive cases are more likely to be female, and self-identified black/African American or Hispanic/Latino patients are more likely to be anti-AQP4 seropositive, congruent with earlier reports.⁵

Furthermore, cases in which anti-AQP4 is detected are older at first attack than those who are seronegative (39.4 vs 35.1, respectively, $p = 0.002$). Although white/Caucasian cases accounted for nearly 63% of the anti-AQP4 seronegative cohort, they account for less than 50% of seropositive cases. The age at first attack also differed by race, with a nearly 5-year disparity between the median age at onset for Hispanics/Latinos compared with white/Caucasians. This finding could reflect socioeconomic skewing of access to specialized medical care, a difference in disease activity/severity, other factor(s), or a combination of factors.

Of note, the prevalence of NMOSD appears to differ in distinct geographic regions. For example, the current estimate of NMOSD prevalence is 3.9 per 100,000 in Olmsted County, Minnesota (USA),⁵ similar to that reported in Denmark (4 per 100,000²⁷). In contrast, the prevalence ranges from 0.72 per 100,000 in England,²⁸ 0.89 per 100,000 in Spain,⁷ and 0.9 per 100,000 in Japan⁶ to 1.96 per 100,000 in Wales,²⁹ 2.5 per 100,000 in the French West Indies,³⁰ 2.6 per 100,000 in India,³¹ and as high as 10 per 100,000 in Martinique.⁵ It is possible that differences in diagnostic criteria could underlie at least some of these apparent differences. In any event, the current CIRCLES data offer insights extending those provided by recent reviews.^{8,32}

The sizable proportion of cases in which anti-AQP4 is not detected suggests that the NMOSD phenotype can result from multiple, independent immunologic events. The emerging recognition of individuals with phenotypes resembling NMOSD, but in whom anti-myelin oligodendrocyte glycoprotein (anti-MOG) autoantibodies are detected in the absence of anti-AQP4,^{10,33,34} suggests at least 2 intriguing possibilities: (1) a broader array of autoantigens than traditionally appreciated may contribute to astrocytopathies and/or (2) patients having anti-MOG autoantibodies may reflect a disease entity that is immunologically distinct from NMOSD, despite largely superimposable clinical manifestations. Thus, patients with anti-MOG antibodies may exhibit disease features that are pathogenically distinct from NMOSD, despite their similar clinical presentations.^{34,35}

Table 3 Case demographic factors by race and ethnicity

	N	Female (N = 599)	Age at onset (N = 593)	ARR (N = 593)
White	315	253 (80.3%)	41.1 (31.1–53.0)	0.5 (0.3–0.9)
Black or African American	141	130 (92.2%)	36.5 (26.1–45.5)	0.5 (0.3–0.9)
Hispanic or Latino	74	67 (90.5%)	36.0 (22.6–46.5)	0.5 (0.3–1.0)
Asian	54	43 (79.6%)	38.3 (25.2–50.2)	0.4 (0.2–0.7)
Other	15	11 (73.3%)	31.7 (22.2–43.1)	0.5 (0.2–0.9)
p Value	—	0.005 ^a	<0.001 ^b	0.342 ^b

Abbreviation: ARR = annualized relapse rate.

^a χ^2 test of association.

^b Analysis of variance.

Nevertheless, 1 recent epidemiologic study showed that individuals with detectable anti-MOG antibody had similar disease prevalence and long-term prognosis when compared with patients lacking detectable anti-AQP4 or anti-MOG antibody.¹⁰ Thus, the relationship between serology and disease phenotype remains to be more clearly understood.

As with all large, multicenter clinical research studies, there are limitations to CIRCLES. Some participants inconsistently followed up or provided incomplete historical information. Complete acquisition of PSF data elements, biospecimens, and fully documented neurologic examination data has proven challenging. In turn, acquiring disability data from incomplete neurologic examinations has emerged as a high priority for improvement. For example, although 90.9% of examinations included completed motor function assessments, only 49.5% assessed visual acuity, and 8.7% completed the Expanded Disability Status Scale. The study has not systematically monitored anti-MOG serostatus,¹⁶ as no approved clinical assay for this autoantibody existed during the study period.

A unique aspect of CIRCLES concerns the detailed clinical history captured at enrollment. For those whose disease is of long duration, recall bias is possible. Beyond basic clinical data, the CIRCLES PSF requests extensive retrospective clinical information, including history of infectious diseases, vaccinations, familial autoimmune diseases, medications, and treatments. Although collection of such extensive information is labor intensive, the resulting data set enables interrogations not possible from smaller or less comprehensive databases. Opportunities for enhancing study performance are currently being addressed through refinements of the study protocol. In particular, increased emphasis has been placed on longitudinal participation, more frequent site monitoring, and systematic methods for disseminating information to study sites pertaining to study performance and efficiency.

The design of CIRCLES allows direct comparisons between the clinical courses of NMOSD and other autoimmune diseases, informing key immunologic events unique to NMOSD. These events in turn facilitate identification of clinically useful biomarkers, including those heralding disease relapse, as well as novel therapeutic targets, agents, and strategies. The CIRCLES biobank contains clinical information and biospecimens from ethnically and geographically diverse cases, including those with heterotypic phenotypes. Thus, CIRCLES represents a unique resource to the academic and drug-discovery communities focused on finding solutions for patients with NMOSD and may enhance parallel efforts in other immune-based diseases.

In summary, areas of research urgently needed in NMOSD include discovery of disease etiology, identification of risk factors, and identification of biomarkers reflecting disease activity and predicting relapse.^{36,37} The ongoing CIRCLES continues to enroll and follow cases and categorical control participants in a systematic and longitudinal manner. This

effort is intended to facilitate breakthroughs regarding the epidemiology and pathogenesis of NMOSD, to reduce barriers to performing well-designed therapeutic trials, and to support postapproval studies of eventually approved therapeutics. Thus, the overarching goal of CIRCLES is to improve patient quality of life through improved diagnosis, relapse prevention, and eventual cures.^{38–40}

The CIRCLES program enables unprecedented opportunities to accelerate breakthroughs in scientific understanding and clinical solutions for NMOSD. Key to the future applicability of CIRCLES will be increased precision in diagnosis and uniformity in the assessment and specification of distinct disease phenotypes. These advances hinge on greater consistency of serologic analysis regarding autoantibodies specific to disease phenotype and standardization in the definition and severity scoring of NMOSD relapses. As these advances are made, they will be incorporated into the definitions used by CIRCLES, attesting to the evolving nature of this research platform. CIRCLES remains an open resource to facilitate hypothesis generation and testing. Given the nature of the biospecimens being collected, CIRCLES enables studies ranging from genomics, transcriptomics, proteomics, and other molecular- and cellular-based research, in addition to clinical investigation.

Acknowledgment

The authors recognize the efforts of the following individuals: Yanet Babcock, Alexandra Kocsik, J. Michael Dean, Colleen Farrell, Haojun Feng, Susan Filomena, Toni Ganaway, Samuel Glaisher, Rivka Green, Elizabeth Gonzales, Elaine Hsu, Catherine J. Johnson, Marlene Keymolen Ramirez, Gloria Rodriguez, Angela Stangorone, Ben W. Thrower, and Gabriella Tosto. The CIRCLES project team, including site PIs and coordinators, the DCC team, and The GJCF and its advisors are grateful to study participants, caregivers, and families who participated in this study. The CIRCLES project is supported by The Guthy-Jackson Charitable Foundation.

Study funding

The CIRCLES project is supported by The Guthy-Jackson Charitable Foundation.

Disclosure

L.J. Cook is supported in part by The Guthy-Jackson Charitable Foundation, which is a sponsor of this research. J.W. Rose is supported in part by The Guthy-Jackson Charitable Foundation, which is a sponsor of this research; he receives research funding from the National Multiple Sclerosis Society, Biogen, and NIH. J.S. Alvey, A.M. Jolley, R. Kuhn, B. Marron, M. Pederson, R. Enriquez, J. Yearley, and S. McKechnie are supported in part by The Guthy-Jackson Charitable Foundation, which is a sponsor of this research. M.H. Han served on the advisory committee of Novartis; received travel funding and/or speaker honoraria from the University of California, San Diego, and CMSC; consulted for Sanofi Genzyme; and received research support from The Guthy-Jackson Charitable Foundation and The Leducq Foundation.

A.J. Tomczak is supported in part by The Guthy-Jackson Charitable Foundation, which is a sponsor of this research. M. Levy is on the scientific advisory boards of Asterias, Chugai, and Alexion; is on the editorial board of *Multiple Sclerosis and Related Disorders*; holds a patent for aquaporin-4 sequence that elicits pathogenic T cell response in animal model of neuromyelitis optica; has consulted for Guidepoint Global, Gerson-Lehrman Group, and Cowen Group; and received research support from Viropharma/Shire, Acorda, ApoPharma, Sanofi, Genzyme, Alnylam, Alexion, Terumo BCT, NIH/NINDS, and The Guthy-Jackson Charitable Foundation. M.A. Mealy and J. Coleman are supported in part by The Guthy-Jackson Charitable Foundation, which is a sponsor of this research. J.L. Bennett is on the editorial boards of the *Journal of Neuro-ophthalmology*, *Multiple Sclerosis*, and *Neurology: Neuroimmunology & Neuroinflammation*; holds a patent for compositions and methods for the treatment of neuromyelitis optica; has consulted for AbbVie, EMD Serono, Mallinckrodt Pharmaceuticals, Equillum, MedImmune, Chugai, Frequency Therapeutics, Genentech, Genzyme, and Clene Nanomedicine; and received research support from EMD Serono, Mallinckrodt, NIH, and The Guthy-Jackson Charitable Foundation. R. Johnson and M. Barnes-Garcia are supported in part by The Guthy-Jackson Charitable Foundation, which is a sponsor of this research. A.L. Traboulsee has the following competing financial interests: research funding from Biogen, Chugai, Novartis, Roche, and Sanofi Genzyme; consultancy honoraria from Biogen, Roche, Sanofi Genzyme, and Teva Neuroscience; and supported in part by The Guthy-Jackson Charitable Foundation, which is a sponsor of this research. R.L. Carruthers is Site Investigator for studies funded by Roche, Novartis, MedImmune, and EMD Serono; receives research support from Teva Innovation Canada, Roche Canada, Vancouver Coastal Health Research Institute; and is supported in part by The Guthy-Jackson Charitable Foundation, which is a sponsor of this research. R.L. Carruthers has received honoraria from Roche, EMD Serono, Sanofi, Biogen, Novartis, and Teva. L.E. Lee, J.L. Schubert, and K. McMullen were supported in part by The Guthy-Jackson Charitable Foundation, which is a sponsor of this research. I. Kister served on the scientific advisory boards of Biogen and Genentech and received research support from The Guthy-Jackson Charitable Foundation, National Multiple Sclerosis Society, Biogen, EMD Serono, Genentech, Genzyme, and Novartis. Z. Rimler and A. Reid are supported in part by The Guthy-Jackson Charitable Foundation, which is a sponsor of this research. N.L. Sicotte is supported in part by the National Multiple Sclerosis Society, the Patient-Centered Outcomes Research Institute (PCORI), and The Guthy-Jackson Charitable Foundation, which is a sponsor of this research. S.M. Planchon is supported in part by The Guthy-Jackson Charitable Foundation, which is a sponsor of this research. J.A. Cohen reports personal compensation for consulting for Alkermes, Biogen, Convelo, EMD Serono, ERT, Gossamer Bio, Novartis, and ProValuate, speaking for Mylan and Synthon, and serving as an editor of *Multiple Sclerosis Journal*, and was supported in part by The Guthy-

Jackson Charitable Foundation, which is a sponsor of this research. D. Ivancic and J.L. Sedlak are supported in part by The Guthy-Jackson Charitable Foundation, which is a sponsor of this research. I.K. Sand receives research support from United States Department of Defense, the National Multiple Sclerosis Society, and The Guthy-Jackson Charitable Foundation. P. Repovic received travel funding and/or honoraria from Biogen, EMD Serono, Genzyme, and Genentech, consulting fees from Biogen, Novartis, Genentech, and EMD Serono, and research support from Genentech, National Multiple Sclerosis Society, and Consortium of MS Centers, and was supported in part by The Guthy-Jackson Charitable Foundation, which is a sponsor of this research. L. Amezcua has received personal compensation for consulting, serving on a scientific advisory board, speaking, or other activities with Genzyme. She receives research funding from Biogen, MedDay, NMSS, NIH NINDS, California Community Foundation, and The Guthy-Jackson Charitable Foundation. A. Pruitt and E. Amundson are supported in part by The Guthy-Jackson Charitable Foundation, which is a sponsor of this research. T. Chitnis served on the clinical advisory boards of Novartis, Celgene, and Sanofi Genzyme; consulted for Biogen, Celgene, Novartis, and Sanofi Genzyme; received research support from Merck Serono, Verily, NIH, MNSS, Peabody Foundation, and Consortium for MS Centers; and was supported in part by The Guthy-Jackson Charitable Foundation, which is a sponsor of the research. D.S. Mullin is supported in part by The Guthy-Jackson Charitable Foundation, which is a sponsor of this research. E.C. Klawiter has received consulting fees from Acorda Therapeutics, Atlas5d, Biogen Idec, EMD Serono, Genentech, and Shire and research support from Atlas5d, Biogen Idec, EMD Serono, and Roche and was supported in part by The Guthy-Jackson Charitable Foundation, which is a sponsor of this research. A.W. Russo is supported in part by The Guthy-Jackson Charitable Foundation, which is a sponsor of this research. C.S. Riley has participated in advisory boards and received honoraria from Novartis, Biogen Idec, Teva, Genentech, Roche, and Genzyme and was supported in part by The Guthy-Jackson Charitable Foundation, which is a sponsor of this research. K.B. Onomichi, L. Levine, K.E. Nelson, N.M. Nealon, and C. Engel are supported in part by The Guthy-Jackson Charitable Foundation, which is a sponsor of this research. M. Kruse-Hoyer reports no disclosures. M. Marcille, L. Tornes, A. Rumpf, A. Greer, M. Kenneally Behne, R.R. Rodriguez, D.W. Behne, D.W. Blackway, and B. Coords are supported in part by The Guthy-Jackson Charitable Foundation, which is a sponsor of this research. T.F. Blaschke is an advisor to The Guthy-Jackson Charitable Foundation, which is a sponsor of this research. J. Sheard is supported in part by The Guthy-Jackson Charitable Foundation, which is a sponsor of this research. T.J. Smith is an advisor to The Guthy-Jackson Charitable Foundation, which is a sponsor of this research. J.M. Behne is supported in part by The Guthy-Jackson Charitable Foundation, which is a sponsor of this research. M.R. Yeaman is supported by research funding from the National Institutes of Health; U.S. Department of Defense; is Founder and

Shareholder of NovaDigm Therapeutics, Inc.; is Founder and Shareholder of Metacin, Inc; and is Advisor to The Guthy-Jackson Charitable Foundation, which is a sponsor of this research. Go to Neurology.org/NN for full disclosures.

Publication history

Received by *Neurology: Neuroimmunology & Neuroinflammation* February 21, 2019. Accepted in final form May 17, 2019.

Appendix 1 Authors

Name	Location	Role	Contribution
Lawrence J. Cook, PhD, MStat	University of Utah School of Medicine, Salt Lake City, UT	Author	Participated in study design; collected study data; analyzed and interpreted data; and drafted and revised the manuscript for intellectual content.
John W. Rose, MD	University of Utah School of Medicine, Salt Lake City, UT	Author	Participated in study design and reviewed and revised the manuscript for intellectual content.
Jessica S. Alvey, MS	University of Utah School of Medicine, Salt Lake City, UT	Author	Participated in study design; collected study data; analyzed and interpreted data; and drafted and revised the manuscript for intellectual content.
Anna Marie Jolley, BS	University of Utah School of Medicine, Salt Lake City, UT	Author	Participated in study design; collected study data; and reviewed and revised the manuscript for intellectual content.
Renee Kuhn, BS, CCRP	University of Utah School of Medicine, Salt Lake City, UT	Author	Participated in study design; collected study data; and reviewed the manuscript for intellectual content.
Brie Marron, BS	University of Utah School of Medicine, Salt Lake City, UT	Author	Collected and analyzed study data and reviewed the manuscript for intellectual content.
Melissa Pederson, MEd	University of Utah School of Medicine, Salt Lake City, UT	Author	Collected and analyzed study data and reviewed the manuscript for intellectual content.
Rene Enriquez, BS	University of Utah School of Medicine, Salt Lake City, UT	Author	Collected and analyzed data and reviewed and revised the manuscript for intellectual content.
Jeff Yearley, BA	University of Utah School of Medicine, Salt Lake City, UT	Author	Collected and analyzed data and reviewed and revised the manuscript for intellectual content.

Appendix 1 (continued)

Name	Location	Role	Contribution
Stephen McKechnie, MS	University of Utah School of Medicine, Salt Lake City, UT	Author	Participated in study design; collected study data; analyzed and interpreted data; and drafted and revised the manuscript for intellectual content.
May H. Han, MD	Department of Neurology and Neurological Sciences, Division of Neuroimmunology and Multiple Sclerosis Center, Stanford University, Stanford, CA	Author	Participated in study design; performed clinical assessments; collected study data and biospecimens; and reviewed and revised the manuscript for intellectual content.
Anna J. Tomczak, MS	Department of Neurology and Neurological Sciences, Stanford School of Medicine, Stanford, CA	Author	Collected study data and reviewed/revised the manuscript for intellectual content.
Michael Levy, MD, PhD	Department of Neurology, Johns Hopkins University School of Medicine, MD; Massachusetts General Hospital, Harvard Medical School, Boston, MA	Author	Participated in study design; performed clinical assessments; collected study data and biospecimens; and reviewed and revised the manuscript for intellectual content.
Maureen A. Mealy, BSN, MSCN	Department of Neurology, Johns Hopkins University School of Medicine, MD	Author	Collected study data and reviewed the manuscript for intellectual content.
Jessica Coleman, BA	Department of Neurology, Johns Hopkins University School of Medicine, MD	Author	Collected study data and reviewed the manuscript for intellectual content.
Jeffrey L. Bennett, MD, PhD	Departments of Neurology and Ophthalmology, University of Colorado School of Medicine, Aurora, CO	Author	Participated in study design; performed clinical assessments; collected study data and biospecimens; and reviewed and revised the manuscript for intellectual content.
Ruth Johnson, BS	Departments of Neurology and Ophthalmology, University of Colorado School of Medicine, Aurora, CO	Author	Collected study data and reviewed the manuscript for intellectual content.
Myka Barnes-Garcia, BA	Departments of Neurology and Ophthalmology, University of Colorado School of Medicine, Aurora, CO	Author	Collected study data and reviewed the manuscript for intellectual content.

Appendix 1 (continued)

Name	Location	Role	Contribution
Anthony L. Traboulsee, MD	Department of Medicine & Neurology, University of British Columbia, Vancouver, BC, Canada	Author	Participated in study design; performed clinical assessments; collected study data and biospecimens; and reviewed the manuscript for intellectual content.
Robert L. Carruthers, MD	Department of Medicine & Neurology, University of British Columbia, Vancouver, BC, Canada	Author	Participated in study design; performed clinical assessments; collected study data and biospecimens; and reviewed the manuscript for intellectual content.
Lisa Eunyong Lee, BS, MSc	Department of Medicine & Neurology, University of British Columbia, Vancouver, BC, Canada	Author	Participated in study design; collected study data; analyzed and interpreted the data; and reviewed/ revised the manuscript for intellectual content.
Julia J. Schubert, BS, BCS	Department of Medicine & Neurology, University of British Columbia, Vancouver, BC, Canada	Author	Collected study data and reviewed the manuscript for intellectual content.
Katrina McMullen, PhD	Department of Medicine & Neurology, University of British Columbia, Vancouver, BC, Canada	Author	Collected study data and reviewed the manuscript for intellectual content.
Ilya Kister, MD	NYU Langone Health, New York, NY	Author	Participated in study design; performed clinical assessments; collected study data and biospecimens; and reviewed and revised the manuscript for intellectual content.
Zoe Rimler, BS	NYU Langone Health, New York, NY	Author	Collected study data and reviewed the manuscript for intellectual content.
Allyson Reid, BA	NYU Langone Health, New York, NY	Author	Collected study data and reviewed the manuscript for intellectual content.
Nancy L. Sicotte, MD	Department of Neurology, Cedars-Sinai Medical Center, Los Angeles, CA	Author	Performed clinical assessments; collected study data and biospecimens; and reviewed and revised the manuscript for intellectual content.
Sarah M. Planchon, PhD	Mellen Center for MS Treatment and Research, Neurological Institute, Cleveland Clinic, Cleveland, OH	Author	Participated in study design; collected study data and biospecimens; and reviewed and revised the manuscript for intellectual content.

Appendix 1 (continued)

Name	Location	Role	Contribution
Jeffery A. Cohen, MD	Mellen Center for MS Treatment and Research, Neurological Institute, Cleveland Clinic, Cleveland, OH	Author	Performed clinical assessments; collected study data and biospecimens; and reviewed and revised the manuscript for intellectual content.
Diane Ivancic, CCRP	Mellen Center for MS Treatment and Research, Neurological Institute, Cleveland Clinic, Cleveland, OH	Author	Collected study data and reviewed the manuscript for intellectual content.
Jennifer L. Sedlak, BSN	Mellen Center for MS Treatment and Research, Neurological Institute, Cleveland Clinic, Cleveland, OH	Author	Collected study data and reviewed the manuscript for intellectual content.
Ilana Katz Sand, MD	Icahn School of Medicine at Mount Sinai, NY	Author	Participated in study design; performed clinical assessments; collected study data and biospecimens; and reviewed and revised the manuscript for intellectual content.
Pavle Repovic, MD, PhD	Multiple Sclerosis Center, Swedish Neuroscience Institute, Seattle, WA	Author	Participated in study design and reviewed and revised the manuscript for intellectual content.
Lilyana Amezcua, MD, MS	Department of Neurology, Keck School of Medicine, University of Southern California, Los Angeles, CA	Author	Participated in study design; performed clinical assessments; collected study data and biospecimens; and reviewed and revised the manuscript for intellectual content.
Ana Pruitt, BS	Department of Neurology, Keck School of Medicine, University of Southern California, Los Angeles, CA	Author	Collected study data and reviewed the manuscript for intellectual content.
Erika Amundson, BA	Department of Neurology, Keck School of Medicine, University of Southern California, Los Angeles, CA	Author	Collected study data and reviewed the manuscript for intellectual content.
Tanuja Chitnis, MD	Department of Neurology, Brigham and Women's Hospital, Harvard Medical School, Boston, MA	Author	Participated in study design; performed clinical assessments; collected study data and biospecimens; and reviewed and revised the manuscript for intellectual content.
Devin S. Mullin, BS	Department of Neurology, Brigham and Women's Hospital, Harvard Medical School, Boston, MA	Author	Collected study data and reviewed the manuscript for intellectual content.

Continued

Appendix 1 (continued)

Name	Location	Role	Contribution
Eric C. Klawiter, MD	Department of Neurology, Massachusetts General Hospital, Harvard Medical School, Boston, MA	Author	Participated in study design; performed clinical assessments; collected study data and biospecimens; and reviewed and revised the manuscript for intellectual content.
Andrew W. Russo, BS	Department of Neurology, Massachusetts General Hospital, Harvard Medical School, Boston, MA	Author	Collected study data and reviewed the manuscript for intellectual content.
Claire S. Riley, MD	Department of Neurology, Columbia University Medical Center, New York, NY	Author	Participated in study design; performed clinical assessments; collected study data and biospecimens; and reviewed and revised the manuscript for intellectual content.
Kaho B. Onomichi, MS	Department of Neurology, Columbia University Medical Center, New York, NY	Author	Collected study data and reviewed the manuscript for intellectual content.
Libby Levine, RN, ANP-BC	Department of Neurology, Columbia University Medical Center, NY, NY	Author	Collected study data and reviewed the manuscript for intellectual content.
Katherine E. Nelson, BA	Department of Neurology, Columbia University Medical Center, NY, NY	Author	Collected study data and reviewed the manuscript for intellectual content.
Nancy M. Nealon, MD	Weill Cornell Medicine, New York, NY	Author	Participated in study design; performed clinical assessments; collected study data and biospecimens; and reviewed and revised the manuscript for intellectual content.
Casey Engel, BA	Weill Cornell Medicine, New York, NY	Author	Collected study data and reviewed the manuscript for intellectual content.
Mason Kruse-Hoyer, MD, MA	Weill Cornell Medicine, New York, NY	Author	Collected study data and reviewed the manuscript for intellectual content.
Melanie Marcille, BA	Weill Cornell Medicine, New York, NY	Author	Collected study data and reviewed the manuscript for intellectual content.
Leticia Tornes, MD	Department of Neurology, Division of Multiple Sclerosis, University of Miami Miller School of Medicine, Miami, FL	Author	Participated in study design; performed clinical assessments; collected study data and biospecimens; and reviewed and revised the manuscript for intellectual content.

Appendix 1 (continued)

Name	Location	Role	Contribution
Anne Rumpf, BS	PPD, Wilmington, NC	Author	Reviewed and curated data and reviewed the manuscript for intellectual content.
Angela Greer, BS	PPD, Wilmington, NC	Author	Reviewed and curated data and reviewed the manuscript for intellectual content.
Megan Kenneally Behne, AS	The Guthy-Jackson Charitable Foundation, Beverly Hills, CA	Author	Reviewed and revised the manuscript for intellectual content.
Renee R. Rodriguez, AS	The Guthy-Jackson Charitable Foundation, Beverly Hills, CA	Author	Reviewed and revised the manuscript for intellectual content.
Daniel W. Behne, MArch	The Guthy-Jackson Charitable Foundation, Beverly Hills, CA	Author	Reviewed and revised the manuscript for intellectual content.
Derek W. Blackway, BA	The Guthy-Jackson Charitable Foundation, Beverly Hills, CA	Author	Reviewed and revised the manuscript for intellectual content.
Brian Coords, MA	The Guthy-Jackson Charitable Foundation, Beverly Hills, CA	Author	Reviewed and revised the manuscript for intellectual content.
Terrence F. Blaschke, MD	Departments of Medicine and of Molecular Pharmacology, Stanford University School of Medicine, Stanford, CA	Author	Participated in study design; analyzed and interpreted data; and reviewed and revised the manuscript for intellectual content.
Judy Sheard, MPH, MA, CCRC	The Guthy-Jackson Charitable Foundation, Beverly Hills, CA	Author	Reviewed and revised the manuscript for intellectual content.
Terry J. Smith, MD	Kellogg Eye Center, University of Michigan Medical School, Ann Arbor, MI	Author	Participated in study design; analyzed and interpreted data; and drafted and revised the manuscript for intellectual content.
Jacinta M. Behne, MA	The Guthy-Jackson Charitable Foundation, Beverly Hills, CA	Author	Designed/ conceptualized study; interpreted the data; and revised the manuscript for intellectual content.
Michael R. Yeaman, PhD	Department of Medicine, University of California, Los Angeles, Los Angeles CA; Harbor-UCLA Medical Center/ LABioMed, Torrance, CA	Author	Participated in study design; analyzed and interpreted data; and drafted and revised the manuscript for intellectual content.

Appendix 2 Coinvestigators

Name	Location	Role	Contribution
Hesham Abboud, MD	Multiple Sclerosis and Neuroimmunology Program, University Hospitals of Cleveland, Case Western Reserve University, Cleveland, OH	Coinvestigator	Reviewed and revised the manuscript for intellectual content.
Orhan Aktas, MD	Heinrich Heine Universität, Düsseldorf, Germany	Coinvestigator	Reviewed and revised the manuscript for intellectual content.
Ayse Altintas, MD	Department of Neurology, Koc University, School of Medicine, Istanbul, Turkey	Coinvestigator	Reviewed and revised the manuscript for intellectual content.
Metha Apiwattanakul, MD	Department of Neurology, Prasat Neurological Institute, Bangkok, Thailand	Coinvestigator	Reviewed and revised the manuscript for intellectual content.
Nasrin Asgari, MD, PhD	Department of Neurology, Slagelse Hospital and Institute of Regional Health Research & Molecular Medicine, University of Southern Denmark, Odense, Denmark	Coinvestigator	Reviewed and revised the manuscript for intellectual content.
Brenda Banwell, MD	Children's Hospital of Philadelphia, Perelman School of Medicine, University of Pennsylvania	Coinvestigator	Reviewed and revised the manuscript for intellectual content.
Denis Bichuetti, MD	Professor of Neurology, Escola Paulista de Medicina-Universidade Federal de São Paulo, Brazil	Coinvestigator	Reviewed and revised the manuscript for intellectual content.
James Bowen, MD	Multiple Sclerosis Center, Swedish Neuroscience Institute, Seattle, WA	Coinvestigator	Reviewed and revised the manuscript for intellectual content.
Simon Broadley, MD, PhD	Griffith University, Queensland, Australia	Coinvestigator	Reviewed and revised the manuscript for intellectual content.
Wolfgang Bruck, MD	University Medical Center Göttingen, Germany	Coinvestigator	Reviewed and revised the manuscript for intellectual content.
Philippe Cabre, MD	CHU Pierre Zobda Quitman, Martinique, French West Indies	Coinvestigator	Reviewed and revised the manuscript for intellectual content.

Appendix 2 (continued)

Name	Location	Role	Contribution
Jeffrey Cohen, MD	Mellen Center for MS Treatment and Research, Neurological Institute, Cleveland Clinic, Cleveland, OH	Coinvestigator	Reviewed and revised the manuscript for intellectual content.
Jerome De Seze, MD, PhD	Chu de Strasbourg, CIC Inserm 1434, France	Coinvestigator	Reviewed and revised the manuscript for intellectual content.
Guillermo Delgado-Garcia, MD	Division of Neurology, National Institute of Neurology and Neurosurgery, Mexico City, Mexico	Coinvestigator	Reviewed and revised the manuscript for intellectual content.
Irena Dujmovic Basuroski, MD, PhD	University of North Carolina at Chapel Hill, Department of Neurology, Chapel Hill, NC	Coinvestigator	Reviewed and revised the manuscript for intellectual content.
Kazuo Fujihara, MD	Department of Multiple Sclerosis Therapeutics, Fukushima Medical University, Fukushima, Japan and Multiple Sclerosis and Neuromyelitis Optica Center, Tohoku Research Institute for Neuroscience, Koriyama, Japan	Coinvestigator	Reviewed and revised the manuscript for intellectual content.
Andrew Goodman, MD	Department of Neurology, University of Rochester Medical Center, Rochester, NY	Coinvestigator	Reviewed and revised the manuscript for intellectual content.
Joachim Havla, MD	Institute of Clinical Neuroimmunology, Ludwig Maximilians University, Munich, Germany	Coinvestigator	Reviewed and revised the manuscript for intellectual content.
Kerstin Hellwig, MD	Department of Neurology, St. Josef Hospital Bochum, Ruhr University, Bochum, Germany	Coinvestigator	Reviewed and revised the manuscript for intellectual content.
Rogier Hintzen, MD, PhD	MS Centre ErasMS, Dept of Neurology, Erasmus MC, Rotterdam, The Netherlands	Coinvestigator	Reviewed and revised the manuscript for intellectual content.
D. Craig Hooper, PhD	Department of Cancer Biology, Thomas Jefferson University, Philadelphia, PA	Coinvestigator	Reviewed and revised the manuscript for intellectual content.
Raffaele Iorio, MD, PhD	Institute of Neurology, Fondazione Policlinico Universitario "A. Gemelli" IRCCS, Università Cattolica del Sacro Cuore, Roma, Italy	Coinvestigator	Reviewed and revised the manuscript for intellectual content.

Continued

Appendix 2 (continued)

Name	Location	Role	Contribution
Anu Jacob, MD	The Walton Centre NHS Trust and University of Liverpool, United Kingdom	Coinvestigator	Reviewed and revised the manuscript for intellectual content.
Sven Jarius, MD	Molecular Neuroimmunology Group, University of Heidelberg, Heidelberg, Germany	Coinvestigator	Reviewed and revised the manuscript for intellectual content.
Jorge Andres Jimenez Arango, MD	Universidad de Antioquia, Neuroclinica, Colombia	Coinvestigator	Reviewed and revised the manuscript for intellectual content.
Gareth John, PhD	The Mount Sinai Hospital, New York, NY	Coinvestigator	Reviewed and revised the manuscript for intellectual content.
Ho Jin Kim, MD, PhD	Department of Neurology, Research Institute and Hospital of National Cancer Center, Goyang, Republic of Korea	Coinvestigator	Reviewed and revised the manuscript for intellectual content.
Sung Min Kim, MD, PhD	Department of Neurology, Seoul National University Hospital, Seoul, Korea	Coinvestigator	Reviewed and revised the manuscript for intellectual content.
Dorian J. Kimbrough, MD	Harvard Medical School, Brigham & Women's Hospital Department of Neurology, Boston, MA	Coinvestigator	Reviewed and revised the manuscript for intellectual content.
Najib Kissani, MD	Department of Neurology, Mohamed VI University Hospital; Neuroscience Research Laboratory, Marrakech Medical School; UCA, Marrakech, Morocco	Coinvestigator	Reviewed and revised the manuscript for intellectual content.
Ingo Kleiter, MD	Marianne-Strauß-Klinik, Behandlungszentrum Kempfenhausen für Multiple Sklerose Kranke, Berg, Germany; Department of Neurology, St. Josef Hospital, Ruhr University Bochum, Bochum, Germany	Coinvestigator	Reviewed and revised the manuscript for intellectual content.
Marco Lana-Peixoto, MD, PhD	Federal University of Minas Gerais Medical School, Belo Horizonte, Brazil	Coinvestigator	Reviewed and revised the manuscript for intellectual content.
Annette Langer-Gould, MD, PhD	Kaiser Permanente Southern California, CA	Coinvestigator	Reviewed and revised the manuscript for intellectual content.

Appendix 2 (continued)

Name	Location	Role	Contribution
M. Isabel Leite, MD, D Phil	Nuffield Department of Clinical Neurosciences, University of Oxford and Oxford University Hospitals, Headley Way, Oxford OX3 9DU, UK	Coinvestigator	Reviewed and revised the manuscript for intellectual content.
Yaou Liu, MD, PhD	Department of Radiology, Beijing Tiantan Hospital, Capital Medical University, Beijing 100050; China	Coinvestigator	Reviewed and revised the manuscript for intellectual content.
Fred Lublin, MD	Mount Sinai Medical Center, New York, NY	Coinvestigator	Reviewed and revised the manuscript for intellectual content.
Yousseoufa Maiga, MD	Faculty of Medicine, University of Technical Sciences and Technologies, Bamako, Mali	Coinvestigator	Reviewed and revised the manuscript for intellectual content.
Yang Mao-Draayer, MD, PhD	Graduate Program in Immunology, Program in Biomedical Sciences, Department of Neurology, University of Michigan Medical School	Coinvestigator	Reviewed and revised the manuscript for intellectual content.
Romain Marignier, MD, PhD	Service de neurologie, sclérose en plaques, pathologies de la myéline et neuro-inflammation and Centre de référence pour les maladies inflammatoires rares du cerveau et de la moelle (MIRCEM) – Hôpital Neurologique Pierre Wertheimer Hospices Civils de Lyon, Lyon, F-69777, France	Coinvestigator	Reviewed and revised the manuscript for intellectual content.
Marcelo Matiello, MD	Assistant Professor of Neurology, Harvard Medical School	Coinvestigator	Reviewed and revised the manuscript for intellectual content.
Callene Momtazee, MD	Department of Neurology, University of California, Los Angeles, Los Angeles, CA	Coinvestigator	Reviewed and revised the manuscript for intellectual content.
Mark Morrow, MD	Department of Neurology, Harbor-UCLA Medical Center, Torrance, CA; David Geffen School of Medicine, Los Angeles, CA	Coinvestigator	Reviewed and revised the manuscript for intellectual content.
Ichiro Nakashima, MD	Department of Neurology, Tohoku Medical and Pharmaceutical University, Sendai, Japan	Coinvestigator	Reviewed and revised the manuscript for intellectual content.

Appendix 2 (continued)

Name	Location	Role	Contribution
Kevin O'Connor, PhD	Yale University School of Medicine, New Haven, CT	Coinvestigator	Reviewed and revised the manuscript for intellectual content.
Celia Oreja-Guevara, MD, PhD	Hospital Clinico San Carlos, Neurology and Universidad Complutense Madrid, Spain	Coinvestigator	Reviewed and revised the manuscript for intellectual content.
Jacqueline Palace, MD	Department of Neurology, Oxford University Hospital Trust, Oxford, United Kingdom	Coinvestigator	Reviewed and revised the manuscript for intellectual content.
Lekha Pandit, MD, PhD	Nitte University, Mangalore, India	Coinvestigator	Reviewed and revised the manuscript for intellectual content.
Friedemann Paul, MD	Charité – Universitätsmedizin Berlin, corporate member of Freie Universität Berlin, Humboldt-Universität zu Berlin, and Berlin Institute of Health, NeuroCure Cluster of Excellence, NeuroCure Clinical Research Center, Berlin, Germany; Experimental and Clinical Research Center, Charité – Universitätsmedizin Berlin corporate member of Freie Universität Berlin, Humboldt-Universität zu Berlin, and Berlin Institute of Health and Max Delbrück Center for Molecular Medicine, Berlin-Buch, Germany	Coinvestigator	Reviewed and revised the manuscript for intellectual content.
Naraporn Prayoonwiwat, MD	Division of Neurology, Department of Medicine, Faculty of Medicine Siriraj Hospital, Mahidol University, Bangkok, Thailand	Coinvestigator	Reviewed and revised the manuscript for intellectual content.
Anne-Katrin Pröbstel, MD	Neurologic Clinic and Polyclinic, Departments of Medicine, and Biomedicine, University Hospital, University of Basel, Basel, Switzerland; UCSF Weill Institute for Neurosciences, Department of Neurology, University of California San Francisco, San Francisco, CA	Coinvestigator	Reviewed and revised the manuscript for intellectual content.
Peiqing Qian, MD	MS Center at Swedish Medical Center, Seattle, WA	Coinvestigator	Reviewed and revised the manuscript for intellectual content.

Appendix 2 (continued)

Name	Location	Role	Contribution
Chao Quan, MD	Department of Neurology, Huashan Hospital, Shanghai Medical College, Fudan University	Coinvestigator	Reviewed and revised the manuscript for intellectual content.
Marius Ringelstein, MD	Department of Neurology, Medical Faculty, Heinrich Heine University Düsseldorf, Düsseldorf, Germany	Coinvestigator	Reviewed and revised the manuscript for intellectual content.
Victor Rivera, MD	Baylor College of Medicine, Houston, TX	Coinvestigator	Reviewed and revised the manuscript for intellectual content.
Dalia L. Rotstein, MD	University of Toronto, Department of Medicine, Division of Neurology	Coinvestigator	Reviewed and revised the manuscript for intellectual content.
Klemens Ruprecht, MD	Department of Neurology, Charité – Universitätsmedizin Berlin, corporate member of Freie Universität Berlin, Humboldt-Universität zu Berlin, and Berlin Institute of Health, Berlin, Germany	Coinvestigator	Reviewed and revised the manuscript for intellectual content.
Maria José Sá, MD, PhD	Neurology Department, Centro Hospitalar de São João, FP-ENAS (UFP Energy, Environment and Health Research Unit), Faculty of Health Sciences, University Fernando Pessoa, Porto, Portugal	Coinvestigator	Reviewed and revised the manuscript for intellectual content.
Albert Saiz, MD, PhD	Hospital Clinic and Institut d' Investigació August Pi i Sunyer (IDIBAPS), University of Barcelona, Barcelona, Spain	Coinvestigator	Reviewed and revised the manuscript for intellectual content.
Ché Serguera, MD, PhD	CEA, Molecular Imaging Research Center (MIRCen), INSERM, Fontenay-aux-Roses, France	Coinvestigator	Reviewed and revised the manuscript for intellectual content.
Eslam Shosha, MD	College of Medicine, Al Majmaah University, Riyadh, Saudi Arabia	Coinvestigator	Reviewed and revised the manuscript for intellectual content.
Sasitorn Siritho, MD	Multiple Sclerosis and Related Disorder Clinics, Siriraj Hospital, Mahidol University, Bangkok, Thailand; Bumrungrad International Hospital, Bangkok, Thailand	Coinvestigator	Reviewed and revised the manuscript for intellectual content.

Continued

Appendix 2 (continued)

Name	Location	Role	Contribution
Aksel Siva, MD	Department of Neurology, Clinical Neuroimmunology Unit, Istanbul University Cerrahpasa School of Medicine, Istanbul, Turkey	Coinvestigator	Reviewed and revised the manuscript for intellectual content.
Ibis Soto de Castillo, MD	Neurology Department, Hospital Universitario de Maracaibo, Maracaibo, Venezuela	Coinvestigator	Reviewed and revised the manuscript for intellectual content.
Olaf Stuve, MD, PhD	UT Southwestern Medical Center, Dallas, TX	Coinvestigator	Reviewed and revised the manuscript for intellectual content.
Silvia Tenenbaum, MD	Department of Neurology, National Pediatric Hospital Dr. J. P. Garrahan, Buenos Aires, Argentina	Coinvestigator	Reviewed and revised the manuscript for intellectual content.
Pablo Villoslada, MD	Institute d'Investigacions Biomediques August Pi Sunyer (IDIBAPS), Barcelona, Spain	Coinvestigator	Reviewed and revised the manuscript for intellectual content.
Dean Wingerchuk, MD	Mayo Clinic, Scottsdale, AZ	Coinvestigator	Reviewed and revised the manuscript for intellectual content.
Jens Würfel, MD	MIAC AG, Basel, Switzerland	Coinvestigator	Reviewed and revised the manuscript for intellectual content.
E. Ann Yeh, MD	Division of Neurology, Hospital for Sick Children, University of Toronto, Canada	Coinvestigator	Reviewed and revised the manuscript for intellectual content.
Scott S. Zamvil, MD, PhD	Department of Neurology and Program in Immunology, University of California, San Francisco, San Francisco, CA	Coinvestigator	Reviewed and revised the manuscript for intellectual content.

References

- Akaishi T, Nakashima I, Sato DK, Takahashi T, Fujihara K. Neuromyelitis optica spectrum disorders. *Neuroimaging Clin N Am* 2017;27:251–265.
- Marignier R, Cobo Calvo A, Vukusic S. Neuromyelitis optica and neuromyelitis optica spectrum disorders. *Curr Opin Neurol* 2017;30:208–215.
- Weinshenker BG, Wingerchuk DM. Neuromyelitis spectrum disorders. *Mayo Clin Proc* 2017;92:663–679.
- Whittam D, Wilson M, Hamid S, Keir G, Bhojak M, Jacob A. What's new in neuromyelitis optica? A short review for the clinical neurologist. *J Neurol* 2017;264:2330–2344.
- Flanagan EP, Cabre P, Weinshenker BG, et al. Epidemiology of aquaporin-4 autoimmunity and neuromyelitis optica spectrum. *Ann Neurol* 2016;79:775–783.
- Houzen H, Kondo K, Niino M, et al. Prevalence and clinical features of neuromyelitis optica spectrum disorders in northern Japan. *Neurology* 2017;89:1995–2001.

- Sepulveda M, Aldea M, Escudero D, et al. Epidemiology of NMOSD in Catalonia: influence of the new 2015 criteria in incidence and prevalence estimates. *Mult Scler* 2017 Epub 2017 Oct 1.
- Pandit L, Asgari N, Apiwattanakul M, et al. Demographic and clinical features of neuromyelitis optica: a review. *Mult Scler* 2015;21:845–853.
- Borisow N, Kleiter I, Gahlen A, et al. Influence of female sex and fertile age on neuromyelitis optica spectrum disorders. *Mult Scler* 2017;23:1092–1103.
- Sepulveda M, Armangue T, Sola-Valls N, et al. Neuromyelitis optica spectrum disorders: comparison according to the phenotype and serostatus. *Neurol Neuroimmunol Neuroinflamm* 2016;3:e225. doi:10.1212/NXI.0000000000000225.
- Kim HJ, Paul F, Lana-Peixoto MA, et al. MRI characteristics of neuromyelitis optica spectrum disorder: an international update. *Neurology* 2015;84:1165–1173.
- Paul F, Murphy O, Pardo S, Levy M. Investigational drugs in development to prevent neuromyelitis optica relapses. *Expert Opin Investig Drugs* 2018;27:265–271.
- Trebst C, Jarius S, Berthele A, et al. Update on the diagnosis and treatment of neuromyelitis optica: recommendations of the Neuromyelitis Optica Study Group (NEMOS). *J Neurol* 2014;261:1–16.
- Wingerchuk DM, Banwell B, Bennett JL, et al. International consensus diagnostic criteria for neuromyelitis optica spectrum disorders. *Neurology* 2015;85:177–189.
- Jarius S, Kleiter I, Ruprecht K, et al. MOG-IgG in NMO and related disorders: a multicenter study of 50 patients. Part 3: brainstem involvement—frequency, presentation and outcome. *J Neuroinflammation* 2016;13:281.
- Jarius S, Ruprecht K, Kleiter I, et al. MOG-IgG in NMO and related disorders: a multicenter study of 50 patients. Part 2: epidemiology, clinical presentation, radiological and laboratory features, treatment responses, and long-term outcome. *J Neuroinflammation* 2016;13:280.
- Jarius S, Ruprecht K, Kleiter I, et al. MOG-IgG in NMO and related disorders: a multicenter study of 50 patients. Part 1: frequency, syndrome specificity, influence of disease activity, long-term course, association with AQP4-IgG, and origin. *J Neuroinflammation* 2016;13:279.
- Kleiter I, Gahlen A, Borisow N, et al. Neuromyelitis optica: evaluation of 871 attacks and 1,153 treatment courses. *Ann Neurol* 2016;79:206–216.
- Mealy MA, Wingerchuk DM, Greenberg BM, Levy M. Epidemiology of neuromyelitis optica in the United States: a multicenter analysis. *Arch Neurol* 2012;69:1176–1180.
- Pache F, Zimmermann H, Mikolajczak J, et al. MOG-IgG in NMO and related disorders: a multicenter study of 50 patients. Part 4: afferent visual system damage after optic neuritis in MOG-IgG-seropositive versus AQP4-IgG-seropositive patients. *J Neuroinflammation* 2016;13:282.
- Collongues N, Marignier R, Zephir H, et al. Neuromyelitis optica in France: a multicenter study of 125 patients. *Neurology* 2010;74:736–742.
- Jarius S, Ruprecht K, Wildemann B, et al. Contrasting disease patterns in seropositive and seronegative neuromyelitis optica: a multicenter study of 175 patients. *J Neuroinflammation* 2012;9:14.
- Ngo ST, Steyn FJ, McCombe PA. Gender differences in autoimmune disease. *Front Neuroendocrinol* 2014;35:347–369.
- Tackley G, O'Brien F, Rocha J, et al. Neuromyelitis optica relapses: race and rate, immunosuppression and impairment. *Mult Scler Relat Disord* 2016;7:21–25.
- Zhangbao J, Zhou L, Li X, et al. The clinical characteristics of AQP4 antibody positive NMO/SD in a large cohort of Chinese Han patients. *J Neuroimmunol* 2017;302:49–55.
- Wingerchuk DM, Lennon VA, Pittock SJ, Lucchinetti CF, Weinshenker BG. Revised diagnostic criteria for neuromyelitis optica. *Neurology* 2006;66:1485–1489.
- Asgari N, Lillevang ST, Skejoe HP, Falah M, Stenager E, Kyvik KO. A population-based study of neuromyelitis optica in Caucasians. *Neurology* 2011;76:1589–1595.
- Jacob A, Panicker J, Lythgoe D, et al. The epidemiology of neuromyelitis optica amongst adults in the Merseyside county of United Kingdom. *J Neurol* 2013;260:2134–2137.
- Cosburn M, Tackley G, Baker K, et al. The prevalence of neuromyelitis optica in South East Wales. *Eur J Neurol* 2012;19:655–659.
- Cabre P, Gonzalez-Quevedo A, Lannuzel A, et al. Descriptive epidemiology of neuromyelitis optica in the Caribbean basin [in French]. *Rev Neurol (Paris)* 2009;165:676–683.
- Pandit L, Kundapur R. Prevalence and patterns of demyelinating central nervous system disorders in urban Mangalore, South India. *Mult Scler* 2014;20:1651–1653.
- Mori M, Kuwabara S, Paul F. Worldwide prevalence of neuromyelitis optica spectrum disorders. *J Neurol Neurosurg Psychiatry* 2018;89:555–556.
- Spadaro M, Gerdes LA, Krumbholz M, et al. Autoantibodies to MOG in a distinct subgroup of adult multiple sclerosis. *Neurol Neuroimmunol Neuroinflamm* 2016;3:e257. doi:10.1212/NXI.0000000000000257.
- Zamvil SS, Slavin AJ. Does MOG Ig-positive AQP4-seronegative opticospinal inflammatory disease justify a diagnosis of NMO spectrum disorder? *Neurol Neuroimmunol Neuroinflamm* 2015;2:e62. doi:10.1212/NXI.0000000000000062.
- Reindl M, Rostasy K. MOG antibody-associated diseases. *Neurol Neuroimmunol Neuroinflamm* 2015;2:e60. doi:10.1212/NXI.0000000000000060.
- Kremer S, Renard F, Achard S, et al. Use of advanced magnetic resonance imaging techniques in neuromyelitis optica spectrum disorder. *JAMA Neurol* 2015;72:815–822.
- Melamed E, Levy M, Waters PJ, et al. Update on biomarkers in neuromyelitis optica. *Neurol Neuroimmunol Neuroinflamm* 2015;2:e134. doi:10.1212/NXI.0000000000000134.
- Chavarro VS, Mealy MA, Simpson A, et al. Insufficient treatment of severe depression in neuromyelitis optica spectrum disorder. *Neurol Neuroimmunol Neuroinflamm* 2016;3:e286. doi:10.1212/NXI.0000000000000286.
- Penner IK, Paul F. Fatigue as a symptom or comorbidity of neurological diseases. *Nat Rev Neurol* 2017;13:662–675.
- Zhao S, Mutch K, Elson L, Nurmikko T, Jacob A. Neuropathic pain in neuromyelitis optica affects activities of daily living and quality of life. *Mult Scler* 2014;20:1658–1661.

Neurology® Neuroimmunology & Neuroinflammation

Collaborative International Research in Clinical and Longitudinal Experience Study in NMOSD

Lawrence J. Cook, John W. Rose, Jessica S. Alvey, et al.
Neurol Neuroimmunol Neuroinflamm 2019;6;
DOI 10.1212/NXI.0000000000000583

This information is current as of June 28, 2019

Updated Information & Services	including high resolution figures, can be found at: http://nn.neurology.org/content/6/5/e583.full.html
References	This article cites 39 articles, 1 of which you can access for free at: http://nn.neurology.org/content/6/5/e583.full.html##ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): All Demyelinating disease (CNS) http://nn.neurology.org/cgi/collection/all_demyelinating_disease_cns Case control studies http://nn.neurology.org/cgi/collection/case_control_studies Optic neuritis; see Neuro-ophthalmology/Optic Nerve http://nn.neurology.org/cgi/collection/optic_neuritis Transverse myelitis http://nn.neurology.org/cgi/collection/transverse_myelitis
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://nn.neurology.org/misc/about.xhtml#permissions
Reprints	Information about ordering reprints can be found online: http://nn.neurology.org/misc/addir.xhtml#reprintsus

Neurol Neuroimmunol Neuroinflamm is an official journal of the American Academy of Neurology. Published since April 2014, it is an open-access, online-only, continuous publication journal. Copyright Copyright © 2019 The Author(s). Published by Wolters Kluwer Health, Inc. on behalf of the American Academy of Neurology.. All rights reserved. Online ISSN: 2332-7812.

