



CORRECTION

Correction to: Tobramycin Clearance Is Best Described by Renal Function Estimates in Obese and Nonobese Individuals: Results of a Prospective Rich Sampling Pharmacokinetic Study

Cornelis Smit^{1,2} • Roeland E. Wasmann³ • Marinus J. Wiezer⁴ • Hendricus P. A. van Dongen⁵ • Johan W. Mouton⁶ • Roger J. M. Brüggemann³ • Catherijne A. J. Knibbe^{1,2} 

© Springer Science+Business Media, LLC, part of Springer Nature 2020

Correction to: Pharm Res (2019) 36: 112
<https://doi.org/10.1007/s11095-019-2651-2>

There was a mistake in the units of CL and Q_v and in the parentheses of the formula for CL in the final model in Table 2. The corrected Table appears below.

The online version of the original article can be found at <https://doi.org/10.1007/s11095-019-2651-2>

✉ Catherijne A. J. Knibbe
c.knibbe@antoniusziekenhuis.nl

¹ Department of Clinical Pharmacy, St. Antonius Hospital, Koekoekslaan 1, 3435, CM Nieuwegein, The Netherlands

² Department of Systems Biomedicine and Pharmacology, Leiden Academic Centre for Drug Research Leiden University, Leiden, The Netherlands

³ Department of Pharmacy, Radboud Institute for Health Sciences, Radboudumc, Nijmegen, The Netherlands

⁴ Department of Surgery, St. Antonius Hospital, Nieuwegein, The Netherlands

⁵ Department of Anesthesiology, St. Antonius Hospital, Nieuwegein, The Netherlands

⁶ Department of Medical Microbiology and Infectious Diseases, Erasmus MC, Rotterdam, The Netherlands

Table 2 Population pharmacokinetic parameters of the base and final tobramycin model and results of the bootstrap analysis

	Base model (%RSE)		Final model (%RSE)		Bootstrap final model ($n = 1000$)		
					(95% Confidence interval)		
					Mean	Lower	Upper
V_c (L)	17.2	(7.3)	—				
$V_c = V_{c70\text{kg}} * (TBW/70)$							
$V_{c70\text{kg}}$ (L)	—		10.6	(11)	10.6	8.94	12.4
CL (L/h)	6.42	(4.3)	—				
$CL = CL_{MDRD\ 115} * (1 + Z * (MDRD - 115))$							
$CL_{MDRD\ 115}$ (L/h)	—		6.33	(2.6)	6.33	6.02	6.63
Z	—		0.00990	(10)	0.0100	0.0880	0.0122
V_p (L)	4.24	(15)	4.34	(18)	4.41	2.84	5.98
Q (L/h)	6.4	(5.1)	6.69	(12)	6.77	2.63	10.91
Inter-individual variability (IIV, %)							
V_c	42.9	(9.3)	24.9 ^a	(17)	24.1	14.9	30.8
CL	25.2	(14)	12.0 ^a	(13)	11.7	7.90	14.5
Residual variability							
Proportional error	0.112	(12)	0.116	(12)	0.115	0.0880	0.141
Additive error (mg/L)	0.369	(13)	0.346	(12)	0.342	0.239	0.445
OFV	351.7		289.6		276.6	185.9	367.2

^a η -shrinkage in the final model is 8% for IIV on CL and 6% for IIV on V_c . CL Clearance from the central compartment, $CL_{MDRD\ 115}$ Clearance from the central compartment for a person with a MDRD of 115 ml/min, MDRD De-indexed Modification of Diet in Renal Disease (in ml/min), OFV Objective Function Value, Q intercompartmental clearance between V_c and V_p , RSE Relative standard error, TBW Total body weight in kg, V_c Central volume of distribution, $V_{c70\text{kg}}$ Central volume of distribution for a 70 kg person