

# Body composition is not related to structural or vascular brain changes

Pauline H. Croll, Daniel Bos, M. Arfan Ikram, Fernando Rivadeneira, Trudy Voortman, Meike W. Vernooij

*Frontiers in Neurology, 2019*

## ABSTRACT

### Background

It is known that obesity [measured with body mass index (BMI)] relates to brain structure and markers of cerebral small vessel disease (CSVD). However, BMI may not adequately represent body composition. Furthermore, whether those cross-sectional associations hold longitudinally remains uncertain.

### Methods

Three thousand six hundred and forty-eight participants underwent baseline (2006-2014) dual-energy X-ray absorptiometry (DXA)-scan to obtain detailed measures of body composition and a magnetic resonance imaging (MRI) scan to assess brain structure. One thousand eight hundred and forty-four participants underwent a second MRI-scan at follow-up (2010-2017; median follow-up: 5.5 years). To assess cross-sectional and longitudinal associations (measures of change have been calculated) between body composition [BMI, fat mass index (FMI), fat-free mass index (FFMI)], and brain tissue volume (grey matter, white matter, hippocampus), white matter microstructure [fractional anisotropy (FA), mean diffusivity (MD)], and CSVD markers (white matter hyperintensity volume, lacunes, microbleeds) we used multivariable linear and logistic regression models.

### Results

A higher BMI and FMI were cross-sectionally associated with smaller white matter volumes, (difference in Z-score per SD higher BMI: -0.064 [95% CI: -0.094, -0.035]) and FMI: -0.067 [95% CI: -0.099, -0.034], higher FA and MD. A higher FFMI was associated larger grey matter volume (difference: 0.060 [95% CI: 0.018, 0.101]). There was no statistically significant or clinically relevant association between body composition and brain changes.

### Conclusions

Body composition, distinguishing between fat mass and fat-free mass, does not directly influence changes in brain tissue volume, white matter integrity and markers of CSVD. Cross-sectional associations between body composition and brain tissue volume likely reflect cumulative risk or shared etiology.

## INTRODUCTION

The increasing prevalence of obesity is accompanied by numerous adverse health effects, including cognitive decline and dementia.<sup>1,2</sup> Moreover, obesity has been linked with smaller brain tissue volumes, and decreased white matter integrity.<sup>1,3-5</sup> On top of that, it has been found that obesity is associated with focal brain pathology in the form of cerebral small vessel disease (CSVD).<sup>6</sup>

An important limitation of previous studies is that obesity is generally assessed by body mass index (BMI).<sup>1</sup> However, ageing is associated with a decrease in lean mass and an increase in fat mass, making BMI less suitable as an approximation of obesity in the elderly.<sup>1,7</sup> In a study looking at the association between body composition and metabolic syndrome the same “BMI problem” was encountered. Researchers found that fat mass index (FMI) was independently and positively associated with the presence of metabolic syndrome regardless of BMI and concluded that FMI is a better approximation of body composition.<sup>8</sup> A second limitation is that previous studies mainly assessed cross-sectional relations, precluding inference on directionality.

We studied the association of body composition, divided into body mass, fat mass and fat-free mass, with brain tissue volume, white matter integrity and markers of CSVD (white matter lesion hyperintensities (WMH), lacunes, and microbleeds) to evaluate whether fat mass and fat-free mass is a better approximation of body composition. Moreover, we aimed to assess the longitudinal association between body composition and brain health.

## MATERIALS AND METHODS

### Setting and study population

This study was embedded within the population-based Rotterdam Study. At study entry and subsequently every 3-4 years, all participants were invited to undergo extensive examinations. For the present study, 4,104 participants (2006-2014) underwent body composition assessment with a dual-energy X-ray absorptiometry (DXA) scan and a brain magnetic resonance imaging (MRI) scan.<sup>9</sup> From this group we excluded participants with cortical infarcts on MRI ( $N = 164$ ), cancer ( $N = 263$ ) and dementia ( $N = 29$ ), leaving 3,648 participants for the cross-sectional analyses. Between 2010 and 2017 in a subsequent examination-round, 1,844 participants had a follow-up MRI-scan available for the longitudinal analysis. Reasons of dropout at follow-up or unavailability of follow up MRI information have been included in Supplemental figure 1.

The Rotterdam Study has received medical ethical approval according to the Population Screening Act: Rotterdam Study, executed by the Ministry of Health, Welfare and Sports of the Netherlands. All participants provided written informed consent.

### **Body composition**

Body weight and length were measured and BMI was calculated ( $\text{kg}/\text{m}^2$ ).<sup>9</sup> A DXA – scan (iDXA, GE Lunar Healthcare, USA) was performed to determine fat-free mass and fat mass in kilograms.<sup>9</sup> From this, we calculated fat mass index (FMI) ( $\text{kg}/\text{m}^2$ ), and fat-free mass index (FFMI) ( $\text{kg}/\text{m}^2$ ).

### **Brain tissue volume, white matter integrity, and markers of CSVD**

Brain-MRI was performed on a 1.5-tesla MRI scanner with a 8-channel head coil, using a standardized scan protocol which has been described in detail before.<sup>9,10</sup> To quantify brain tissue volume, grey matter volume, white matter volume, WMH volume and intracranial volume, automated brain tissue classification was used. This quantification strategy was based on a *k*-nearest neighbour classifier algorithm, extended with an in-house-developed WMH segmentation.<sup>10</sup> Total brain volume did not include the cerebellum (used MRI segmentation only segments supratentorial brain tissue and not all MRI scans incorporated the entire cerebellum in the field of view). T1-weighted MR images were processed using FreeSurfer (version 5.1) to obtain the hippocampus volume.<sup>9</sup> White matter microstructure [fractional anisotropy (FA), mean diffusivity (MD)] was assessed with diffusion weighted imaging.<sup>9,11</sup> Visual ratings were performed for presence of lacunes or microbleeds.<sup>9</sup>

### **Covariables**

Information on energy intake, smoking, alcohol, education and physical activity was obtained from questionnaires and interviews as described in detail in the supplement.<sup>9</sup>

### **Statistics**

We investigated associations of body composition with brain tissue volumes, WMH volume (log-transformed), and FA and MD as a ratio of intracranial volume (to adjust for absolute head size differences) using linear regression models. We calculated changes by subtracting baseline measurement from follow-up measurement, which were eventually standardized and used to assess the longitudinal association with linear regression models. The association of body composition with presence of lacunes and microbleeds at baseline and the progression of lacunes or microbleeds at follow-up for the longitudinal analysis was assessed with logistic regression models. Adjustments were made for age, age<sup>2</sup>, sex, and education (model 1). We additionally adjusted for energy intake, smoking, physical activity, and alcohol (model 2). Adding height, hyper-

**Table 1.** Population characteristics

Characteristics	Population for cross-sectional analysis	Population for longitudinal analysis
	Baseline N = 3,648	Baseline N = 1,844
Age, years	65.9 (11.1)	60.9 (9.9)
Age range, years	45.7 – 97.8	44.1 – 90.2
Female, %	57.3	55.9
Education level, %		
Primary	8.1	7.1
Lower	38.5	35.4
Middle	30.1	29.3
Higher	22.4	27.9
Hypertension, %	44.1	34.3
Hypercholesterolemia, %	50.1	46.9
Diabetes, %	8.4	7.4
Smoking, %		
Never	33.0	33.3
Former	52.1	49.7
Current	14.4	16.4
Physical activity, MET-hours per week	44.7 (18.0-83.8) <sup>a</sup>	46.3 (19.6 – 83.8) <sup>a</sup>
Energy intake, kcal/day	2,090.5 (1,689.6 - 2,571.6) <sup>a</sup>	2,2567.0 (1,777.8 – 2,647.2) <sup>a</sup>
Alcohol intake, g/day	7.3 (1.1 - 17.6) <sup>a</sup>	8.5 (1.7 – 20.0) <sup>a</sup>
<b>Body composition</b>		
Length, cm	168.4 (9.4)	169.5 (9.4)
Weight, kg	77.4 (13.6)	77.9 (13.6)
Body mass index, kg/m <sup>2</sup>	27.2 (3.9)	27.0 (3.8)
Fat mass index, kg/m <sup>2</sup>	9.6 (3.3)	9.3 (3.3)
Fat-free mass index, kg/m <sup>2</sup>	17.6 (2.1)	17.4 (2.2)
<b>Brain MRI volumetry</b>		
Brain tissue volume, mL	930.2 (103.3)	949.4 (101.8)
Grey matter volume, mL	528.3 (60.8)	536.2 (61.9)
White matter volume, mL	401.8 (64.9)	413.2 (62.9)
Fractional anisotropy	0.3 (0.0)	0.3 (0.0)
Mean diffusivity	0.7 (0.0)	0.7 (0.0)
<b>Markers of cerebral small vessel disease</b>		
Lacunae, presence %	7.2	4.7
Microbleeds, presence %	19.7	16.1
White hyperintensity volume	8.2 (1.1)*	7.9 (1.0)*

MET: metabolic equivalent of task. kcal: kilocalories. g: gram. cm: centimetre. kg: kilogram. m: meter. mL: millilitre. Values are mean (standard deviation) for continuous variables or median (interquartile range) when indicated (<sup>a</sup>), percentages for dichotomous variables. \*; ln-transformed.

**Table 2.** The associations between body composition and brain tissue volume and CSVD markers – model 1

	Brain tissue volume	Grey matter volume	White matter volume	Hippocampus volume	Fractional anisotropy	Mean diffusivity	White matter hyperintensity volume*	Lacunes	Microbleeds
	Difference in Z-score (95% CI) p-value	Odds ratio (95% CI) p-value	Odds ratio (95% CI) p-value						
<b>Cross-sectional</b>									
<b>Body mass index (kg/m<sup>2</sup>)</b>	-0.025 (-0.050, 0.001) 0.056	0.042 (0.010, 0.073) 0.009	-0.064 (-0.094, -0.035) 0.000	-0.022 (-0.050, 0.006) 0.128	0.061 (0.031, 0.091) 0.000	0.045 (0.018, 0.072) 0.001	0.008 (-0.018, 0.034) 0.565	1.075 (0.944, 1.223) 0.276	0.931 (0.854, 1.015) 0.103
<b>Fat mass index (kg/m<sup>2</sup>)</b>	-0.033 (-0.061, -0.005) 0.020	0.034 (-0.001, 0.068) 0.055	-0.067 (-0.099, -0.034) 0.000	-0.019 (-0.015, 0.012) 0.224	0.098 (0.065, 0.130) 0.000	0.075 (0.045, 0.104) 0.000	0.000 (-0.028, 0.029) 0.982	1.080 (0.933, 1.250) 0.301	0.945 (0.859, 1.040) 0.250
<b>Fat-free mass index (kg/m<sup>2</sup>)</b>	-0.007 (-0.040, 0.027) 0.697	0.060 (0.018, 0.101) 0.005	-0.059 (-0.098, -0.021) 0.003	-0.028 (-0.065, 0.0094) 0.137	-0.022 (-0.062, 0.017) 0.268	-0.021 (-0.056, 0.014) 0.246	0.024 (-0.010, 0.058) 0.171	1.070 (0.905, 1.264) 0.430	0.899 (0.804, 1.005) 0.061
<b>Longitudinal</b>									
<b>Body mass index (kg/m<sup>2</sup>)</b>	0.004 (-0.029, 0.036) 0.810	-0.006 (-0.048, 0.036) 0.789	0.011 (-0.031, 0.054) 0.604	-0.050 (-0.097, -0.004) 0.034	0.029 (-0.031, 0.089) 0.339	0.052 (-0.007, 0.111) 0.084	-0.004 (-0.046, 0.038) 0.861	1.003 (0.816, 1.232) 0.979	0.978 (0.852, 1.123) 0.754
<b>Fat mass index (kg/m<sup>2</sup>)</b>	0.012 (-0.022, 0.047) 0.485	0.004 (-0.040, 0.049) 0.846	0.011 (-0.035, 0.056) 0.645	-0.033 (-0.082, 0.017) 0.192	0.048 (-0.014, 0.110) 0.130	0.061 (0.000, 0.122) 0.050	-0.009 (-0.054, 0.036) 0.686	0.906 (0.732, 1.120) 0.362	0.921 (0.800, 1.061) 0.257
<b>Fat-free mass index (kg/m<sup>2</sup>)</b>	-0.017 (-0.061, 0.027) 0.448	-0.031 (-0.087, 0.026) 0.291	0.012 (-0.046, 0.069) 0.684	-0.089 (-0.151, -0.026) 0.006	-0.026 (-0.108, 0.055) 0.527	0.017 (-0.063, 0.098) 0.672	0.010 (-0.047, 0.067) 0.726	1.149 (0.946, 1.397) 0.162	1.080 (0.945, 1.233) 0.259

\* In-transformed. CI: confidence interval. Cross-sectional differences represent the difference in standard deviation brain tissue volume (total brain tissue, grey matter, white matter, hippocampus volume, fractional anisotropy, mean diffusivity and white matter hyperintensity volume) as a ratio of intracranial volume per standard deviation higher body mass index, fat mass index, or fat-free mass index. Cross-sectional odds ratios represent the odds of the presence of lacunes and microbleeds per standard deviation higher body mass index, fat mass index, or fat-free mass index. Longitudinal differences represent the difference in standard deviation change in brain tissue volume (total brain tissue, grey matter, white matter, hippocampus volume, fractional anisotropy, mean diffusivity and white matter hyperintensity volume) as a ratio of intracranial volume per standard deviation higher body mass index, fat mass index, or fat-free mass index. Longitudinal odds ratios represent the odds of progressing from no presence of lacunes and microbleeds to presence or the odds of progressing to more lacunes or microbleeds at follow-up per standard deviation higher body mass index, fat mass index, or fat-free mass index. Effect estimates were considered statistically significant with a p-value < 0.05. Cross sectional models are adjusted for age, age<sup>2</sup>, sex, and education. Longitudinal models were additionally adjusted for time between MRI scans.

**Table 3.** The associations between body composition and brain tissue volume and CSVD markers – model 2

	Brain tissue volume	Grey matter volume	White matter volume	Hippocampus volume	Fractional anisotropy	Mean diffusivity	White matter hyperintensity volume*	Lacunes	Microbleeds
	Difference in Z-score (95% CI) p-value	Odds ratio (95% CI) p-value	Odds ratio (95% CI) p-value						
<b>Cross-sectional</b>									
<b>Body mass index (kg/m<sup>2</sup>)</b>	-0.028 (-0.060, 0.003) 0.075	0.039 (0.001, 0.076) 0.043	-0.066 (-0.101, -0.030) 0.000	-0.016 (-0.049, 0.018) 0.357	0.078 (0.043, 0.112) 0.000	0.060 (0.029, 0.091) 0.000	0.013 (-0.018, 0.043) 0.409	1.134 (0.980, 1.131) 0.091	0.918 (0.830, 1.015) 0.095
<b>Fat mass index (kg/m<sup>2</sup>)</b>	-0.035 (-0.070, -0.001) 0.045	0.033 (-0.008, 0.075) 0.113	-0.069 (-0.108, -0.030) 0.001	-0.008 (-0.044, 0.029) 0.685	0.114 (0.076, 0.152) 0.000	0.092 (0.058, 0.127) 0.000	0.006 (-0.028, 0.039) 0.728	1.137 (0.962, 1.344) 0.131	0.927 (0.829, 1.038) 0.188
<b>Fat-free mass index (kg/m<sup>2</sup>)</b>	-0.014 (-0.056, 0.027) 0.498	0.051 (0.002, 0.101) 0.041	-0.061 (-0.107, -0.014) 0.010	-0.034 (-0.077, 0.010) 0.130	-0.002 (-0.047, 0.044) 0.944	-0.010 (-0.051, 0.032) 0.646	0.028 (-0.012, 0.068) 0.165	1.149 (0.948, 1.392) 0.156	0.895 (0.786, 1.018) 0.092
<b>Longitudinal</b>									
<b>Body mass index (kg/m<sup>2</sup>)</b>	0.029 (-0.012, 0.069) 0.165	0.008 (-0.041, 0.057) 0.756	0.028 (-0.021, 0.077) 0.270	-0.041 (-0.098, 0.016) 0.163	0.004 (-0.070, 0.078) 0.918	0.025 (-0.043, 0.092) 0.475	0.005 (-0.045, 0.055) 0.840	0.943 (0.725, 1.227) 0.662	0.979 (0.831, 1.154) 0.804
<b>Fat mass index (kg/m<sup>2</sup>)</b>	0.039 (-0.005, 0.083) 0.080	0.005 (-0.011, 0.021) 0.567	0.032 (-0.021, 0.085) 0.233	-0.019 (-0.080, 0.042) 0.546	0.014 (-0.063, 0.091) 0.728	0.021 (-0.049, 0.091) 0.557	-0.004 (-0.057, 0.0105) 0.897	0.835 (0.632, 1.104) 0.206	0.948 (0.801, 1.122) 0.534
<b>Fat-free mass index (kg/m<sup>2</sup>)</b>	0.003 (-0.052, 0.058) 0.920	-0.011 (-0.078, 0.055) 0.741	0.016 (-0.051, 0.083) 0.643	-0.093 (-0.171, -0.015) 0.019	-0.023 (-0.126, 0.079) 0.655	0.030 (-0.063, 0.124) 0.521	0.027 (-0.041, 0.095) 0.444	1.160 (0.896, 1.501) 0.260	1.041 (0.885, 1.225) 0.629

\* In-transformed. CI: confidence interval. kg: kilogram. m: meter. Cross-sectional differences represent the difference in standard deviation brain tissue volume (total brain tissue, grey matter, white matter, and hippocampus volume) and the difference in standard deviation fractional anisotropy, mean diffusivity and white matter hyperintensity volume as a ratio of intracranial volume per standard deviation higher body mass index, fat mass index, or fat-free mass index. Cross-sectional odds ratios represent the odds of the presence of lacunes and microbleeds per standard deviation higher body mass index, fat mass index, or fat-free mass index. Longitudinal differences represent the difference in standard deviation change in brain tissue volume (total brain tissue, grey matter, white matter, and hippocampus volume) and the difference in standard deviation fractional anisotropy, mean diffusivity and white matter hyperintensity volume as a ratio of intracranial volume per standard deviation higher body mass index, fat mass index, or fat-free mass index. Longitudinal odds ratios represent the odds of progressing from no presence of lacunes and microbleeds to presence or the odds of progressing to more lacunes or microbleeds at follow-up per standard deviation higher body mass index, fat mass index, or fat-free mass index. Effect estimates were considered statistically significant with a p-value < 0.05. Cross sectional models are adjusted for age, sex, education, energy intake, smoking, physical activity, and alcohol consumption. Longitudinal models were additionally adjusted for time between MRI scans.

tension, hypercholesterolemia, and diabetes to the regression models did not change the effect estimates and were therefore left out of the final analysis. For the longitudinal analysis we additionally adjusted for time between MRI-scans. We ran the same analyses, with brain tissue volumes, WMH volume, and FA and MD unstandardized (while adjusting for intracranial volume in the regression model) to evaluate clinical relevance of the associations. We compared our results with previously reported brain tissue changes (independent from inter-individual differences in head size) ranging from 3.6 millilitres to 5.4 millilitres with one year of ageing.<sup>12,13</sup> We checked for interaction between body composition and sex and between body composition and age (< 60 versus ≥ 60 years) as effects of BMI might differ between mid- and late-life.<sup>7</sup> We were powered (alpha level: 0.05; power level: 0.80) to detect minimal differences of 2.028 millilitre brain tissue volume. A p-value < 0.05 was considered statistically significant. IBM SPSS statistics version 24.0 (IBM Corp, Armonk, NY, USA) was used for analyses.

## RESULTS

Baseline characteristics are described in Table 1. Mean age was 65.9 years (standard deviation (SD): 9.8), 57.3% was female. Mean BMI was 27.2 kg/m<sup>2</sup> (SD: 3.9). Participants with a follow-up MRI scan compared to those with only baseline MRI were younger, higher educated, and healthier (Supplementary table 1).

We found that cross-sectionally a higher BMI and FMI were related with smaller white matter volumes (difference in Z-score per SD higher BMI: -0.064 [95% CI: -0.094, -0.035]; FMI: -0.067 [95% CI: -0.099, -0.034]), higher FA (BMI difference: 0.061 [95% CI: 0.031, 0.091]; FMI difference: 0.098 [95% CI: 0.065, 0.130]) and higher MD (BMI difference: 0.045 [95% CI: 0.018, 0.072]; FMI difference: 0.075 [95% CI: 0.045, 0.104]), and a higher FFMI was related with larger grey matter volumes (difference: 0.060 [95% CI: 0.018, 0.101]) (Table 2, model 1). Effect estimates did not change between the different models (Table 3, model 2), nor were there differences in effects of body composition indices between the various outcomes of interest. Though statistically significant, cross-sectional results might not be clinically relevant as effect estimates of change in brain tissue volumes, as well as their corresponding confidence intervals, reported in millilitre difference per one point increase in BMI, FMI, or FFMI are small and do not exceed clinically relevant amounts of tissue changes (Supplementary tables 2 and 3).

Only the longitudinal association between FFMI and change in hippocampus volume was statistically significant (difference: -0.089 [95% CI: -0.151, -0.026]) (Table 2; model 1), and did not change across models. However, it is questionable whether the direction of this association is biological plausible, and when reporting this association in millilitre (Supplementary table 3; model 2), it is borderline statistically significant (difference in

millilitre per one point increase in FFMI: -0.027 [95% CI: -0.055, 0.000]). We did not find any statistically significant or clinically relevant longitudinal associations between body composition, brain tissue volumes, white matter integrity, or markers of CSVD (Tables 2 and 3; Supplementary tables 2 and 3). Both cross-sectional and longitudinal results did not differ between midlife and late life (data not shown), or between males and females (Supplementary table 4).

## DISCUSSION

We found that detailed measures of body composition (higher BMI and higher FMI) related to smaller white matter volume and decreased white matter microstructure and that higher FFMI related to higher grey matter volumes. However, expressing those cross-sectional associations in millilitre difference, raises the question whether those are also clinically relevant. No statistically significant or clinically relevant associations were found longitudinally (median follow-up: 5.5 years).

Strengths of this study are the large sample size, the distinction of body composition, the wide range of brain imaging markers and the longitudinal design. However, some limitations should also be acknowledged. First, due to the observational design we cannot exclude the possibility of residual confounding. Second, we did not adjust for possible changes in body composition over time in the longitudinal analysis.

Previous studies showed that body composition relates to brain structure,<sup>1,2</sup> which is replicated in some of our cross-sectional results. Currently there are no established thresholds for rates of brain atrophy. However, in two other population-based studies, change in brain tissue volume with one year of ageing were found to range from 3.6 millilitres to 5.4 milliliters.<sup>3,4</sup> Against this background, it is questionable whether a (cross-sectional) 0.496 millilitre smaller brain volume per one-point increase in BMI (independent of age) and other (longitudinal) effect estimates, are also clinically relevant. The absence of a longitudinal association did not result from insufficient power, as we were able to minimally detect a change of 2.028 millilitre brain tissue volume, which is smaller than a clinically relevant change of 5 millilitre brain volume.<sup>3</sup> Though, the absence of a longitudinal association might result in part from selection bias. Indeed, it appeared that participants with MRI at follow-up compared to those without follow-up were younger, higher educated and healthier. Another explanation might be that the effects of body composition in midlife and late life may be reversed.<sup>5,6</sup> But, we could not confirm effect modification by age in our study. Other longitudinal studies are inconclusive in their results, both reporting non-significant associations, and significant associations between body composition and various brain measurements,<sup>6-10</sup> probably due to different methodologies. In a study investigating whether the effect of obesity

on the brain increased with age they found no significant results and thus suggested that obesity may act as a modifier of brain atrophy rather than it being a direct cause.<sup>11</sup> Cardiovascular risk factors (hypertension, glucose, cholesterol, plasma triglyceride) are known to co-occur frequently with obesity and have independent negative effects on brain structure.<sup>12-15</sup> Therefore, although we did not find significant associations between body composition and markers of CSVD, it might be that the effects of vascular risk factors may be less pronounced in their effects on brain structure. And it might also be that solely body composition does not influence brain health, but it is rather the interplay of cardiovascular risk factors altogether that may impact brain health negatively.<sup>16, 17</sup>

Since BMI may not be a good approximation for body composition,<sup>18</sup> we used FMI and FFMI. Surprisingly, both cross-sectionally and longitudinally, regression estimates between the different body composition measurement (BMI, FMI and FFMI) and brain MRI data did not differ.

Based on our findings it seems that cross-sectional associations between body composition and brain health likely reflect cumulative risk or shared etiology. Moreover, it seems that body composition, decomposed into fat mass and fat-free mass, does not influence brain changes or the presence of CSVD. Therefore, we cannot conclude that FMI and FFMI is a better approximation of adiposity than BMI to study brain health. Other longitudinal studies are needed to replicate our findings.

## REFERENCES

1. Stillman CM, Weinstein AM, Marsland AL, Gianaros PJ, Erickson KI. Body-Brain Connections: The Effects of Obesity and Behavioral Interventions on Neurocognitive Aging. *Front Aging Neurosci* 2017;9:115.
2. Collaborators GBDO. Health Effects of Overweight and Obesity in 195 Countries over 25 Years. *N Engl J Med* 2017.
3. Kullmann S, Schweizer F, Veit R, Fritsche A, Preissl H. Compromised white matter integrity in obesity. *Obes Rev* 2015;16:273-281.
4. Cherbuin N, Sargent-Cox K, Fraser M, Sachdev P, Anstey KJ. Being overweight is associated with hippocampal atrophy: the PATH Through Life Study. *Int J Obes (Lond)* 2015;39:1509-1514.
5. Dekkers IA, Jansen PR, Lamb HJ. Obesity, Brain Volume, and White Matter Microstructure at MRI: A Cross-sectional UK Biobank Study. *Radiology* 2019;181012.
6. Karcher HS, Holzwarth R, Mueller HP, et al. Body fat distribution as a risk factor for cerebrovascular disease: an MRI-based body fat quantification study. *Cerebrovasc Dis* 2013;35:341-348.
7. Spauwen PJ, Murphy RA, Jonsson PV, et al. Associations of fat and muscle tissue with cognitive status in older adults: the AGES-Reykjavik Study. *Age Ageing* 2017;46:250-257.
8. Liu P, Ma F, Lou H, Liu Y. The utility of fat mass index vs. body mass index and percentage of body fat in the screening of metabolic syndrome. *BMC Public Health* 2013;13:629.
9. Ikram MA, Brusselle GGO, Murad SD, et al. The Rotterdam Study: 2018 update on objectives, design and main results. *Eur J Epidemiol* 2017.
10. Ikram MA, van der Lugt A, Niessen WJ, et al. The Rotterdam Scan Study: design update 2016 and main findings. *Eur J Epidemiol* 2015;30:1299-1315.
11. de Groot M, Ikram MA, Akoudad S, et al. Tract-specific white matter degeneration in aging: the Rotterdam Study. *Alzheimers Dement* 2015;11:321-330.
12. Resnick SM, Pham DL, Kraut MA, Zonderman AB, Davatzikos C. Longitudinal magnetic resonance imaging studies of older adults: a shrinking brain. *J Neurosci* 2003;23:3295-3301.
13. Croll PH, Voortman T, Ikram MA, et al. Better diet quality relates to larger brain tissue volumes: The Rotterdam Study. *Neurology* 2018;90:e2166-e2173.
14. van Bloemendaal L, Ijzerman RG, Ten Kulve JS, et al. Alterations in white matter volume and integrity in obesity and type 2 diabetes. *Metab Brain Dis* 2016;31:621-629.
15. Ronan L, Alexander-Bloch AF, Wagstyl K, et al. Obesity associated with increased brain age from midlife. *Neurobiol Aging* 2016;47:63-70.
16. Shaw ME, Sachdev PS, Abhayaratna W, Anstey KJ, Cherbuin N. Body mass index is associated with cortical thinning with different patterns in mid- and late-life. *Int J Obes (Lond)* 2017.
17. Gustafson D, Lissner L, Bengtsson C, Bjorkelund C, Skoog I. A 24-year follow-up of body mass index and cerebral atrophy. *Neurology* 2004;63:1876-1881.
18. Brooks SJ, Benedict C, Burgos J, et al. Late-life obesity is associated with smaller global and regional gray matter volumes: a voxel-based morphometric study. *Int J Obes (Lond)* 2013;37:230-236.
19. Driscoll I, Beydoun MA, An Y, et al. Midlife obesity and trajectories of brain volume changes in older adults. *Hum Brain Mapp* 2012;33:2204-2210.
20. Yates KF, Sweat V, Yau PL, Turchiano MM, Convit A. Impact of metabolic syndrome on cognition and brain: a selected review of the literature. *Arterioscler Thromb Vasc Biol* 2012;32:2060-2067.
21. Sala M, de Roos A, van den Berg A, et al. Microstructural brain tissue damage in metabolic syndrome. *Diabetes Care* 2014;37:493-500.

22. Ikram MA, Vrooman HA, Vernooij MW, et al. Brain tissue volumes in the general elderly population. The Rotterdam Scan Study. *Neurobiol Aging* 2008;29:882-890.
23. DeBette S, Beiser A, Hoffmann U, et al. Visceral fat is associated with lower brain volume in healthy middle-aged adults. *Annals of neurology* 2010;68:136-144.
24. Pichler A, Khalil M, Langkammer C, et al. The impact of vascular risk factors on brain volume and lesion load in patients with early multiple sclerosis. *Mult Scler* 2017:1352458517736149.
25. Goyal A, Nimmakayala KR, Zonszein J. Is There a Paradox in Obesity? *Cardiology in review* 2014;22:163-170.