



EXPERT STATEMENT



Nutritional support for children during critical illness: European Society of Pediatric and Neonatal Intensive Care (ESPNIC) metabolism, endocrine and nutrition section position statement and clinical recommendations

Lyvonne N. Tume^{1,2*} , Frederic V. Valla³ , Koen Joosten⁴, Corinne Jotterand Chaparro^{5,6}, Lynne Latten⁷, Luise V. Marino⁸, Isobel Macleod⁹, Clémence Moullet^{5,6}, Nazima Pathan¹⁰, Shancy Rooze¹¹, Joost van Rosmalen¹² and Sascha C. A. T. Verbruggen⁴

© 2020 The Author(s)

Abstract

Background: Nutritional support is considered essential for the outcome of paediatric critical illness. There is a lack of methodologically sound trials to provide evidence-based guidelines leading to diverse practices in PICUs worldwide. Acknowledging these limitations, we aimed to summarize the available literature and provide practical guidance for the paediatric critical care clinicians around important clinical questions many of which are not covered by previous guidelines.

Objective: To provide an ESPNIC position statement and make clinical recommendations for the assessment and nutritional support in critically ill infants and children.

Design: The metabolism, endocrine and nutrition (MEN) section of the European Society of Pediatric and Neonatal Intensive Care (ESPNIC) generated 15 clinical questions regarding different aspects of nutrition in critically ill children. After a systematic literature search, the Scottish Intercollegiate Guidelines Network (SIGN) grading system was applied to assess the quality of the evidence, conducting meta-analyses where possible, to generate statements and clinical recommendations, which were then voted on electronically. Strong consensus (> 95% agreement) and consensus (> 75% agreement) on these statements and recommendations was measured through modified Delphi voting rounds.

Results: The final 15 clinical questions generated a total of 7261 abstracts, of which 142 publications were identified relevant to develop 32 recommendations. A strong consensus was reached in 21 (66%) and consensus was reached in 11 (34%) of the recommendations. Only 11 meta-analyses could be performed on 5 questions.

Conclusions: We present a position statement and clinical practice recommendations. The general level of evidence of the available literature was low. We have summarised this and provided a practical guidance for the paediatric critical care clinicians around important clinical questions.

Keywords: Enteral nutrition, Parenteral nutrition, Child, Paediatric, Intensive care, Guidelines

*Correspondence: lyvonnetume@gmail.com

¹ Faculty of Health and Society, University of Salford, Manchester M6 6PU, UK

Full author information is available at the end of the article

Lyvonne N. Tume and Frédéric V. Valla: joint first authors.

Introduction

Critical illness induces profound metabolic and endocrine changes in close interaction with the alterations in autonomic and immune systems. The metabolic and endocrine changes are characterized by catabolism, insulin resistance and shifts in substrate utilisation [1]. These changes evolve during the course of illness, where the acute changes are assumed to be advantageous for survival. However, following the acute phase these changes might become harmful [1,2]. Parallel to these changes, critically ill children frequently experience feeding difficulties, caused by (perceived) feed intolerance and feeding interruptions [3,4]. This often leads to undernourishment with a cumulative macronutrient deficit during the course of their Pediatric Intensive Care Unit (PICU) stay [5, 6]. Malnutrition at PICU admission is frequent (15–25% prevalence rates) in developing countries; nutritional status deterioration is also an early and frequent phenomenon in this setting with almost one-third of critically ill children presenting with nutritional indices decline [7–9]. Muscle wasting is also a constant, intense and rapid phenomenon [10]. Malnourishment and macronutrient deficits during critical illness have been associated with increased morbidity (infections, weakness, prolonged mechanical ventilation and delayed recovery) as well as increased mortality. However, overfeeding has also been shown to pose harm to critically ill children, especially during the acute phase. As the metabolic and endocrine response evolves during the course of critical illness, possibly the nutritional support should also accommodate these changes and differ during the different phases of paediatric critical illness as well.

Although optimal nutrition is considered essential to improve outcomes in critically ill children, large well-designed randomized controlled trials (RCTs) with clinically relevant outcome measures are scarce [11, 12]. The limited evidence has led to a wide variation in nutritional practices worldwide, between individual clinicians, PICUs and countries [13, 14]. Yet the evidence is increasing, and the number of publications on nutritional support in paediatric critical illness in 2018 has doubled when compared with 2012 and tripled since 2007. In 2017 the American Society of Parenteral and Enteral Nutrition (ASPEN) and the Society of Critical Care Medicine (SCCM) published their guidelines for the provision of nutritional support in the PICU [15]. However, several important clinical topics remained unanswered [16]. For instance, term neonates (defined as >37–44 weeks' gestational age) which comprise around 32% of the PICU population, were excluded from these recommendations [17]. As a multidisciplinary research group within Europe, the ESPNIC metabolism, endocrinology and

Take-home message

There is a lack of high-quality evidence to guide nutrition in paediatric critical illness. This position statement and clinical recommendations summarise the existing evidence around 15 of the most important clinical questions, and where no evidence is available, suggest good clinical practice.

nutrition (MEN) section, therefore, felt it was timely to address unanswered clinical questions and review new evidence to produce a position statement and recommendations on artificial nutrition in critically ill children.

Methodology

Selection of members

The working group was composed of a multidisciplinary team of 11 European specialists (five paediatric intensivists, two nurses and four dietitians) in nutritional support for critically ill children, who are members of the MEN section of the European Society of Pediatric and Neonatal Intensive Care (ESPNIC). Four members (LT/CJ/KJ/SV) were well trained and experienced in the development and methodology of systematic reviews and development of recommendations. A biostatistician (JvR) was added to the multi-disciplinary team specifically for the expertise in meta-analyses, but did not participate in development of the recommendations or the voting process.

Question development and search strategy

The working group met initially, in June 2017, to discuss the project, and generate 15 broad clinical questions. The systematic literature search was performed by biomedical information specialists (EK, SG, GdJ and WB; see acknowledgements) of the Erasmus Medical Centre Library (Rotterdam, The Netherlands) in four databases (Embase.com; Medline Epub (Ovid); Cochrane Central; Web of Science) and included all articles published from 1997 until May 2018 and updated in November 30 2018. Supplement file 1 describes the search terms used per question.

Inclusion and exclusion criteria were agreed by the group. Inclusion criteria were RCTs, case-control, before and after and cohort studies including critically ill term neonates and children (aged ≥ 37 weeks' gestational age–18 years). We only included manuscripts written in English or French, which excluded three papers, one in Russian two in Chinese. Publications describing studies in pre-term infants were excluded, unless the question specifically related to neonatal PICU patients and no evidence existed in term neonates (Question 4). In addition to reviews, animal studies, case reports, editorials, commentaries, conference abstracts and letters were

excluded. Separate publications presenting outcomes from the same study population were included, but seen as one study, and the study that provided the most complete data to answer the question was included. For each of the 15 questions, key search words were defined, and specific search combinations were developed for the four databases (Supplementary file 1).

Selection of studies

In order to select the eligible studies, the results from each database were combined and exported to Endnote, followed by removal of duplicates and exportation to a Word document, allowing at least two working group members to separately undertake the screening of the abstracts in a standardized way. Abstracts were screened for eligibility by the group members, and those which were thought to be eligible were automatically exported as final abstract. Areas of disagreement were resolved by discussion. Abstracts that determined to be eligible by one of the two members were discussed with a third reviewer before decision of inclusion or exclusion. If eligibility criteria were met, full manuscripts were procured. Similarly, if a disagreement on the eligibility of the paper occurred, a discussion took place with a third reviewer. We also examined reference lists from included articles for suitable studies. A PRISMA diagram is shown in Supplementary file 2.

Data extraction and assessment of study quality and evidence grading

Data from eligible papers were extracted by two reviewers with the primary reviewer not an author on the paper. In addition, the risk of bias was assessed by two reviewers independently using the SIGN critical appraisal checklists available for each study design (<https://www.sign.ac.uk/checklists-and-notes.html>) (Supplementary file 3). Any disagreement with grading was discussed and the two lead authors (LT/FV) reviewed all the evidence grading. The classification of the literature into levels of evidence was performed according to the SIGN grading system (Supplementary file 3).

Data analysis including meta-analyses

In some questions, the data were combined statistically in a meta-analysis if they met the following criteria: there was more than one study, the combined studies (in one analysis) were either randomised trials or observational studies, the population and the intervention were sufficiently similar to combine and the outcomes were the same, or for continuous outcome variables, if we had data on the distribution of the variable. To perform the

meta-analyses, we a priori defined clinically relevant outcome variables on which the meta-analyses would be performed. These were mortality, new infections, gastrointestinal complications (vomiting aspiration/diarrhoea/NEC-ischemia), length of ventilation and length of stay (PICU/hospital). Anticipating a broad inconsistency of these outcome variables we chose a pragmatic meta-analysis. The risk of bias tables are presented in Supplementary File 4. For dichotomous outcomes, we used a random effects model for the relative risk of the intervention to compute a pooled relative risk and its 95% CI. The Hartung–Knapp–Sidik–Jonkman method was used to estimate the between-study variance, and a continuity correction of 0.5 was applied in case of zero cell frequencies. The heterogeneity of combined study results was assessed using the inconsistency statistic and tested using Cochran's Q test. The meta-analyses were performed using R version 3.6.1 with the package meta.

Consensus methodology and grading of the recommendations

Based on the results from the systematic review and meta-analyses, a first draft of recommendations was composed, including the supporting text and grade of recommendation. The classification of the grades of recommendation (A–D, Good Clinical Practice) was undertaken according to the SIGN grading system (Supplementary file 3) [18]. In May 2018, a second meeting took place to discuss all questions and review the evidence quality and recommendations. The group generated the position statement and a draft guideline with a total of 32 recommendations, which was followed by a round of electronic voting to gain consensus using a Delphi method in June 2018 [19, 20]. The survey involved voting on each recommendation on 3-point scale with categories: disagree, agree and unsure. This was created and distributed via a proprietary electronic online platform hosted by the University of Southampton (<https://www.isurvey.soton.ac.uk/>) and checked by one of the authors (LM) without identifying features to ensure anonymity. In round 2, we provided the group results and asked the group to re-vote. We defined strong consensus as agreement of >95%, consensus as agreement of 75–95% and no consensus as agreement <75%. Feedback received during the first round of online voting was used to modify and improve the recommendations in order to reach a higher degree of consensus at the final online voting in September 2018. Any recommendations with an agreement equal to or lower than 95% were discussed at a consensus meeting which took place on 31 October 2018. Following a revised meta-analysis, a last and final meeting of a core group of four members took place in November 2019, which was followed by a final round of

electronic voting. The AGREE reporting checklist for guideline development was followed (Supplementary File 5). The ESPNIC process for endorsement of guidelines was undertaken.

Results

A total of 7261 abstracts were screened. Subsequently 142 publications were reviewed, and data were extracted (Supplementary file 2) and included in the development of 32 recommendations (Table 1). The general level of evidence was low: out of the 142 publications, 5 (3.5%) were graded 1+ according to the SIGN grading system, 27 (18.9%) were graded 1−, six (4.2%) were graded 2++, 20 (14.0%) were graded 2+, 82 (58.0%) were graded 2−, one (0.7%) was graded 3 and one (0.7%) was graded 4. Furthermore, the data were suitable for meta-analysis for only 11 (sub)questions, all of which had dichotomous outcome measures. All forest plots of the meta-analyses have been provided in supplement file 6. Overall, heterogeneity of the studies suitable for meta-analysis varied with I^2 0–91% (p value 0.13–0.83), and two meta-analyses with a I^2 of higher than 50%, 53% and 55%, respectively, for the impact of gastric versus post-pyloric feeding on aspiration and intermittent versus continuous feeding on diarrhoea. The pooled relative risk showed a significant difference between groups in only 1 out of these 11 meta-analyses. Enteral feeding versus no enteral feeding in children on haemodynamic support resulted in a lower risk of mortality (RR 0.41 [95%CI 0.20–0.86]). Accordingly, the grading of the 32 recommendations was as follows: five recommendations were graded as B, five as C, 12 as D and 10 were GCP.

A strong consensus was reached in 21 (66%) and consensus was reached in 11 (34%) of the recommendations. A detailed discussion of the clinical questions, the recommendations with evidence grading, and level of consensus achieved are presented in Supplementary file 7 with a full reference list. The table of evidence is presented in Supplementary file 8. A summary of all recommendations is shown in Table 1. A summary of comparisons between our recommendations and those presented by ASPEN/SCCM is shown in Table 2.

Discussion

This position statement with clinical recommendations provides new guidance based on new evidence, as well as reinforcing most of the existing 2017 ASPEN Guidelines. These ASPEN PICU nutrition guidelines published in 2017 were based on a literature search from January 1995 to March 2016 and consisted of 17 recommendations. These ESPNIC clinical recommendations are based on an updated literature search until November 2018. Both the American (ASPEN) and our European guidelines

provide expert opinion which is essential in this setting where limited data are available. Our recommendations are predominantly consistent with the ASPEN guideline recommendations (Table 2) which helps assist in the uptake and implementation of guidelines into practice [21]. Implementation of evidence into clinical practice remains problematic, in 2017 a European survey of 59 PICUs found that 69% of PICUs still had no local feeding guidelines [13]. Additionally, this position statement generated new clinical guidance as half of our clinical questions differed from the ASPEN guidelines. These included guidance on feeding neonates with arterial umbilical arterial catheters; the type of enteral formula to be used; the amount or type of each macronutrient to provide; the value of gastric residual volume to assess feeding tolerance; the use of prokinetics to enhance feeding tolerance and the use of feeding protocols to improve outcomes. Furthermore, these new ESPNIC recommendations covered in more detail the indications for enteral nutrition in various subgroups of patients in clinicians are in general uncertain on how to progress feeding (i.e. term neonates and children on haemodynamic support and after cardiac surgery) [13]. In addition, our position statement provides a different stand on two recommendations as compared with the ASPEN guidelines, based on new available research. In contrast with the ASPEN guidelines we recommend to consider withholding parenteral nutrition during the first week in neonates and children, independent of their nutritional state [8, 15, 22]. Furthermore, there was also strong consensus in our working group that there is insufficient evidence to recommend a protein/amino acid intake of 1.5 g/kg/day or higher during the acute phase of disease to benefit clinical outcomes [15]. The intake of 1.5 g/kg/day or higher has shown to prevent cumulative negative protein balance [23, 24]. However, future research should consider that the exact threshold is unknown and might overestimate protein/amino acid requirements during acute critical illness; thus further work should, therefore, also investigate low protein/amino acid intakes during that phase [25].

Overall, as expected, the general level of evidence was low, and the meta-analyses provided little value because of the heterogeneity in interventions and outcomes, population and the type of study designs (few RCTs). This resulted in few studies able to be pooled for this analysis. Despite these limitations, we formulated 32 recommendations which can guide PICU healthcare professionals. There are a few key messages to be taken from our position statement. Although hardly any methodologically sound studies exist, recent developments have shown that nutritional interventions in our PICUs are capable of impacting on the short- and long-term

Table 1 Summary of recommendations for nutritional support for children during critical illness

Question	Recommendation	SIGN recommendation grade	Consensus	References used in synthesis of recommendations
In critically ill children, should nutritional status be assessed and what is the optimal method to assess nutritional status?	1.1: The assessment of nutritional status is recommended in critically ill children at admission and throughout their PICU admission	GCP	Strong consensus	[31–57]
	1.2: It is recommended to perform anthropometric measurements on admission and regularly during admission, and to express these measurements in z-scores, including weight, height/length mid upper arm circumference and head circumference in young children	GCP	Strong consensus	[6, 10, 48–55]
In critically ill children, when should enteral nutrition be commenced and how should it be increased?	2.1: It is recommended to commence early enteral nutrition within 24 h of admission unless contraindicated	D	Strong consensus	[58, 61, 64–69, 71]
	2.2: It is recommended to increase enteral nutrition in a stepwise fashion until goal for delivery is achieved using a feeding protocol or guideline	D	Strong consensus	[59, 60, 62, 63, 71, 72, 74–76]
In critically ill children on haemodynamic support (vasoactive medications, extra-corporeal life support ECLS) does enteral feeding compared to no enteral feeding affect outcomes?	3.1: Early enteral nutrition is recommended in term neonates who are stable on ECLS	D	Consensus	[77–80]
	3.2: Early enteral nutrition is recommended in children who are stable on ECLS	D	Strong consensus	[82]
	3.3: Early enteral nutrition is recommended in term neonates who are stable on pharmaceutical haemodynamic support	GCP	Consensus	[83, 85, 86]
	3.4: Early enteral nutrition is recommended in children who are stable on pharmaceutical haemodynamic support	D	Strong consensus	[83, 85, 86]
	3.5: Early enteral nutrition is recommended in children after cardiac surgery	C	Consensus	[87–94]
In critically ill term neonates with umbilical arterial catheters and/or PGE1 infusions, does enteral feeding impact on adverse events?	4.1: Enteral nutrition should be considered in term neonates with umbilical arterial catheters	D	Strong consensus	[95, 96, 100]
	4.2: Enteral nutrition should be considered in critically ill term neonates on PGE1 infusion if managed in a critical care unit with adequate observation and monitoring	D	Strong consensus	[97–99]
In critically ill children what are their energy requirements?	5.1 In the acute phase, energy intake provided to critically ill children should not exceed resting energy expenditure	C	Strong consensus	[26, 101–104]
	5.2. After the acute phase, energy intake provided to critically ill children should account for energy debt, physical activity, rehabilitation and growth	GCP	Strong consensus	[105–113]
In critically ill children, what is the most accurate method of determining or predicting energy expenditure?	6.1 Measuring resting energy expenditure using a validated indirect calorimeter should be considered to guide nutritional support in critically ill infants and children after the acute phase	GCP	Strong consensus	[114–119]
	6.2 Schofield equation (for age and gender and using an accurate weight) is recommended to estimate resting energy expenditure	C	Strong consensus	[120–125]
In critically ill children, what are the macro-nutrient requirements?				
What is the recommended glucose intake?	7.1. Parenteral glucose provision should be sufficient to avoid hypoglycemia but not excessive to prevent hyperglycemia	D	Strong consensus	[126, 127]

Table 1 (continued)

Question	Recommendation	SIGN recommendation grade	Consensus	References used in synthesis of recommendations
What is the recommended lipid intake or type?	7.2: When parenteral nutrition is used, composite lipid emulsions, with or without fish oil, should be considered as the first-choice treatment	GCP	Strong consensus	[128]
What is the recommended protein/amino acid intake?	7.3a: For critically ill infants and children on enteral nutrition a minimum enteral protein intake of 1.5 g/kg/d can be considered to avoid negative protein balance	B	Strong consensus	[23, 24, 106–108, 129, 130]
	7.3b: There is insufficient evidence available to support the use of additional protein/amino acid intake during the acute phase of illness (Strong consensus)	D	Strong consensus	[131–137]
In critically ill children, do different feed formulas (polymeric vs. semi-elemental feed, standard vs. enriched formula) impact on clinical outcomes?	8.1 Polymeric feeds should be considered as the first choice for enteral nutrition in most critically ill children, unless there are contraindications	GCP	Strong consensus	
	8.2 Protein and energy-dense formulations may be considered to support achievement of nutritional requirements in fluid-restricted critically ill children	B	Consensus	[138, 139]
	8.3 Peptide-based formulations may be considered to improve tolerance and progression of enteral feeding in children for whom polymeric formulations are poorly tolerated or contra-indicated	GCP	Strong consensus	[141]
In critically ill children, does pharmaconutrition (glutamine, lipids and/or micronutrients) impact on clinical outcomes?	9.1 There is insufficient evidence to recommend the use of pharmaconutrition in critically ill children	B	Strong consensus	[81, 141–148]
In critically ill children, does continuous feeding compared to intermittent bolus gastric feeding impact on outcomes?	10.1: There is no evidence to suggest that either continuous or intermittent/bolus feeds are superior in delivering gastric feeds in critically ill children	B	Strong consensus	[70, 149–152]
In critically ill children, does gastric feeding compared to post-pyloric feeding impact on clinical outcomes?	11.1: Gastric feeding is as safe as post pyloric feeding in the majority of critically ill children	C	Strong consensus	[83, 150, 151, 153]
	11.2: Gastric feeding is not inferior to post pyloric feeding in the majority of critically ill children	D	Strong consensus	[150, 151, 153]
	11.3 Post-pyloric feeding can be considered for critically ill children at high risk of aspiration or requiring frequent fasting for surgery or procedures	GCP	Strong consensus	
In critically ill children does routine Gastric Residual Volume (GRV) to guide enteral feeding impact on outcomes?	12.1: Routine measurement of GRV in critically ill children is not recommended	D	Strong consensus	[154]
In critically ill children, do prokinetics impact on clinical outcomes?	13.1: There is insufficient evidence to support the use of prokinetics in critically ill children to improve gastric emptying and feed tolerance	GCP	Strong consensus	[144, 145]
In critically ill children, when should Parenteral Nutrition (PN) be started?	14.1: Withholding parenteral nutrition for up to one week can be considered in critically ill term neonates and children, independent of nutritional status, while providing micronutrients	B	Consensus	[8, 9, 22, 25–27, 156]
In critically ill children, does the use of a feeding protocols impact on clinical outcomes?	15.1: Enteral feeding protocols are recommended to improve time to initiation of EN and nutritional intake	C	Strong consensus	[30, 59, 60, 62, 72, 74–, 76, 87, 90, 157–, 162]
	15.2: Enteral feeding protocols are recommended for high-risk populations to improve nutritional intake and reduce adverse events	D	Strong consensus	[30, 59, 60, 62, 72, 74–76, 87, 90, 157–162]

Table 2 Comparison between American and European guidelines on paediatric intensive care nutrition support

ESPNIC Recommendations (European Society of Paediatric and Neonatology Intensive care)	SCCM and ASPEN recommendations (Society of Critical Care Medicine and American Society for Parenteral and Enteral Nutrition)
Q1: In critically ill children, should nutritional status be assessed and what is the optimal method to assess nutritional status?	Q1A. What is the impact of nutritional status on outcomes in critically ill children? Q1B. What are the best practices to screen and identify patients with malnutrition or those at risk of nutritional deterioration in the PICU?
R1.1: The assessment of nutritional status is recommended in critically ill children at admission and throughout their PICU admission	R1A. Based on observational studies, malnutrition, including obesity, is associated with adverse clinical outcomes including longer periods of ventilation, higher risk of hospital-acquired infection, longer PICU and hospital stay, and increased mortality. We recommend that patients in the PICU undergo detailed nutritional assessment within 48 h of admission. Furthermore, as patients are at risk of nutritional deterioration during hospitalization, which can adversely affect clinical outcomes, we suggest that the nutritional status of patients be re-evaluated at least weekly throughout hospitalization
R1.2: It is recommended to perform anthropometric measurements on admission and regularly during admission, and to express these measurements in z-scores, including weight, height/length mid upper arm circumference and head circumference in young children	R1B. Based on observational studies and expert consensus, we recommend that weight and height/length be measured on admission to the PICU, and z scores for body mass index for-age (weight-for-length < 2 years), or weight-for-age (if accurate, height is not available), be used to screen for patients at extremes of these values. In children < 36-months, head circumference must be documented. Validated screening methods for the PICU population to identify patients at risk of malnutrition must be developed. Screening methods might allow limited resources to be directed to high-risk patients who are most likely to benefit from early nutritional assessment and interventions
Q2: In critically ill children, when should enteral nutrition be commenced and how should it be increased?	Q6B. When should EN be initiated? Q4A. Is EN feasible in critically ill children? Q4B. What is the benefit of EN in this group? Q5A. What is the optimum method for advancing EN in the PICU population?
R2.1: It is recommended to commence early enteral nutrition within 24 h of admission unless contraindicated	R6B. Based on expert opinion, we suggest that EN be initiated in all critically ill children, unless it is contraindicated. Based on observational studies, we suggest early initiation of EN, within the first 24–48 h after admission to the PICU, in eligible patients. We suggest the use of institutional EN guidelines and stepwise algorithms that include criteria for eligibility for EN, timing of initiation, and rate of increase as well as a guide to detecting and managing EN intolerance R4A. Based on observational studies, we recommend EN as the preferred mode of nutrient delivery to the critically ill child. Observational studies support the feasibility of EN, which can be safely delivered to critically ill children with medical and surgical diagnoses, and to those receiving vasoactive medications. Common barriers to EN in the PICU include delayed initiation, interruptions due to perceived intolerance, and prolonged fasting around procedures. Based on observational studies, we suggest that interruptions to EN be minimized to achieve nutrient delivery goals by the enteral route R4B. Although the optimal dose of macronutrients is unclear, some amount of nutrient delivered as EN has been beneficial for gastrointestinal mucosal integrity and motility. Based on large cohort studies, early initiation of EN (within 24–48 h of PICU admission) and achievement of up to two thirds of the nutrient goal in the first week of critical illness have been associated with improved clinical outcomes
R2.2: It is recommended to increase enteral nutrition in a stepwise fashion until goal for delivery is achieved using a feeding protocol or guideline	R5A. Based on observational studies, we suggest the use of a stepwise algorithmic approach to advance EN in children admitted to the PICU. The stepwise algorithm must include bedside support to guide the detection and management of EN intolerance and the optimal rate of increase in EN delivery
Q3: In critically ill children on haemodynamic support (vasoactive medications, extracorporeal life support ECLS) does enteral feeding compared to no enteral feeding affect outcomes?	Q4A. Is EN feasible in critically ill children?

Table 2 (continued)

ESPNIC Recommendations (European Society of Paediatric and Neonatology Intensive care)	SCCM and ASPEN recommendations (Society of Critical Care Medicine and American Society for Parenteral and Enteral Nutrition)
R3.1: Early enteral nutrition is recommended in term neonates who are stable on ECLS	NA
R3.2: Early enteral nutrition is recommended in children who are stable on ECLS	NA
R3.3: Early enteral nutrition is recommended in term neonates who are stable on pharmaceutical haemodynamic support	
R3.4: Early enteral nutrition is recommended in children who are stable on pharmaceutical haemodynamic support	R4A. Based on observational studies, we recommend EN as the preferred mode of nutrient delivery to the critically ill child. Observational studies support the feasibility of EN, which can be safely delivered to critically ill children with medical and surgical diagnoses, and to those receiving vasoactive medications
R3.5: Early enteral nutrition is recommended in children after cardiac surgery	
Q4: In critically ill term neonates with umbilical arterial catheters and/or PGE1 infusions, does enteral feeding impact on adverse events?	NA
R4.1: Enteral nutrition should be considered in term neonates with umbilical arterial catheters	NA
R4.2: Enteral nutrition should be considered in critically ill term neonates on PGE1 infusion if managed in a critical care unit with adequate observation and monitoring	NA
Q5: In critically ill children what are their energy requirements?	Q2C. What is the target energy intake in critically ill children?
R5.1 In the acute phase, energy intake provided to critically ill children should not exceed resting energy expenditure	R2C. Based on observational cohort studies, we suggest achieving delivery of at least two thirds of the prescribed daily energy requirement by the end of the first week in the PICU. Cumulative energy deficits during the first week of critical illness may be associated with poor clinical and nutritional outcomes. Based on expert consensus, we suggest attentiveness to individualized energy requirements, timely initiation and attainment of energy targets, and energy balance to prevent unintended cumulative caloric deficit or excesses
R5.2. After the acute phase, energy intake provided to critically ill children should account for energy debt, physical activity, rehabilitation and growth	
Q6: In critically ill children, what is the most accurate method of determining or predicting energy expenditure?	Q2A. What is the recommended energy requirement for critically ill children?
	Q2B. How should energy requirement be determined in the absence of IC?
R6.1 Measuring resting energy expenditure using a validated indirect calorimeter should be considered to guide nutritional support in critically ill infants and children after the acute phase	R2A. Based on observational cohort studies, we suggest that measured energy expenditure by indirect calorimetry (IC) be used to determine energy requirements and guide prescription of the daily energy goal
R6.2 Schofield equation (for age and gender and using an accurate weight) is recommended to estimate resting energy expenditure	R2B. If IC measurement of resting energy expenditure (REE) is not feasible, we suggest that the Schofield or Food Agriculture Organization/World Health Organization/United Nations University equations may be used "without" the addition of stress factors to estimate energy expenditure. Multiple cohort studies have demonstrated that most published predictive equations are inaccurate and lead to unintended overfeeding or underfeeding. The Harris-Benedict equations and the RDAs, which are suggested by the Dietary Reference Intakes, should not be used to determine energy requirements in critically ill children
Q7.1: What is the recommended glucose intake?	NA
R7.1. Parenteral glucose provision should be sufficient to avoid hypoglycaemia but not excessive to prevent hyperglycaemia	NA
Q7.2: What is the recommended lipid intake or type?	NA
R7.2: When parenteral nutrition is used, composite lipid emulsions, with or without fish oil, should be considered as the first-choice treatment	NA

Table 2 (continued)

ESPNIC Recommendations (European Society of Paediatric and Neonatology Intensive care)	SCCM and ASPEN recommendations (Society of Critical Care Medicine and American Society for Parenteral and Enteral Nutrition)
Q7.3: What is the recommended protein/amino acid intake?	Q3A. What is the minimum recommended protein requirement for critically ill children?
	Q3B. What is the optimal protein delivery strategy in the PICU?
	Q3C. How should protein delivery goals be determined in critically ill children?
R7.3a: For critically ill infants and children on enteral nutrition a minimum enteral protein intake of 1.5 g/kg/d can be considered to avoid negative protein balance	R3A. Based on evidence from RCTs and supported by observational cohort studies, we recommend a minimum protein intake of 1.5 g/kg/d. Protein intake higher than this threshold has been shown to prevent cumulative negative protein balance in RCTs. In critically ill infants and young children, the optimal protein intake required to attain a positive protein balance may be much higher than this minimum threshold. Negative protein balance may result in loss of lean muscle mass, which has been associated with poor outcomes in critically ill patients. Based on a large observational study, higher protein intake may be associated with lower 60-d mortality in mechanically ventilated children
R7.3b: There is insufficient evidence available to support the use of additional protein/amino acid intake during the acute phase of illness (Strong consensus)	R3B. Based on results of randomized trials, we suggest provision of protein early in the course of critical illness to attain protein delivery goals and promote positive nitrogen balance. Delivery of a higher proportion of the protein goal has been associated with positive clinical outcomes in observational studies
Q8: In critically ill children, do different feed formulas (polymeric vs. semi-elemental feed, standard vs. enriched formula) impact on clinical outcomes?	NA
R8.1 Polymeric feeds should be considered as the first choice for enteral nutrition in most critically ill children, unless there are contraindications	NA
R8.2 Protein and energy-dense formulations may be considered to support achievement of nutritional requirements in fluid-restricted critically ill children	NA
R8.3 Peptide-based formulations may be considered to improve tolerance and progression of enteral feeding in children for whom polymeric formulations are poorly tolerated or contra-indicated	NA
Q9: In critically ill children, does pharmaconutrition (glutamine, lipids and/or micronutrients) impact on clinical outcomes?	Q8. What is the role of immunonutrition in critically ill children?
R9.1 There is insufficient evidence to recommend the use of pharmaconutrition in critically ill children	R8. Based on available evidence, we do not recommend the use of immunonutrition in critically ill children
Q10: In critically ill children, does continuous feeding compared to intermittent bolus gastric feeding impact on outcomes?	NA
R10.1: There is no evidence to suggest that either continuous or intermittent/bolus feeds are superior in delivering gastric feeds in critically ill children	NA
Q11: In critically ill children, does gastric feeding compared to post-pyloric feeding impact on clinical outcomes?	Q6A. What is the best site for EN delivery—gastric or small bowel?
R11.1: Gastric feeding is as safe as post pyloric feeding in most critically ill children	R6A. Existing data are insufficient to make universal recommendations regarding the optimal site to deliver EN to critically ill children. Based on observational studies, we suggest the gastric route be the preferred site for EN in patients in the PICU. The post-pyloric or small intestinal site for EN may be used in patients unable to tolerate gastric feeding or those at high risk for aspiration. Existing data are insufficient to make recommendations regarding the use of continuous vs intermittent gastric feeding
R11.2: Gastric feeding is not inferior to post pyloric feeding in the majority of critically ill children	

Table 2 (continued)

ESPNIC Recommendations (European Society of Paediatric and Neonatology Intensive care)	SCCM and ASPEN recommendations (Society of Critical Care Medicine and American Society for Parenteral and Enteral Nutrition)
R11.3 Post-pyloric feeding can be considered for critically ill children at high risk of aspiration or requiring frequent fasting for surgery or procedures	
Q12: In critically ill children does routine Gastric Residual Volume (GRV) to guide enteral feeding impact on outcomes?	NA
R12.1: Routine measurement of GRV in critically ill children is not recommended	NA
Q13: In critically ill children, do prokinetics impact on clinical outcomes?	NA
R13.1: There is insufficient evidence to support the use of prokinetics in critically ill children to improve gastric emptying and feed tolerance	NA
Q14: In critically ill children, when should Parenteral Nutrition (PN) be started?	Q7A. What is the indication for and optimal timing of PN in critically ill children?
R14.1: Withholding parenteral nutrition for up to one week can be considered in critically ill term neonates and children, independent of nutritional status, while providing micronutrients	Q7B. What is the role of PN as a supplement to inadequate EN?
	R7A. Based on a single RCT, we do not recommend the initiation of PN within 24 h of PICU admission
	R7B. In children tolerating EN, we suggest stepwise advancement of nutrient delivery via the enteral route and delaying commencement of PN. Based on current evidence, the role of supplemental PN to reach a specific goal for energy delivery is not known. The time when PN should be initiated to supplement insufficient EN is also unknown. The threshold for and timing of PN initiation should be individualized. Based on a single RCT, supplemental PN should be delayed until 1 week after PICU admission in patients with normal baseline nutritional state and low risk of nutritional deterioration. Based on expert consensus, we suggest PN supplementation in children who are unable to receive any EN during the first week in the PICU. In patients who are severely malnourished or at risk of nutritional deterioration, PN may be supplemented in the first week if they are unable to advance past low volumes of EN
Q15: In critically ill children, does the use of a feeding protocols impact on clinical outcomes?	NA
R15.1: Enteral feeding protocols are recommended to improve time to initiation of EN and nutritional intake	NA
R15.2: Enteral feeding protocols are recommended for high-risk populations to improve nutritional intake and reduce adverse events	NA
NA	Q5B. What is the role of a nutrition support team or a dedicated dietitian in optimizing nutrition therapy?
NA	5B. Based on observational studies, we suggest a nutrition support team, including a dedicated dietitian, be available on the PICU team, to facilitate timely nutritional assessment, and optimal nutrient delivery and adjustment to the patients

NA not applicable, EN enteral nutrition, IC indirect calorimetry, PN parenteral nutrition, RCT randomized controlled trial, RDA recommended daily allowance

outcome in critically ill children [26, 27]. Despite the lack of effect shown of protocols on mortality and NEC in the meta-analysis, as the level of this evidence was low, all individual studies did show positive effect on other variables such as time to initiate feeding and achievement of energy goals, but it was not possible to pool these in a meta-analysis. Therefore, despite this, we still recommend PICUs use feeding protocols which provide guidance on the assessment of nutritional status and the start and advancement of feeding. A final key messages from this position statement is to encourage the

enteral feeding of critically ill neonates and children early wherever possible, unless clear contraindications exist. Although starting early EN is recommended, no evidence exists to support high nutritional intake during the acute phase of critical illness and withholding supplemental PN during the first week in PICU may be considered when enteral nutrition is insufficient.

Limitations

We acknowledge that these clinical recommendations are based at times on sparse paediatric evidence. Moreover,

for many questions and clinical recommendations we could not be age-specific, although the (patho)physiology of nutritional and metabolic changes during critical illness is age-dependent. For instance, the recommendations specifically for neonates were partially based upon studies in preterm neonates as no evidence existed in term neonates. The threshold of > 37 weeks in our recommendations is recognized as rigid and we cannot exclude that some of our recommendations also apply for late preterm (> 34 weeks) or early term (> 36 weeks) neonates. Similarly, the same arguments can be raised for adolescents, where for certain (older) adolescents, recommendations from adult guidelines might be suitable. However, the mean age in adult ICUs is 60.9 years [28] and it, therefore, cannot be assumed that critically ill young adults are similar in their (patho)physiologic response to nutritional and metabolic changes to elderly patients. We further acknowledge that, as a priori anticipated, pragmatic meta-analyses were required due to the inconsistencies in the outcome variables. Another limitation is that our consensus voting was based only on the views of our study team of 11 experts. Finally, as already elaborated on in our discussion, aside from several novel features our recommendations have an overlap with the American (ASPEN) guidelines published in 2017. A future collaboration between the American and European societies might improve upcoming guidelines and help implement the recommendations worldwide. Despite these limitations, this work has provided an updated summary of the existing evidence, including questions around term neonates, which are not dealt with by other recommendations or guidelines, yet comprise a significant amount of the European PICU population [29, 30].

Conclusion

This ESPNIC position statement with recommendations provides a 'best-available-evidence' guide for clinicians working in PICU to provide nutritional support in this setting. The lack of methodologically sound trials and the heterogeneous character of studies available were important barriers in the generation of these recommendations. Many recommendations are based on expert consensus and have a low level of evidence. Nonetheless, our recommendations support the use of a formal nutritional assessment and a feeding protocol in all PICUs.

Electronic supplementary material

The online version of this article (<https://doi.org/10.1007/s00134-019-05922-5>) contains supplementary material, which is available to authorized users.

Author details

¹ Faculty of Health and Society, University of Salford, Manchester M6 6PU, UK. ² Pediatric Intensive Care Unit, Alder Hey Children's NHS Foundation Trust, East Prescott Road, Liverpool L12 2AP, UK. ³ Pediatric Intensive Care Unit, Hôpital Femme Mère Enfant, CarMEN INSERM UMR, 1060 Hospices Civils de Lyon, Lyon-Bron, France.

⁴ Intensive Care, Department of Pediatrics and Pediatric Surgery, Erasmus Medical Centre, Sophia Children's Hospital, Rotterdam, The Netherlands. ⁵ Geneva School of Health Sciences, HES-SO University of Applied Sciences and Arts Western Switzerland, Delémont, Switzerland. ⁶ Pediatric Intensive Care Unit, University Hospital of Lausanne, Lausanne, Switzerland. ⁷ Nutrition and Dietetics, Alder Hey Children's Hospital Liverpool, Liverpool, UK. ⁸ Department of Dietetics/Speech and Language Therapy, NIHR Biomedical Research Centre Southampton, University Hospital Southampton, Faculty of Environmental and Life Sciences, University of Southampton, Southampton, UK. ⁹ Pediatric Intensive Care Unit, Royal Hospital for Children, Glasgow, UK. ¹⁰ Department of Pediatrics, University of Cambridge, Hills Road, Cambridge, UK. ¹¹ Pediatric Intensive Care Unit, Queen Fabiola Children's University Hospital, Brussels, Belgium. ¹² Department of Biostatistics, Erasmus Medical Centre Rotterdam, Rotterdam, The Netherlands.

Acknowledgements

We want to thank Elise Krabbendam, Sabrina Gunput, Gerdien de Jong and Wichor Bramer, Biomedical Information Specialists of the Medical Library of the Erasmus Medical Centre, Rotterdam, The Netherlands. The costs covering the open access publication of this article are covered by ESPNIC.

Funding

No industry funding was provided. The European Society of Pediatric and Neonatal Intensive Care (ESPNIC) provided logistical support and funding for the meetings.

Compliance with ethical standards

Conflicts of interest

The authors of these guidelines have reported all potential conflicts or financial disclosures. For a complete overview of the potential conflicts and financial disclosures see below. No funding or contribution from industry was involved in the completion of these recommendations, nor were any industry representatives present at any of the committee meetings. LNT is a member of the NIHR. SCV is supported by Nutricia Advanced Medical Nutrition (unrestricted research grant). SCV has received speakers' fees from Nutricia Advanced Medical Nutrition and Baxter in the past. KJ is supported by Nutricia Advanced Medical Nutrition (unrestricted research grant). FVW has received speaker fees from Fresenius Kabi and Nutricia (past) and consultant fees from Baxter (current). LVM is supported by a Health Education England/NIHR Clinical Lectureship (ICA-CL-2016-02-001) supported by the National Institute for Health Research. LVM has also received speaker's fees from Abbott Laboratories and Nutricia in the past. NP is supported by research funding from the National Institute for Health Research. CJC is supported by the Marisa Sophie Foundation and has received travel fees from Nutricia and Baxter (past). CM is supported by the Marisa Sophie Foundation and has received travel fees from Nutricia and Baxter (past).

Open Access

This article is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License, which permits any non-commercial use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by-nc/4.0/>.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 26 November 2019 Accepted: 28 December 2019
Published online: 20 February 2020

References

- Joosten KF, Kerklaan D, Verbruggen SC (2016) Nutritional support and the role of the stress response in critically ill children. *Curr Opin Clin Nutr Metab Care* 19:226–233
- Joosten KFM, Eveleens RD, Verbruggen S (2019) Nutritional support in the recovery phase of critically ill children. *Curr Opin Clin Nutr Metab Care* 22(2):152–158. <https://doi.org/10.1097/MCO.0000000000000549>
- Keehn A, O'Brien C, Mazurak V et al (2015) Epidemiology of interruptions to nutrition support in critically ill children in the pediatric intensive care unit. *JPEN J Parenter Enter Nutr* 39:211–217
- Leong AY, Cartwright KR, Guerra GG, Joffe AR, Mazurak VC, Larsen BM (2014) A Canadian survey of perceived barriers to initiation and continuation of enteral feeding in PICUs. *Pediatr Crit Care Med* 15:e49–55
- Hulst J, Joosten K, Zimmermann L et al (2004) Malnutrition in critically ill children: from admission to 6 months after discharge. *Clin Nutr* 23:223–232
- Hulst JM, van Goudoever JB, Zimmermann LJ et al (2004) The effect of cumulative energy and protein deficiency on anthropometric parameters in a pediatric ICU population. *Clin Nutr* 23:1381–1389
- Prince NJ, Brown KL, Mebrahtu TF, Parslow RC, Peters MJ (2014) Weight-for-age distribution and case-mix adjusted outcomes of 14,307 pediatric intensive care admissions. *Intensive Care Med* 40:1132–1139
- van Puffelen E, Hulst JM, Vanhorebeek I, Dulfer K, Van den Berghe G, Verbruggen SCAT, Joosten KFM (2018) Outcomes of delaying parenteral nutrition for 1 week vs initiation within 24 hours among undernourished children in pediatric intensive care. *JAMA Netw Open* 1:e182668
- van Puffelen E, Hulst JM, Vanhorebeek I et al (2020) Effect of late versus early initiation of parenteral nutrition on weight deterioration during PICU stay: secondary analysis of the PEPaNIC randomised controlled trial. *Clin Nutr* 39(1):104–109. <https://doi.org/10.1016/j.clnu.2019.02.014>
- Valla FV, Young DK, Rabilloud M et al (2017) Thigh ultrasound monitoring identifies decreases in quadriceps femoris thickness as a frequent observation in critically ill children. *Pediatr Crit Care Med* 18:e339–e347
- Joffe A, Anton N, Lequier L et al (2009) Nutritional support for critically ill children. *Cochrane Database Syst Rev* (5):CD005144. <https://doi.org/10.1002/14651858.CD005144>
- Fivez T, Kerklaan D, Mesotten D, Verbruggen S, Joosten K, Van den Berghe G (2017) Evidence for the use of parenteral nutrition in the pediatric intensive care unit. *Clin Nutr* 36:218–223
- Tume LN, Balmaks R, da Cruz E et al (2018) Enteral feeding practices in infants with congenital heart disease across European PICUs: a European Society of Pediatric and Neonatal Intensive Care Survey. *Pediatr Crit Care Med* 19:137–144
- Kerklaan D, Fivez T, Mehta NM et al (2016) Worldwide survey of nutritional practices in PICUs. *Pediatr Crit Care Med* 17:10–18
- Mehta NM, Skillman HE, Irving SY et al (2017) Guidelines for the provision and assessment of nutrition support therapy in the pediatric critically ill patient: Society of Critical Care Medicine and American Society for Parenteral and Enteral Nutrition. *Pediatr Crit Care Med* 18:675–715
- Valla FV, Gaillard Le Roux B, Tume LN (2017) Pediatric intensive care nutrition guidelines 2017: key questions remain unanswered. *JPEN J Parenter Enter Nutr* 2018(42):9
- https://www.picanet.org.uk/Audit/Annual-Reporting/PICANet_2017_Annual_Report_Tables_and_Figures_FINAL_v2.0.pdf. Accessed PICANET, 2018. Accessed November 2019, 2018
- Harbour R, Miller J (2001) A new system for grading recommendations in evidence based guidelines. *BMJ* 323:334–336
- Hasson F, Keeney S, McKenna H (2000) Research guidelines for the Delphi survey technique. *J Adv Nurs* 32:1008–1015
- Keeney S, Hasson F, McKenna HP (2001) A critical review of the Delphi technique as a research methodology for nursing. *Int J Nurs Stud* 38:195–200
- van Puffelen E, Jacobs A, Verdoorn CJM et al (2019) International survey of de-implementation of initiating parenteral nutrition early in pediatric intensive care units. *BMC Health Serv Res* 19:379
- van Puffelen E, Vanhorebeek I, Joosten KFM, Wouters PJ, Van den Berghe G, Verbruggen S (2018) Early versus late parenteral nutrition in critically ill, term neonates: a preplanned secondary subgroup analysis of the PEPaNIC multicentre, randomised controlled trial. *Lancet Child Adolesc Health* 2:505–515
- Jotterand Chaparro C, Laure Depeyre J, Longchamp D, Perez MH, Taffe P, Cotting J (2016) How much protein and energy are needed to equilibrate nitrogen and energy balances in ventilated critically ill children? *Clin Nutr* 35:460–467
- Bechard LJ, Parrott JS, Mehta NM (2012) Systematic review of the influence of energy and protein intake on protein balance in critically ill children. *J Pediatr* 161(333–9):e1
- Vanhorebeek I, Verbruggen S, Casaer MP et al (2017) Effect of early supplemental parenteral nutrition in the pediatric ICU: a preplanned observational study of post-randomisation treatments in the PEPaNIC trial. *Lancet Respir Med* 5:475–483
- Fivez T, Kerklaan D, Mesotten D et al (2016) Early versus late parenteral nutrition in critically ill children. *N Engl J Med* 374:1111–1122
- Verstraete S, Verbruggen SC, Hordijk JA et al (2019) Long-term developmental effects of withholding parenteral nutrition for 1 week in the pediatric intensive care unit: a 2-year follow-up of the PEPaNIC international, randomised, controlled trial. *Lancet Respir Med* 7:141–153
- UK Intensive Care National Audit and Research centre. Key statistics from the Case Mix Programme—adult, general critical care units 2015–2016. <https://www.icnarc.org.uk>. Accessed Dec 2019
- Simsic JM, Carpenito KR, Kirchner K et al (2017) Reducing variation in feeding newborns with congenital heart disease. *Congenit Heart Dis* 12:275–281
- Furlong-Dillard J, Neary A, Marietta J et al (2018) Evaluating the impact of a feeding protocol in neonates before and after biventricular cardiac surgery. *Pediatr Qual Saf* 3:e080
- Wakita M, Fukatsu A, Amagai T (2011) Nutrition assessment as a predictor of clinical outcomes for infants with cardiac surgery: using the prognostic nutritional index. *Nutr Clin Pract* 26:192–198
- Vermilyea S, Slicker J, El-Chammas K et al (2013) Subjective global nutritional assessment in critically ill children. *JPEN J Parenter Enter Nutr* 37:659–666
- Valla FV, Ford-Chessel C, Meyer R et al (2015) A training program for anthropometric measurements by a dedicated nutrition support team improves nutritional status assessment of the critically ill child. *Pediatr Crit Care Med* 16:e82–e88
- Irving SY, Seiple S, Nagle M, Falk S, Mascarenhas M, Srinivasan V (2015) Perceived barriers to anthropometric measurements in critically ill children. *Am J Crit Care* 24:e99–e107
- Valla FV, Berthiller J, Gaillard-Le-Roux B et al (2018) Faltering growth in the critically ill child: prevalence, risk factors, and impaired outcome. *Eur J Pediatr* 177:345–353
- Grippa RB, Silva PS, Barbosa E, Bresolin NL, Mehta NM, Moreno YM (2017) Nutritional status as a predictor of duration of mechanical ventilation in critically ill children. *Nutrition* 33:91–95
- Sotoudeh M, Khalili M, Azizian M, Imani B (2016) Prevalence of malnutrition based on under weight inpatients in pediatric intensive care unit. *Int J Pharm Technol* 8:12333–12340
- Bechard LJ, Duggan C, Touger-Decker R et al (2016) Nutritional status based on body mass index is associated with morbidity and mortality in mechanically ventilated critically ill children in the PICU. *Crit Care Med* 44:1530–1537
- Radman M, Mack R, Barnoya J et al (2014) The effect of preoperative nutritional status on postoperative outcomes in children undergoing surgery for congenital heart defects in San Francisco (UCSF) and Guatemala City (UNICAR). *J Thorac Cardiovasc Surg* 147:442–450
- Anton-Martin P, Papacostas M, Lee E, Nakonezny PA, Green ML (2018) Underweight status is an independent predictor of in-hospital mortality in pediatric patients on extracorporeal membrane oxygenation. *JPEN J Parenter Enter Nutr* 42:104–111
- Leite HP, de Lima LF, de Oliveira Iglesias SB, Pacheco JC, de Carvalho WB (2013) Malnutrition may worsen the prognosis of critically ill children with hyperglycemia and hypoglycemia. *JPEN J Parenter Enter Nutr* 37:335–341
- Zamberlan P, Leone C, Tannuri U, Carvalho WB, Delgado AF (2012) Nutritional risk and anthropometric evaluation in pediatric liver transplantation. *Clinics (Sao Paulo)* 67:1387–1392
- de Souza MF, Leite HP, Koch Nogueira PC (2012) Malnutrition as an independent predictor of clinical outcome in critically ill children. *Nutrition* 28:267–270

44. Castillo A, Santiago MJ, Lopez-Herce J et al (2012) Nutritional status and clinical outcome of children on continuous renal replacement therapy: a prospective observational study. *BMC Nephrol* 13:125
45. Numa A, McAweeney J, Williams G, Awad J, Ravindranathan H (2011) Extremes of weight centile are associated with increased risk of mortality in pediatric intensive care. *Crit Care* 15:R106
46. Botran M, Lopez-Herce J, Mencía S et al (2011) Relationship between energy expenditure, nutritional status and clinical severity before starting enteral nutrition in critically ill children. *Br J Nutr* 105:731–737
47. Srinivasan V, Nadkarni VM, Helfaer MA, Carey SM, Berg RA (2010) American Heart Association National Registry of Cardiopulmonary Resuscitation I Childhood obesity and survival after in-hospital pediatric cardiopulmonary resuscitation. *Pediatrics* 125:e481–e488
48. Anderson JB, Beekman RH 3rd, Border WL et al (2009) Lower weight-for-age z score adversely affects hospital length of stay after the bidirectional Glenn procedure in 100 infants with a single ventricle. *J Thorac Cardiovasc Surg* 138(397–404):e1
49. Eskedal LT, Hagemo PS, Seem E et al (2008) Impaired weight gain predicts risk of late death after surgery for congenital heart defects. *Arch Dis Child* 93:495–501
50. Kelleher DK, Laussen P, Teixeira-Pinto A, Duggan C (2006) Growth and correlates of nutritional status among infants with hypoplastic left heart syndrome (HLHS) after stage 1 Norwood procedure. *Nutrition* 22:237–244
51. Hulst JM, van Goudoever JB, Zimmermann LJ, Tibboel D, Joosten KF (2006) The role of initial monitoring of routine biochemical nutritional markers in critically ill children. *J Nutr Biochem* 17:57–62
52. Fivez T, Hendrickx A, Van Herpe T et al (2016) An analysis of reliability and accuracy of muscle thickness ultrasonography in critically ill children and adults. *JPEN J Parenter Enter Nutr* 40:944–949
53. Briassoulis G, Zavras N, Hatzis T (2001) Malnutrition, nutritional indices, and early enteral feeding in critically ill children. *Nutrition* 17:548–557
54. Davis ET, Xie L, Levenbrown Y (2017) Impact of obesity on outcomes in critically ill children. *JPEN J Parenter Enter Nutr*. <https://doi.org/10.1177/0148607117725043>
55. Ross PA, Newth CJ, Leung D, Wetzel RC, Khemani RG (2016) Obesity and mortality risk in critically ill children. *Pediatrics* 137:e20152035
56. Ward SL, Gildengorin V, Valentine SL et al (2016) Impact of weight extremes on clinical outcomes in pediatric acute respiratory distress syndrome. *Crit Care Med* 44:2052–2059
57. Goh VL, Wakeham MK, Brazauskas R, Mikhailov TA, Goday PS (2013) Obesity is not associated with increased mortality and morbidity in critically ill children. *JPEN J Parenter Enter Nutr* 37:102–108
58. Abdul Manaf Z, Kassim N, Hamzaidd NH, Razali NH (2013) Delivery of enteral nutrition for critically ill children. *Nutr Diet* 70:120–125
59. Ang B, Han WM, Wong JJM, Lee AN, Chan YH, Lee JH (2016) Impact of a nurse-led feeding protocol in a pediatric intensive care unit. *Proc Singap Healthc* 25:35–42
60. Braudis NJ, Curley MA, Beaupre K et al (2009) Enteral feeding algorithm for infants with hypoplastic left heart syndrome poststage I palliation. *Pediatr Crit Care Med* 10:460–466
61. Canarie MF, Barry S, Carroll CL et al (2015) Risk Factors for delayed enteral nutrition in critically ill children. *Pediatr Crit Care Med* 16:e283–e289
62. Carpenito KR, Prusinski R, Kirchner K et al (2016) Results of a feeding protocol in patients undergoing the hybrid procedure. *Pediatr Cardiol* 37:852–859
63. Lee H, Koh SO, Kim H, Sohn MH, Kim KE, Kim KW (2013) Avoidable causes of delayed enteral nutrition in critically ill children. *J Korean Med Sci* 28:1055–1059
64. Mikhailov TA, Kuhn EM, Manzi J et al (2014) Early enteral nutrition is associated with lower mortality in critically ill children. *JPEN J Parenter Enter Nutr* 38:459–466
65. Tripathi S, Kaur H, Varayil JE, Hurt RT (2015) Effects of enteral nutrition on clinical outcomes among mechanically ventilated and sedated patients in the pediatric intensive care unit. *Signa Vitae* 10(2):131–148. <https://doi.org/10.22514/SV102.122015.8>
66. Mikhailov TA, Gertz SJ, Kuhn EM, Scanlon MC, Rice TB, Goday PS (2018) Early enteral nutrition is associated with significantly lower hospital charges in critically ill children. *JPEN J Parenter Enter Nutr* 42:920–925
67. Bagci S, Keles E, Girgin F et al (2018) Early initiated feeding versus early reached target enteral nutrition in critically ill children: an observational study in pediatric intensive care units in Turkey. *J Paediatr Child Health* 54:480–486
68. Balakrishnan B, Flynn-O'Brien KT, Simpson PM, Dasgupta M, Hanson SJ (2019) Enteral nutrition initiation in children admitted to pediatric intensive care units after traumatic brain injury. *Neurocrit Care* 30:193–200
69. Malakouti A, Sookplung P, Siriussawakul A et al (2012) Nutrition support and deficiencies in children with severe traumatic brain injury. *Pediatr Crit Care Med* 13:e18–24
70. Fayazi S, Adineh M, Zahraei Fard S, Farokh Payam S, Ahmadi Batvandy Z (2016) Comparing two methods of enteral nutrition in terms of their complications and the time needed to reach goal calorie in children hospitalized in ICU. *Int J Pediatr* 4:2119–2130
71. Gurgueira GL, Leite HP, Taddei JA, de Carvalho WB (2005) Outcomes in a pediatric intensive care unit before and after the implementation of a nutrition support team. *JPEN J Parenter Enter Nutr* 29:176–185
72. Hamilton S, McAleer DM, Ariagno K et al (2014) A stepwise enteral nutrition algorithm for critically ill children helps achieve nutrient delivery goals*. *Pediatr Crit Care Med* 15:583–589
73. Mara J, Gentles E, Alfheda HA et al (2014) An evaluation of enteral nutrition practices and nutritional provision in children during the entire length of stay in critical care. *BMC Pediatr* 14:186
74. Meyer R, Harrison S, Sargent S, Ramnarayan P, Habibi P, Labadarios D (2009) The impact of enteral feeding protocols on nutritional support in critically ill children. *J Hum Nutr Diet* 22:428–436
75. Petrillo-Albarano T, Pettignano R, Asfaw M, Easley K (2006) Use of a feeding protocol to improve nutritional support through early, aggressive, enteral nutrition in the pediatric intensive care unit. *Pediatr Crit Care Med* 7:340–344
76. Yoshimura S, Miyazu M, Yoshizawa S et al (2015) Efficacy of an enteral feeding protocol for providing nutritional support after pediatric cardiac surgery. *Anaesth Intensive Care* 43:587–593
77. Hanekamp MN, Spoel M, Sharman-Koendjibiharie M et al (2005) Gut hormone profiles in critically ill neonates on extracorporeal membrane oxygenation. *J Pediatr Gastroenterol Nutr* 40:175–179
78. Hanekamp MN, Spoel M, Sharman-Koendjibiharie I, Peters JW, Albers MJ, Tibboel D (2005) Routine enteral nutrition in neonates on extracorporeal membrane oxygenation. *Pediatr Crit Care Med* 6:275–279
79. Piena M, Albers MJ, Van Haard PM, Gischler S, Tibboel D (1998) Introduction of enteral feeding in neonates on extracorporeal membrane oxygenation after evaluation of intestinal permeability changes. *J Pediatr Surg* 33:30–34
80. Wertheim HF, Albers MJ, Piena-Spoel M, Tibboel D (2001) The incidence of septic complications in newborns on extracorporeal membrane oxygenation is not affected by feeding route. *J Pediatr Surg* 36:1485–1489
81. Ong EG, Eaton S, Wade AM et al (2012) Randomized clinical trial of glutamine-supplemented versus standard parenteral nutrition in infants with surgical gastrointestinal disease. *Br J Surg* 99:929–938
82. Pettignano R, Heard M, Davis R, Labuz M, Hart M (1998) Total enteral nutrition versus total parenteral nutrition during pediatric extracorporeal membrane oxygenation. *Crit Care Med* 26:358–363
83. López-Herce J, Mencía S, Sánchez C, Santiago MJ, Bustinza A, Vigil D (2008) Postpyloric enteral nutrition in the critically ill child with shock: a prospective observational study. *Nutr J* 7:6. <https://doi.org/10.1186/1475-2891-7-6>
84. Iannucci GJ, Oster ME, Mahle WT (2013) Necrotizing enterocolitis in infants with congenital heart disease: the role of enteral feeds. *Cardiol Young* 23:553–559
85. King W, Petrillo T, Pettignano R (2004) Enteral nutrition and cardiovascular medications in the pediatric intensive care unit. *JPEN J Parenter Enter Nutr* 28:334–338
86. Panchal AK, Manzi J, Connolly S et al (2016) Safety of enteral feedings in critically ill children receiving vasoactive agents. *JPEN J Parenter Enter Nutr* 40:236–241
87. del Castillo SL, McCulley ME, Khemani RG et al (2010) Reducing the incidence of necrotizing enterocolitis in neonates with hypoplastic left heart syndrome with the introduction of an enteral feed protocol. *Pediatr Crit Care Med* 11:373–377

-
88. Jeffries HE, Wells WJ, Starnes VA, Wetzel RC, Moromisato DY (2006) Gastrointestinal morbidity after Norwood palliation for hypoplastic left heart syndrome. *Ann Thorac Surg* 81:982–987
 89. Luce WA, Schwartz RM, Beauseau W et al (2011) Necrotizing enterocolitis in neonates undergoing the hybrid approach to complex congenital heart disease. *Pediatr Crit Care Med* 12:46–51
 90. Manuri L, Morelli S, Agati S et al (2017) Early hybrid approach and enteral feeding algorithm could reduce the incidence of necrotizing enterocolitis in neonates with ductus-dependent systemic circulation. *Cardiol Young* 27:154–160
 91. Natarajan G, Reddy Anne S, Aggarwal S (2010) Enteral feeding of neonates with congenital heart disease. *Neonatology* 98:330–336
 92. Sánchez C, López-Herce J, Carrillo A, Bustinza A, Sancho L, Vigil D (2006) Transpyloric enteral feeding in the postoperative of cardiac surgery in children. *J Pediatr Surg* 41:1096–1102
 93. Sahu MK, Singal A, Menon R et al (2016) Early enteral nutrition therapy in congenital cardiac repair postoperatively: a randomized, controlled pilot study. *Ann Card Anaesth* 19:653–661
 94. Typpo KV, Larmonier CB, Deschenes J, Redford D, Kiela PR, Ghishan FK (2015) Clinical characteristics associated with postoperative intestinal epithelial barrier dysfunction in children with congenital heart disease. *Pediatr Crit Care Med* 16:37–44
 95. Davey AM, Wagner CL, Cox C, Kendig JW (1994) Feeding premature infants while low umbilical artery catheters are in place: a prospective, randomized trial. *J Pediatr* 124:795–799
 96. Barrington KJ (2000) Umbilical artery catheters in the newborn: effects of position of the catheter tip. *Cochrane Database Syst Rev* (2):CD000505
 97. Willis L, Thurene P, Kaufman J, Wymore E, Skillman H, da Cruz E (2008) Enteral feeding in prostaglandin-dependent neonates: is it a safe practice? *J Pediatr* 153:867–869
 98. Lucron H, Chipaux M, Bossier G et al (2005) [Complications of prostaglandin E1 treatment of congenital heart disease in pediatric medical intensive care] Complications du traitement par prostaglandines E1 des cardiopathies congénitales en réanimation médicale pédiatrique. *Arch Mal Coeur Vaiss* 98:524–530
 99. Becker KC, Hornik CP, Cotten CM et al (2015) Necrotizing enterocolitis in infants with ductal-dependent congenital heart disease. *Am J Perinatol* 32:633–638
 100. Havranek T, Johanboeke P, Madramootoo C, Carver JD (2007) Umbilical artery catheters do not affect intestinal blood flow responses to minimal enteral feedings. *J Perinatol* 27:375–379
 101. Larsen BMK, Beggs MR, Leong AY, Kang SH, Persad R, Garcia GG (2018) Can energy intake alter clinical and hospital outcomes in PICU? *Clin Nutr ESPEN* 24:41–46
 102. Joosten KF, Verhoeven JJ, Hazelzet JA (1999) Energy expenditure and substrate utilization in mechanically ventilated children. *Nutrition* 15:444–448
 103. Dylewski M, Wibbenmeyer L, Bessey PQ et al (2013) Nutrition outcomes. *J Burn Care Res* 34:371–375
 104. Coss-Bu JA, Jefferson LS, Walding D, David Y, Smith EO, Klish WJ (1998) Resting energy expenditure and nitrogen balance in critically ill pediatric patients on mechanical ventilation. *Nutrition* 14:649–652
 105. de Betue CT, van Waardenburg DA, Deutz NE et al (2011) Increased protein-energy intake promotes anabolism in critically ill infants with viral bronchiolitis: a double-blind randomised controlled trial. *Arch Dis Child* 96:817–822
 106. van Waardenburg DA, de Betue CT, Goudoever JB, Zimmermann LJ, Joosten KF (2009) Critically ill infants benefit from early administration of protein and energy-enriched formula: a randomized controlled trial. *Clin Nutr* 28:249–255
 107. Coss-Bu JA, Klish WJ, Walding D, Stein F, Smith EO, Jefferson LS (2001) Energy metabolism, nitrogen balance, and substrate utilization in critically ill children. *Am J Clin Nutr* 74:664–669
 108. Briassoulis G, Tsorva A, Zavras N, Hatzis T (2002) Influence of an aggressive early enteral nutrition protocol on nitrogen balance in critically ill children. *J Nutr Biochem* 13:560
 109. Mehta NM, Bechard LJ, Cahill N et al (2012) Nutritional practices and their relationship to clinical outcomes in critically ill children—an international multicenter cohort study*. *Crit Care Med* 40:2204–2211
 110. Mtaweh H, Smith R, Kochanek PM et al (2014) Energy expenditure in children after severe traumatic brain injury. *Pediatr Crit Care Med* 15:242–249
 111. De Wit B, Meyer R, Desai A, Macrae D, Pathan N (2010) Challenge of predicting resting energy expenditure in children undergoing surgery for congenital heart disease. *Pediatr Crit Care Med* 11:496–501
 112. Sy J, Gourishankar A, Gordon WE et al (2008) Bicarbonate kinetics and predicted energy expenditure in critically ill children. *Am J Clin Nutr* 88:340–347
 113. Framson CM, LeLeiko NS, Dallal GE, Roubenoff R, Snelling LK, Dwyer JT (2007) Energy expenditure in critically ill children. *Pediatr Crit Care Med* 8:264–267
 114. Havalad S, Quaid MA, Sapiaga V (2006) Energy expenditure in children with severe head injury: lack of agreement between measured and estimated energy expenditure. *Nutr Clin Pract* 21:175–181
 115. van der Kuip M, de Meer K, Oosterveld MJ, Lafeber HN, Gemke RJ (2004) Simple and accurate assessment of energy expenditure in ventilated pediatric intensive care patients. *Clin Nutr* 23:657–663
 116. Vazquez Martinez JL, Martinez-Romillo PD, Diez Sebastian J, Ruza TF (2004) Predicted versus measured energy expenditure by continuous, online indirect calorimetry in ventilated, critically ill children during the early postinjury period. *Pediatr Crit Care Med* 5:19–27
 117. Hardy CM, Dwyer J, Snelling LK, Dallal GE, Adelson JW (2002) Pitfalls in predicting resting energy requirements in critically ill children: a comparison of predictive methods to indirect calorimetry. *Nutr Clin Pract* 17:182–189
 118. White MS, Shepherd RW, McEniery JA (2000) Energy expenditure in 100 ventilated, critically ill children: improving the accuracy of predictive equations. *Crit Care Med* 28:2307–2312
 119. Briassoulis G, Venkataraman S, Thompson AE (2000) Energy expenditure in critically ill children. *Crit Care Med* 28:1166–1172
 120. Verhoeven JJ, Hazelzet JA, van der Voort E, Joosten KF (1998) Comparison of measured and predicted energy expenditure in mechanically ventilated children. *Intensive Care Med* 24:464–468
 121. Jotterand Chaparro C, Moullet C, Taffe P et al (2018) Estimation of resting energy expenditure using predictive equations in critically ill children: results of a systematic review. *JPEN J Parenter Enter Nutr* 42:976–986
 122. Mehta NM, Smallwood CD, Joosten KF, Hulst JM, Tasker RC, Duggan CP (2015) Accuracy of a simplified equation for energy expenditure based on bedside volumetric carbon dioxide elimination measurement—a two-center study. *Clin Nutr* 34:151–155
 123. Meyer R, Kulinskaya E, Briassoulis G et al (2012) The challenge of developing a new predictive formula to estimate energy requirements in ventilated critically ill children. *Nutr Clin Pract* 27:669–676
 124. Lopez-Herce Cid J, Sanchez Sanchez C, Mencia Bartolome S, Santiago Lozano MJ, Carrillo Alvarez A, Bellon Cano JM (2007) [Energy expenditure in critically ill children: correlation with clinical characteristics, caloric intake, and predictive equations] Consumo calorico en el niño crítico: relacion con las características clínicas, el aporte calórico y las formulas teoricas de calculo de las necesidades. *An Pediatr (Barc)* 66:229–239
 125. Taylor RM, Cheeseman P, Preedy V, Baker AJ, Grimble G (2003) Can energy expenditure be predicted in critically ill children? *Pediatr Crit Care Med* 4:176–180
 126. Verbruggen SC, de Betue CT, Schierbeek H et al (2011) Reducing glucose infusion safely prevents hyperglycemia in post-surgical children. *Clin Nutr* 30:786–792
 127. de Betue CT, Verbruggen SC, Schierbeek H et al (2012) Does a reduced glucose intake prevent hyperglycemia in children early after cardiac surgery? a randomized controlled crossover study. *Crit Care* 16:R176
 128. Larsen BM, Goonewardene LA, Joffe AR, Van Aerde JE, Field CJ, Olstad DL, Clandinin MT (2012) Pre-treatment with an intravenous lipid emulsion containing fish oil (eicosapentaenoic and docosahexaenoic acid) decreases inflammatory markers after open-heart surgery in infants: a randomized, controlled trial. *Clin Nutr* 31:322–329
 129. Geukens VG, Dijsselhof ME, Jansen NJ et al (2015) The effect of short-term high versus normal protein intake on whole-body protein synthesis and balance in children following cardiac surgery: a randomized double-blind controlled clinical trial. *Nutr J* 14:72

130. Botran M, Lopez-Herce J, Mencia S, Urbano J, Solana MJ, Garcia A (2011) Enteral nutrition in the critically ill child: comparison of standard and protein-enriched diets. *J Pediatr* 159(27–32):e1
131. Verbruggen SC, Coss-Bu J, Wu M et al (2011) Current recommended parenteral protein intakes do not support protein synthesis in critically ill septic, insulin-resistant adolescents with tight glucose control. *Crit Care Med* 39:2518–2525
132. Hauschild DB, Oliveira LDA, Farias MS et al (2019) Enteral protein supplementation in critically ill children: a randomized controlled pilot and feasibility study. *JPEN J Parenter Enter Nutr* 43:281–289
133. Kyle UG, Akcan-Arikan A, Silva JC, Goldsworthy M, Shekerdemian LS, Coss-Bu JA (2017) Protein feeding in pediatric acute kidney injury is not associated with a delay in renal recovery. *J Ren Nutr* 27:8–15
134. Wong JJ, Han WM, Sultana R, Loh TF, Lee JH (2017) Nutrition delivery affects outcomes in pediatric acute respiratory distress syndrome. *JPEN J Parenter Enter Nutr* 41:1007–1013
135. Mehta NM, Bechard LJ, Zurawski D, Duggan CP, Heyland DK (2015) Adequate enteral protein intake is inversely associated with 60-d mortality in critically ill children: a multicenter, prospective, cohort study. *Am J Clin Nutr* 102:199–206
136. Merhar SL, Meinen-Derr J, Sprague J et al (2015) Safety and tolerability of enteral protein supplementation for infants with brain injury. *Nutr Clin Pract* 30:546–550
137. de Betue CT, Joosten KF, Deutz NE, Vreugdenhil AC, van Waardenburg DA (2013) Arginine appearance and nitric oxide synthesis in critically ill infants can be increased with a protein-energy-enriched enteral formula. *Am J Clin Nutr* 98:907–916
138. Zhang H, Gu Y, Mi Y, Jin Y, Fu W, Latour JM (2019) High-energy nutrition in pediatric cardiac critical care patients: a randomized controlled trial. *Nurs Crit Care* 24(2):97–102. <https://doi.org/10.1111/nicc.12400>
139. Eveleens RD, Dungen DK, Verbruggen S, Hulst JM, Joosten KFM (2018) Weight improvement with the use of protein and energy enriched nutritional formula in infants with a prolonged PICU stay. *J Hum Nutr Diet* 32(1):3–10. <https://doi.org/10.1111/jhn.12603>
140. Vidigal MV, Leite HP, Nogueira PC (2012) Factors associated with peptide-based formula prescription in a pediatric intensive care unit. *J Pediatr Gastroenterol Nutr* 54:620–623
141. Barbosa E, Moreira EA, Goes JE, Faintuch J (1999) Pilot study with a glutamine-supplemented enteral formula in critically ill infants. *Rev Hosp Clin Sao Paulo* 54:21–24
142. Briassoulis G, Filippou O, Hatz E, Papassotiropoulos I, Hatzis T (2005) Early enteral administration of immunonutrition in critically ill children: results of a blinded randomized controlled clinical trial. *Nutrition* 21:799–807
143. Carcillo JA, Dean JM, Holubkov R et al (2012) The randomized comparative pediatric critical illness stress-induced immune suppression (CRISIS) prevention trial. *Pediatr Crit Care Med* 13:165–173
144. Carcillo JA, Dean JM, Holubkov R et al (2017) Interaction between 2 nutraceutical treatments and host immune status in the pediatric critical illness stress-induced immune suppression comparative effectiveness trial. *JPEN J Parenter Enter Nutr* 41:1325–1335
145. Jacobs BR, Nadkarni V, Goldstein B et al (2013) Nutritional immunomodulation in critically ill children with acute lung injury: feasibility and impact on circulating biomarkers. *Pediatr Crit Care Med* 14:e45–56
146. Jordan I, Balaguer M, Esteban ME et al (2016) Glutamine effects on heat shock protein 70 and interleukines 6 and 10: randomized trial of glutamine supplementation versus standard parenteral nutrition in critically ill children. *Clin Nutr* 35:34–40
147. McNally JD, Nama N, O'Hearn K et al (2017) Vitamin D deficiency in critically ill children: a systematic review and meta-analysis. *Crit Care* 21:287
148. Cvijanovich NZ, King JC, Flori HR, Gildengorin G, Vinks AA, Wong HR (2016) Safety and dose escalation study of intravenous zinc supplementation in pediatric critical illness. *JPEN J Parenter Enter Nutr* 40:860–868
149. Horn D, Chaboyer W (2003) Gastric feeding in critically ill children: a randomized controlled trial. *Am J Crit Care* 12:461–468
150. Meert KL, Daphtary KM, Metheny NA (2004) Gastric vs small-bowel feeding in critically ill children receiving mechanical ventilation: a randomized controlled trial. *Chest* 126:872–878
151. Sonmez Duzkaya D, Yildiz S (2016) Effect of two different feeding methods on preventing ventilator associated pneumonia in the pediatric intensive care unit (PICU): a randomised controlled study. *Aust Crit Care* 29:139–145
152. Horn D, Chaboyer W, Schluter PJ (2004) Gastric residual volumes in critically ill pediatric patients: a comparison of feeding regimens. *Aust Crit Care* 17(98–100):2–3
153. Kamat P, Favaloro-Sabatier J, Rogers K, Stockwell JA (2008) Use of methylene blue spectrophotometry to detect subclinical aspiration in enterally fed intubated pediatric patients. *Pediatr Crit Care Med* 9:299–303
154. Tume LN, Bickerdike A, Latten L et al (2017) Routine gastric residual volume measurement and energy target achievement in the PICU: a comparison study. *Eur J Pediatr* 176:1637–1644
155. Gharpure V, Meert KL, Sarnaik AP (2001) Efficacy of erythromycin for postpyloric placement of feeding tubes in critically ill children: a randomized, double-blind, placebo controlled study. *JPEN J Parenter Enter Nutr* 25:160–165
156. van Puffelen E, Hulst JM, Vanhorebeek I et al (2018) Outcomes of delaying parenteral nutrition for 1 week vs initiation within 24 hours among undernourished children in pediatric intensive care: a subanalysis of the PEPaNIC randomized clinical trial. *JAMA Netw Open* 1:e182668
157. Kyle UG, Lucas LA, Mackey G et al (2016) Implementation of nutrition support guidelines may affect energy and protein intake in the pediatric intensive care unit. *J Acad Nutr Diet* 116(844–51):e4
158. Brown A-M, Forbes ML, Vitale VS, Tiroider UH, Zeller R (2012) Effects of a gastric feeding protocol on efficiency of enteral nutrition in critically ill infants and children. *ICAN: Infant Child Adolesc Nutr* 2012;4:175–80.
159. Wong JJ, Ong C, Han WM, Lee JH (2014) Protocol-driven enteral nutrition in critically ill children: a systematic review. *JPEN J Parenter Enter Nutr* 38:29–39
160. Gentles E, Mara J, Diamantidi K et al (2014) Delivery of enteral nutrition after the introduction of practice guidelines and participation of dietitians in pediatric critical care clinical teams. *J Acad Nutr Diet* 114(1974–80):e3
161. Kaufman J, Vichayavilas P, Rannie M et al (2015) Improved nutrition delivery and nutrition status in critically ill children with heart disease. *Pediatrics* 135:e717–e725
162. Geukens V, Neef M, Dijkshof M, Sauerwein H, Bos A (2012) Effect of a nurse-driven feeding algorithm and the institution of a nutritional support team on energy and macronutrient intake in critically ill children. *e-SPEN J* 7(1):e35–e40. <https://doi.org/10.1016/j.eclnm.2011.12.002>