

Support services for immigrant people living with HIV; health care providers` perspective

This chapter has been published in similar form as:

Been SK, Verbon A. Support Services for Immigrant People Living with HIV: Health Care Provider's Perspective. *Journal of AIDS and Clinical Research*. 2018. 9: 752.

ABSTRACT

To evaluate support services for immigrant people living with HIV, a survey was done among HIV care providers of all 27 HIV treatment centers in the Netherlands. Although multiple support services are available, there is a need for additional care in the form of peer support, psychosocial care, and care for refugees and undocumented immigrants in rural areas. With the increase of immigrants in Europe, evaluation of support services in other European countries is recommended.

INTRODUCTION

In Europe, migrants are among the key populations disproportionately affected by HIV (1). This group is a vulnerable population with barriers to optimally access HIV prevention and treatment services, resulting in more late presentation of HIV, less retention in care, and ultimately suboptimal viral suppression rates (2-5). In the Netherlands, more than 40% of people living with HIV (PLWH) is of immigrant origin (6). In 2014, only 70% of the immigrant PLWH (IPLWH) who were diagnosed and linked to care had reached viral suppression (HIV RNA <100copies/ml) compared to 82% of the Dutch PLWH (Figure 1).

The availability of medical care and support services is crucial for adequate HIV treatment. According to UNAIDS, these services are important to, among others, support adherence to treatment, enhance prevention and management of HIV-related infections, and to enhance coping with the challenges of living with HIV (7). Access for IPLWH to these services starts with the awareness of HIV care providers (HCPs) of the available support services. Currently it is unclear which services can be utilized for IPLWH in the HIV treatment centers (HTCs) in the Netherlands, which challenges HCPs encounter, and what is needed to optimize the care for the increasing population of IPLWH in the Netherlands and in Europe. The aim of this study was to evaluate the availability of support services for IPLWH that HCPs working in all Dutch HTCs can refer to.




Figure 1. 'Cascade of Care' in the Netherlands (Stichting HIV Monitoring, 2014)

METHODS

Participants and data collection

Between April and June 2015, a questionnaire was sent to physicians and nurses from all 27 HTCs in the Netherlands. This questionnaire included items about their view on the care and support

services for first and second generation IPLWH offered in both their HTC and nationally. Additionally, HCPs were asked if it was possible to offer additional support services to their patients, like interpreters, legal aid, spiritual aid, health and HIV education, psychological or psychiatric aid, sex counseling, social work, buddy care, and peer support.

Data analyses

Results were analyzed per HTC. If respondents from one HTC gave opposite answers, the result was counted as 'inconclusive' for that center. In addition, a distinction was made between HTCs located in urban and rural areas as refugees are primarily located in the rural areas and immigrants with a residence permit mostly reside in urban areas. Lastly, a comparison was made between HTCs with more and less than 600 PLWH in care.

RESULTS

HCPs from 25 of the 27 HTCs (93%) completed the questionnaire, 13 HTCs were located in urban areas and 12 in rural areas (Table 1). Thirteen HTCs had more than 600 PLWH in care. HCPs often underestimated the percentage of IPLWH in care, they estimated that 33% (mean, range: 2-60) of the PLWH in clinical care were of immigrant origin. The care provided within the HTC and within the Netherlands was graded as 7.7 and 7.1 (mean) out of 10, respectively (Table 1). HCPs of 80% of the HTCs (20 HTCs) could refer IPLWH to support services such as interpreters, psychological or psychiatric aid, social work, buddy care, and peer support. In multiple HCPs, no referrals were possible to services like legal aid (32%, 8 HTCs), spiritual aid (28%, 7 HTCs), health and HIV education (44%, 11 HTCs), and sex counseling (24%, 6 HTCs).

Barriers for appropriate care on a patients' level (e.g., language barriers, cultural differences, psychosocial problems, and no show at outpatient clinic appointments) were experienced in 72% of the HTCs (18 HTCs). In 32% of the HTCs (8 HTCs), barriers were experienced on the level of the HTC (e.g., only medically necessary treatment of uninsured patients, and no social workers available in the HTC). On a regional level (province), barriers were experienced in 24% of the HTCs (6 HTCs) (e.g., lack in overview of available support services). HCPs from 48% of the HTCs (12 HTCs) experienced barriers on a national level (e.g., difficulties in continuum of care after replacement of refugees, and complex regulations for asylum seekers and undocumented IPLWH).

Although it is possible to refer patients to psychological and/or psychiatric aid in all HTCs, HCPs from 5 HTCs indicated a need for additional psychosocial care (Table 1). HCPs from 2 HTCs mentioned that IPLWH had barriers such as disclosure concerns and HIV-related stigma that prevented them from using support services.

Table 1. HIV-treatment centers in the Netherlands

	Total N = 25	Rural HTC's N = 12	Urban HTC's N = 13	≥600 PLWH in care N = 13	<600 PLWH in care N = 12
1. Awarded grade for the care provided for IPLWH within the HTC. (median, min-max)	7.7 (6-9)	7.3 (6-8)	7.8 (6-9)	7.8 (6-9)	7.7 (6-9)
2. Awarded grade for the care provided for IPLWH nationally. ^a (median, min-max)	7.1 (5-8)	6.9 (5-8)	7.3 (5-8)	7.0 (5-8)	7.2 (6-8)
3. Satisfied with the care provided for IPLWH in the HTC? ^a (N, % 'yes')	17 (68%)	8 (67%)	9 (69%)	10 (77%)	7 (58%)
4. Satisfied with the care provided for IPLWH nationally? ^a (N, % 'yes')	14 (56%)	6 (50%)	8 (62%)	8 (62%)	6 (50%)
5. Experienced gaps in care and support services for IPLWH. ^a (N, % 'yes')	12 (48%)	8 (67%)	4 (31%)	5 (39%)	7 (58%)
• Peer support and/or buddy care (N, %)	8 (32%)	6 (50%)	2 (15%)	4 (31%)	4 (33%)
• Psychosocial care (N, %)	5 (20%)	2 (17%)	3 (23%)	2 (15%)	3 (25%)
6. Concerns for care for refugees and undocumented IPLWH. (N, %)	16 (64%)	9 (75%)	7 (54%)	9 (69%)	7 (58%)
7. Sufficient visibility to available support services for IPLWH? ^a (N, % 'yes')	11 (44%)	5 (42%)	6 (46%)	2 (15%)	7 (58%)

IPLWH, immigrant people living with HIV; PLWH, people living with HIV; HTC, HIV-Treatment center; HCP, HIV care providers

^a Inconclusive: Q3, N=4; Q4, N=4; Q5, N=5; Q7, N=5.

Urban versus rural areas

In 67% of the 12 HTC's located in rural areas (N=8), respondents missed specific aspects in available care provided to immigrants, compared to only 31% of the 13 urban area HTC's (N=4) (Table 1). The area of care most frequently missed by HCP's from HTC's in rural areas (50%, N=6) was peer support and/or buddy care. HCP's from 42% of the HTC's located in rural areas (N=5) and 46% of the HTC's located in urban areas (N=6) indicated that they have insufficient insight into the available support services in their region.

Comparison based on the size of the HIV-population

In 58% of the HTC's with <600 patients in care (N=7), respondents missed specific aspects in available care provided to immigrants, compared to 39% of the HTC's with ≥600 patients in care (N=5) (Table 1). HCP's from 58% of the HTC's with <600 patients in care (N=7) and 15% of the HTC's with ≥600 patients in care (N=2) indicated that they have insufficient insight into the available support services in their region.

Asylum seekers and undocumented IPLWH

The most frequently addressed concern on different levels throughout the questionnaire was care for refugees and undocumented IPLWH (Table 1), mentioned by HCP's from 75% of the HTC's located in rural areas (N=9) and 54% of the HTC's located in urban areas (N=7). HCP's indicated that replacements of refugees to other asylum seeker centers throughout the nation is a challenge, hampering adequate HIV care. In addition, only contracted hospitals (15 out of 27 HTC's) are allowed to provide non-emergency HIV-care to refugees and undocumented IPLWH. This may result in loss of patient-provider relationships, and subsequently loss of patients retained in care when patients have to relocate, and higher travel costs for this population with minimal financial resources.

CONCLUSIONS

In conclusion, our data indicate that there is a need for additional support services for IPLWH such as peer support and/or buddy care, psychosocial care, and specifically for refugees and undocumented IPLWH in HTC's in the Netherlands. The need for these support services is higher in rural areas than in urban areas. Peer and/or buddy support projects from urban areas need to be expanded to rural areas. In addition, a reassessment of the current Dutch policy on HIV care for refugees and undocumented IPLWH, and psychosocial care for all IPLWH should take place, since access to HIV care and support services is not only beneficial for the individual patient but also decreases the risk of transmission. With the increase of immigrants in Europe over the past years, it is important for all European countries to evaluate the specific areas of care provided in

and outside HTC's, on a regional and on a national level and these services should be adjusted if needed.

New contribution to the literature

This study evaluated the availability of support services for immigrant people living with HIV in the Netherlands. HIV care providers indicate a need for additional support services for immigrant people living with HIV. This need is higher in rural areas. Policy on care for refugees and undocumented people living with HIV should be reassessed in the Netherlands, but also in other European countries.

Acknowledgements

We thank the HIV care providers of the Dutch HIV treatment centers for their contribution to the questionnaire. Additionally, we thank colleagues from the following organizations that have contributed to the questionnaire: Soa Aids Nederland, Ethnic Minority program, Amsterdam; Dutch HIV Association, Netherlands; ShivA, Amsterdam; Foundation Mara, Rotterdam; Dutch Association of HIV physicians, Netherlands; Dutch Association of HIV-nurses, Netherlands; Erasmus University Medical Center, Rotterdam.

The results from this study have been presented at the following meetings:

- ECCMID 2016, Amsterdam, the Netherlands, April 10th 2016.
- Spring meeting of the Dutch Association of HIV physicians, Utrecht, the Netherlands, June 23rd 2016.

REFERENCES

1. Europe EW. HIV/AIDS Surveillance in Europe, 2015. Stockholm: European Centre for Disease Prevention and Control, 2016. Available from: <http://ecdc.europa.eu/en/publications/Publications/HIV-AIDS-surveillance-Europe-2015.pdf>.
2. Deblonde J, De Koker P, Hamers FF, Fontaine J, Luchters S, Temmerman M. Barriers to HIV testing in Europe: a systematic review. *Eur J Public Health*. 2010;20(4):422-32.
3. Op de Coul EL, Van Sighem A, Brinkman K, Van Benthem BH, Van der Ende ME, Geerlings S, et al. Factors associated with presenting late or with advanced HIV disease in the Netherlands, 1996-2014: results from a national observational cohort. *BMJ Open*. 2016;6(1):e009688.
4. Schouten M, van Velde AJ, Snijdewind IJ, Verbon A, Rijnders BJ, van der Ende ME. [Late diagnosis of HIV positive patients in Rotterdam, the Netherlands: risk factors and missed opportunities] Late diagnose van hiv-patienten in Rotterdam: risicofactoren en gemiste kansen. *Ned Tijdschr Geneeskd*. 2013;157(15):A5731.
5. van Andel E, Been SK, Rokx C, van der Ende ME. Risk factors in an HIV-infected population for refraining from specialist care. *AIDS Care*. 2016;28(10):1255-60.
6. van Sighem A.I. BTS, Wit F.W.N.M., Smit C., Matser A., Reiss P. Monitoring Report 2016. Human Immunodeficiency Virus (HIV) Infection in the Netherlands. Amsterdam: Stichting HIV Monitoring, 2016 2016. Available from: Available online at www.hiv-monitoring.nl.
7. UNAIDS. HIV care and support taking into account the WHO consolidated guidelines. Switzerland: Joint United Nations Programme on HIV/AIDS, 2016. Available from: http://www.unaids.org/sites/default/files/media_asset/JC2741_HIV-care-and-support_en.pdf.