

REVIEW ARTICLE (META-ANALYSIS)

Epidemiology of Cerebral Palsy in Adulthood: A Systematic Review and Meta-analysis of the Most Frequently Studied Outcomes

Marloes van Gorp, PhD,^{a,b} Sander R. Hilberink, PhD,^{a,c} Suzie Noten, MSc,^{a,b}
Joyce L. Benner, PhD,^{a,b} Henk J. Stam, MD, PhD,^a Wilma M.A. van der Slot, MD, PhD,^b
Marij E. Roebroek, PhD^{a,b}

From the ^aDepartment of Rehabilitation Medicine, Erasmus MC University Medical Center, Rotterdam, The Netherlands; ^bRijndam Rehabilitation, Rotterdam, The Netherlands; and ^cResearch Centre Innovations in Care, Rotterdam University of Applied Sciences, Rotterdam, The Netherlands.

Abstract

Objective: To describe the epidemiology of health status, impairments, activities and participation in adults with cerebral palsy (CP).

Data Sources: Embase, MEDLINE, Web of Science, PsycINFO, Cumulative Index to Nursing and Allied Health, Cochrane, and Google Scholar were searched for 3 themes (“cerebral palsy,” “adult,” and “outcome assessment”) in literature published between January 2000 and December 2018.

Study Selection: Full-article peer-reviewed English journal articles on descriptive, observational, or experimental studies describing the most studied outcomes in adults with CP ($n \geq 25$, age ≥ 18 y) were included. Studies were included in the analyses if frequently studied outcomes were described in at least 3 studies using similar methods of assessment.

Data Extraction: Data were extracted independently by 2 authors from 65 articles (total $N = 28,429$) using a standardized score sheet.

Data Synthesis: Meta-analyses revealed that overall, on average 65.1% (95% confidence interval [CI], 55.1-74.5) of adults with CP experienced pain, 57.9% (95% CI, 51.1-64.6) were ambulant, 65.5% (95% CI, 61.2-69.7) had little or no limitation in manual ability, 18.2% (95% CI, 10.6-27.2) had tertiary education, 39.2% (95% CI, 31.5-47.1) were employed, and 29.3% (95% CI, 9.0-55.3) lived independently. In adults without intellectual disability, proportions of individuals who were ambulant (72.6% [95% CI, 58.8-84.5]) and lived independently (90.0% [95% CI, 83.8-94.9]) were higher ($P = .014$ and $P < .01$, respectively). The Fatigue Severity Scale score was 4.1 (95% CI, 3.8-4.4). Epilepsy (28.8% [95% CI, 20.1-38.4]) and asthma (28.3% [95% CI, 18.7-38.9]) were especially prevalent comorbidities.

Conclusions: The present systematic review and meta-analysis on the epidemiology of adults with CP provided state-of-the-art knowledge on the most frequently studied outcomes. On average, adults with CP are fatigued, and a majority experience pain, are ambulant, and have little or no difficulty with manual ability. On average, 40% are employed and 30% live independently. More uniformity in assessment and reports is advised to improve knowledge on epidemiology and gain insight in more outcomes.

Archives of Physical Medicine and Rehabilitation 2020; ■: ■ ■ ■ ■ - ■ ■ ■ ■

© 2020 by the American Congress of Rehabilitation Medicine

Cerebral palsy (CP) is an umbrella-term describing “a group of permanent disorders of the development of movement and posture, causing activity limitation, that are attributed to nonprogressive disturbances that occurred in the developing fetal or infant brain. The motor disorders of cerebral palsy are often

accompanied by disturbances of sensation, perception, cognition, communication and behavior, by epilepsy and by secondary musculoskeletal problems”^{1(p9)} and it affects 2 to 2.5 per 1000 live births.¹⁻⁴ Approximately three-fourths of persons with CP are adults (ie, 18 years or older).⁵

In the past decades, attention for adults with CP in rehabilitation practice and research increased.⁶ Aging with CP comes with several issues in health and functioning. Recently, several comorbidities were increased in adults with CP, including asthma

Supported by Rijndam Rehabilitation, Rotterdam, The Netherlands.
Disclosures: none

and hypertension.^{7,8} Pain and fatigue are common in adults with CP,⁹ and pain, mobility, and self-care deteriorate over time.^{10,11} In addition, young adults with CP are less often employed or live independently than young adults without CP.^{12,13} Moreover, activity limitations and difficulty in participation increase with aging.^{10,14}

Common impairments and functional limitations in children with CP were previously studied in a systematic review,¹⁵ but systematic knowledge on adults with CP is scarce. Because differences were found in epidemiology between children and adults with CP,¹⁶ results cannot be extrapolated to the adult CP population. Insight into the prevalence of impairments and level of functioning in adults with CP would be helpful to guide rehabilitation follow-up programs. In addition, such insight would contribute to a better understanding of the effect of CP at adult age. Therefore, this review and meta-analysis aims to describe the health status, prevalence of impairments, and level of activities and participation of adults with CP using published literature. We focused on the most frequently studied outcomes in adults with CP from the literature, which were reported in a recently published review by Benner et al.⁶

Methods

Search strategy

The search strategy, which was developed in consultation with an information specialist, was used in 7 databases (Embase, MEDLINE, Web of Science, PsycINFO, Cumulative Index to Nursing and Allied Health, Cochrane, and Google Scholar) and consisted of 3 themes (“cerebral palsy,” “adult,” and “outcome assessment”). Key terms were mapped to controlled headings and expanded to include free text terms, tailored to the specific database. See Benner et al for the complete search strategy.⁶ Literature was searched from 2000 through 2018 because the search used in Benner et al was updated with publications from 2017 and 2018. After deduplication, all publications were screened for eligibility on title and abstract and subsequently full text by 2 independent reviewers (J.B., S.N. for publications in 2000-2016; M.v.G., S.H. for 2017-2018), disagreements were discussed and resolved between the 2 reviewers. If disagreement remained this was discussed with a third reviewer (M.R.) to reach consensus.

Eligibility criteria

Studies were included if they met the following criteria: (1) They reported on the most frequently studied outcomes according to the systematic review of Benner et al.⁶ From this list of outcomes

single-item outcomes (eg, pain) were selected if at least 10 studies provided information on this outcome; multi-item outcomes (eg, Fatigue Severity Scale [FSS]) were selected if at least 5 studies provided information. (2) At least 3 studies reported on an outcome using similar methods of assessment, enabling comparison and meta-analysis. (3) Study design was descriptive, observational, or experimental. Meta-analyses, reviews, case studies, qualitative studies, comments, and study protocols were excluded. (4) They were full-article peer-reviewed journal articles written in English. (5) They described individuals with CP only or in case-control design. (6) They described ≥ 25 adults who were 18 years or older at the time of the first measurement or at follow-up. Studies describing both children and adults with CP were included when at least 50% of the sample were adults. (7) They reported on outcomes of functioning, excluding studies reporting on evaluations of services, for example, evaluation of transition services or complications and adverse events during or after surgery. (8) Study samples were population based or rehabilitation based, or samples addressed a substantial subgroup of individuals with CP (eg, Gross Motor Function Classification System [GMFCS] I-III, bilateral spastic CP, dyskinetic CP). Nonrepresentative samples that consisted of adults selected because of having specific impairments (eg, pain, fatigue, incontinence) or because of previous operations (eg, selective dorsal rhizotomy, constructive hip operations) were excluded. For intervention studies, only baseline observations were included. When outcomes were reported in multiple publications on the same study sample, the most recently observed outcome was included.

Outcomes

Frequently studied outcomes identified by Benner et al were grouped in line with the International Classification of Functioning, Disability, and Health (ICF) as health status, impairments, or activities and participation, with the first group encompassing outcomes that were not covered by the ICF or were linked to more than 1 component.^{6,17} These outcomes are depicted in the first column of [Table 1](#). Health status outcomes addressed weight status (body mass index [BMI, calculated as weight in kilograms divided by height in meters squared], obesity), fat mass (waist circumference), comorbidities, and the domains of the Short Form-36 Health Survey (SF-36) (scale range 0-100, with the physical health component score on a T score metric with a mean of 50 and SD of 10 for the US general population).⁷⁸ Comorbidities included epilepsy, diabetes, hypertension, asthma, stroke, heart disease, and osteoarthritis. The impairments addressed pain (presence of any pain, including the outcomes pain in joints and pain in body part⁶), fatigue (assessed using the FSS, scale range 1-7)⁷⁹ and mobility of joints (limited range of motion in knee or hip). Outcomes referring to the level of activities and participation were walking status (ambulant with or without assistive device as primary method of mobility in daily life), which included the outcomes walking or moving around using assistive devices and assistive devices for mobility⁶, fine hand use (Manual Ability Classification System [MACS] level I or II, manual ability),⁸⁰ education status (attending or completing tertiary education according to the international standard classification of education; ie, higher education, college and university),⁸¹ employment status (having remunerative employment), living status (living independently), and activities and social roles, assessed using the Assessment of Life Habits questionnaire, Life-H (scale range 0-9).⁸² Although frequently studied according to Benner et al,⁶ we

List of abbreviations:

BMI	body mass index
CI	confidence interval
CP	cerebral palsy
FSS	Fatigue Severity Scale
GMFCS	Gross Motor Function Classification System
ICF	International Classification of Functioning, Disability and Health
ID	intellectual disability
MACS	Manual Ability Classification System
SF-36	Short Form-36 Health Survey

Table 1 Characteristics of included studies

Author*	Country [†]	N	ID (%)	GMFCS I-III (%)	Age (y), Mean (Range)	% Male
Ballester-Plané et al ¹⁸	Spain ^a	52	Mixed	56	25 (7-62)	54
Laporta-Hoyos et al ¹⁹	Spain ^a	39	Mixed	64	21 (6-62)	51
Whitney et al ²⁰	USA ^b	1395	Mixed	54	18 to >50	48
Whitney et al ⁸	USA ^b	452	Mixed	51	24 (18-30)	43
Cremer et al ²¹	USA ^b	435	Mixed	54	50 (40-60)	46
Cremer et al a ²¹	USA ^b	201	Mixed	55	48 (40-60)	100
Cremer et al b ²¹	USA ^b	234	Mixed	53	50 (40-60)	0
Ágústsson et al ²²	Sweden ^c	830	Mixed	57	23 (16-73)	56
Alriksson-Schmidt et al ²³	Sweden ^c	102	Mixed	72	21 (18-24)	62
Rodby-Bousquet et al ²⁴	Sweden ^c	102	Mixed	72	21 (19-23)	62
van Gorp et al a ²⁵	The Netherlands ^d	67	31	67	25 (21-27)	67
van Gorp et al b ²⁵	The Netherlands ^d	54	0	82	32 (29-34)	54
van Meeteren et al ²⁶	The Netherlands ^d	83	0	87 [‡]	20 (18-22)	59
Nieuwenhuijsen et al ²⁷	The Netherlands ^d	87	0	90	20 (18-22)	59
Benner et al ¹⁰	The Netherlands ^e	49	22	80	40 (34-45)	55
Benner et al ¹²	The Netherlands ^e	49	22	80	40 (34-45)	55
van der Dussen et al ²⁸	The Netherlands ^e	80	28	71 [§]	21-31	51
Kirk et al a ²⁹	Denmark ^f	22	Mixed	100	34 (18-57)	68
Kirk et al b ²⁹	Denmark ^f	22	Mixed	100	37 (18-59)	55
McPhee et al ³⁰	Canada ^g	42	Mixed	57	34 (18-75)	50
McPhee et al ³¹	Canada ^g	42	Mixed	57	34 (18-75)	50
Slaman et al a ³²	The Netherlands ^h	28	0	100	20 (16-24)	43
Slaman et al b ³²	The Netherlands ^h	29	0	98	20 (16-24)	52
Russchen et al ³³	The Netherlands ^h	56	0	100	20 (16-24)	48
Slaman et al ³⁴	The Netherlands ^h	57	0	98	20 (16-24)	47
Ryan et al ³⁵	Ireland	55	Mixed	75	38 (18-65)	56
Morgan et al ³⁶	Australia	34	0	100	44 (26-65)	44
Morgan et al ³⁷	Australia	25	16	100	41 (30-65)	36
Reddihough et al ³⁸	Australia	335	50	63	25 (20-30)	51
van der Slot et al ³⁹	The Netherlands ⁱ	43	0	95	37 (25-45)	63
van der Slot et al a ³⁹	The Netherlands ⁱ	27	0	95	37 (25-45)	100
van der Slot et al b ³⁹	The Netherlands ⁱ	16	0	95	37 (25-45)	0
van der Slot et al ⁴⁰	The Netherlands ⁱ	56	0	93	36 (25-45)	63
van der Slot et al ⁴¹	The Netherlands ⁱ	56	0	93	36 (25-45)	63
van der Slot et al a ⁴¹	The Netherlands ⁱ	35	0	93	36 (25-45)	100
van der Slot et al b ⁴¹	The Netherlands ⁱ	21	0	93	36 (25-45)	0
Opheim et al ⁴²	Norway ^j	149	0	93	40 (24-76)	51
Opheim et al ¹¹	Norway ^j	149	0	93	40 (24-76)	51
Jahnsen et al ⁴³	Norway ^j	406	0	78 [§]	34 (18-72)	51
Jahnsen et al ⁴⁴	Norway ^j	406	0	78 [§]	34 (18-72)	51
Jahnsen et al ⁴⁵	Norway ^j	766	0	Mixed	34 (18-72)	52
Maltais et al ⁴⁶	Canada	132	Mixed	60	28 (20-41)	50
Michelsen et al ⁴⁷	Denmark ^k	416	20	88 [§]	32 (29-35)	58
Michelsen et al ⁴⁸	Denmark ^k	819	20	84 [§]	29 (21-35)	58
Engel et al ⁴⁹	USA	100	0	18	38 (19-71)	55
Fortuna et al ⁷	USA	229	54	55 [§]	18 to >60	59
Gillet et al ⁵⁰	Australia	33	Mixed	100 [‡]	25 (15-51)	55
Lundh et al ⁵¹	Sweden	50	Mixed	100	32 (22-67)	54
Park et al ⁵²	USA	229	0	51	40 (20-69)	60
Peterson et al ⁵³	USA	2659	Mixed	Mixed	Median (IQR): 36 (25-48)	52
Sienko et al ⁵⁴	USA	97	41 [¶]	65	24 (18-30)	48
Smith et al ⁵⁵	UK	1705	21	Mixed	33±16	53
Daunter et al ⁵⁶	USA	50	0	70	27 (18-35)	48
de Albuquerque Botura et al ⁵⁷	Brazil	93	Mixed	0	18-57	56
Hayward et al ⁵⁸	USA	375	24 [¶]	63	36 (18-76)	0
Park et al ⁵⁹	Korea	53	Mixed	63 [‡]	31±14	62
Vukojevic et al ⁶⁰	Bosnia and Herzegovina	100	55	Mixed	18-58	62

(continued on next page)

Table 1 (continued)

Author*	Country†	N	ID (%)	GMFCS I-III (%)	Age (y), Mean (Range)	% Male
Yildiz et al ⁶¹	Turkey	117	26 [#]	74	25 (18-62)	55
Brunton et al ⁶²	Canada	111	0	78	19 (14-31)	50
Peterson et al ⁶³	USA	1015	Mixed	Mixed	58 (57-60)	66
Peterson et al ⁶⁴	USA	112	Mixed	52	34±13	46
Huang et al ⁶⁵	Taiwan	279	28	50 [§]	26 (20-40)	61
Nedjad et al ⁶⁶	Sweden	156	62 [¶]	27 [§]	37 (19-43)	55
Mesterman et al ⁶⁷	Israel	95	35	83 [§]	23 (18-30)	61
Gaskin et al ⁶⁸	Australia	51	0	45	38 (19-66)	63
Andersson et al a ⁶⁹	Sweden	13	0	100	36 (26-58)	64
Andersson et al b ⁶⁹	Sweden	12	0	100	36 (26-58)	64
Sandstrom et al ⁷⁰	Sweden	48	20 [¶]	71	33 (20-?)	48
Strauss et al ⁷¹	USA	14,806	Mixed	62 [§]	20-85	53
Andersson et al ⁷²	Sweden	221	0	Mixed	36 (20-58)	57
Bottos et al ⁷³	Italy	72	41	76 [§]	33 (19-65)	60
Furukawa et al ⁷⁴	Japan	81	Mixed	73 [§]	36 (26-57)	44
Hodgkinson et al ⁷⁵	France	234	Mixed	0	28 (15-?)	59
Maruishi et al ⁷⁶	Japan	256	43	Mixed	32 (17-83)	55
Murphy et al ⁷⁷	USA	101	Mixed	Mixed	43 (19-74)	52

NOTE. Mixed: no exact proportion reported, also no in- or exclusion criteria regarding cognition or mobility.

Abbreviation: IQR, interquartile range.

* Letters a or b after the authors name (ie, Cremer et al a) indicate subsamples of 1 publication.

† Letters indicate publications on the same study sample in a country.

‡ Proportion of GMFCS level I and II.

§ No GMFCS level reported; proportion walking with or without walking aid.

|| Those unable to complete questionnaires were excluded.

¶ Proportion proxy report.

Proportion illiterate, but individuals with severe cognitive impairments were excluded.

did not estimate overall results for the following outcomes because these were only reported in 1 or 2 included studies using a similar assessment method: spasticity, muscle power, hip displacement, gait pattern, speech, spinal deformities of thoracic, lumbar and sacral column, Barthel Index,⁸³ FIM,⁸⁴ Gross Motor Function Measure,⁸⁵ Functional Mobility Scale,⁸⁶ and Japanese Orthopaedic Association Score.⁸⁷

Data extraction

A standardized data extraction record sheet was used to collect study sample characteristics and outcomes. Sample characteristics included country, sample size, sex, age, proportion with intellectual disability (ID), and proportion with level of GMFCS I-III.⁸⁸ Studies including any proportion of individuals with ID or those that did not mention exclusion based on intellectual function, suggesting individuals with ID could be part of the sample, were reported as mixed samples. Studies including only individuals without ID are further referred to as no ID samples. With respect to the outcomes, the method of assessment and the outcome (proportion [%], or mean/median scores [SD/interquartile range]) were noted.

Analysis

Overall results with corresponding 95% confidence intervals (CIs) were estimated using meta-analysis models. Overall mean proportions and means were estimated with a random-effects meta-analysis model using the DerSimonian and Laird estimator.⁸⁹ The

Freeman-Tukey double arcsine transformation was used on proportions.^{90,91} Means were analyzed untransformed. Meta-analysis modeling was done using the metaprop and metamean functions of the meta package in R 3.2.5.⁹² The random-effects model takes the heterogeneity of samples into account. Statistical heterogeneity was quantified using the I^2 measure, which describes the amount of variation attributed to heterogeneity rather than sampling error across samples.⁹² Subgroup analyses of mixed and no ID samples were conducted, if multiple studies of mixed and no ID samples were available, for outcomes on activities and participation because ID is known to be strongly related to these outcomes.^{25,93} A random-effect Q test was used to test differences between mixed and no ID samples. Sensitivity analyses excluded specific subsamples of the population with CP from analyses if substantial deviations in the outcome could be explained by sample characteristics. These samples were for manual ability: a no ID sample²⁵ and a sample with only individuals with dyskinetic subtype,¹⁹ for epilepsy: a no ID sample⁵² and a sample with only individuals with GMFCS level IV and V,⁵⁷ and for living situation: a young sample (mean age, 20y).²⁷ Results of sensitivity analyses showed substantial influence of deviating samples, which were therefore excluded in the estimation of overall proportions and means; outcomes of the excluded studies were displayed in the forest plots.

Results

After deduplication, the initial search revealed 6662 publications (fig 1). After screening by title and abstract, 810 full-text articles




Fig 1 Flow diagram of study selection process.

were assessed for eligibility. Finally, 65 articles (total $N = 28,429$) were included in the meta-analyses, meaning these reported 1 or more of the selected outcomes comparable to at least 2 other studies. Table 1 displays the characteristics of the studies used in the meta-analyses. Of the samples, 73% were mixed and 27% were no ID. The mean age of the study samples was 32 years, samples consisted on average of 68% individuals with GMFCS level I-III and 53% were male. Table 2 shows the overall proportion and/or mean for each of the outcomes, and fig 2 and supplemental appendix S1 (available online only at <http://www.archives-pmr.org/>) show the forest plots of these analyses. The level of heterogeneity (I^2) was substantial ($>75\%$) for most of the analyses, reflecting considerable variation in results between studies (see fig 2, supplemental appendix S1).

Health status

Mean overall BMI was 25.1 (95% CI, 23.8-26.5), and 22.6% (95% CI, 14.7-31.5) of individuals with CP were obese (BMI >30). Epilepsy (28.8% [95% CI, 20.0-38.4]), asthma (28.3% [95% CI, 18.7-38.9]), and hypertension (21.6% [95% CI: 15.6; 28.4]) were the most prevalent comorbidities. The overall estimated mean SF-36 physical component score was 44.6 (95% CI, 40.7-48.5).

Impairments

Overall presence of pain was estimated at 65.1% (95% CI, 55.1-74.5). For fatigue, the overall estimated mean FSS score was 4.1 (95% CI, 3.8-4.4). Limitations in knee and hip mobility were estimated to be present in 39.5% (95% CI, 21.1-59.5) and 33.4% (95% CI, 15.1-54.6) of adults with CP, respectively.

Level of activities and participation

A majority of adults with CP in mixed samples were estimated to be ambulatory (57.9% [95% CI, 51.1-64.6]) and to have little or no limitation in manual ability (MACS level I or II, 65.5% [95% CI, 61.2-69.7]). The proportion of ambulatory individuals seemed

higher in the 2 no ID samples (72.6% [95% CI, 58.8-84.5], $Q: 3.60$, $P = .058$). Analyses on additional outcomes of those ambulatory included all samples, regardless of ID. Among ambulatory individuals, 31.9% (95% CI, 22.2-42.4) used an assistive device, and 56.2% (95% CI, 37.0-74.5) had reported to perceive a decline in walking function or capacity over time. The overall mean results on the 6-minute walk test of ambulatory adults with CP was 387.9 m (95% CI, 328.5-447.3).

Reports of attendance of tertiary education were remarkably similar between mixed samples and no ID samples ($Q: 1.03$, $P = 0.311$, overall result: 18.2% [95% CI, 10.6-27.2]). The mean estimated proportion of adults with CP with remunerative employment tended to be lower in mixed samples (39.2% [95% CI, 31.5-47.1]) than in no ID samples (56.0% [95% CI, 31.7-78.9], $Q: 1.64$, $P = .200$). A large difference ($Q: 22.11$, $P < 0.01$) was found in the proportions living independently between mixed samples (29.3% [95% CI, 9.0; 55.3]) and no ID samples (90.0% [95% CI, 83.8; 94.8]). Only used in no ID samples, the overall estimated mean Life-H activities score was 8.0 (95% CI, 7.5-8.4), and the social roles score was 8.2 (95% CI, 7.9-8.4).

Discussion

This systematic review and meta-analysis estimated overall health status, and rates of impairments, activities, and participation of adults with CP for the most studied outcomes in the available literature according to a recent systematic review by Benner et al.⁶ This provides state-of-the-art knowledge for this population on a range of topics, such as pain, walking status, living situation, and comorbidities based on large accumulated samples combined from studies across the globe.

Health status

Body composition and comorbidities that are known to be associated with lifestyle, such as obesity, hypertension, and diabetes, vary strongly across regions and age. The overall proportions estimated in the present review are therefore difficult to compare with general population data irrespective of age. Still, the overall estimates that we provided for adults with CP provide a robust indication of the occurrence of several comorbidities in the CP adult population. Epilepsy is a known comorbidity of CP and is included in its definition.¹ The overall occurrence of 28.8% is similar to that reported in a previous review of adults with CP and the more recent review in children with CP.^{2,15} In addition, the overall estimated occurrence of asthma in adults with CP (28.3%) exceeds prevalence rates that were reported across the globe in general populations (ranging from 1.0%-21.5%),⁹⁴ suggesting that asthma may be a common comorbidity in CP.

Impairments

Pain is the most frequently studied outcome in adults with CP.⁶ The estimated occurrence of pain (65%) is in line with another recent review in adults with CP focusing on pain using individual patient data (70%; 95% CI, 62-78)⁹⁵ and seems to be more prevalent compared with the general US population (56%).⁹⁶ For fatigue we estimated an overall score on the FSS over 4.0, which is often used as a cutoff score indicating fatigue, implying that on average adults with CP experience fatigue.⁹⁷ A Swiss study found a similar mean score for patients with a recent stroke and a lower mean score (3.0) for healthy subjects without CP.⁹⁷ Prevention and

Table 2 Meta-analyses results

Outcome	Studies (n) (Reference nos.)	Cases in Analysis (N)	Overall Proportion or Mean (95% CI)
Health status			
Body composition			
Waist circumference (mean, in cm)	5 (31,34,35,39,64)	266	84.0 (82.2-85.7)
BMI (mean)	6 (21,31,33,35,39,64)	743	25.1 (23.8-26.5)
Obese (% BMI \geq 30)	7 (7,21,30,35,39,59,64)	968	22.6% (14.7-31.5)
Comorbidities (%)			
Epilepsy	6 (7,10,18,55,60,73)	2189	28.8% (20.0-38.4)
Diabetes	7 (7,8,30,39,52,53,55,63)	6594	6.7% (3.8-10.3)
Hypertension	8 (7,8,30,39,52,53,59,63)	4646	21.6% (15.6-28.4)
Asthma	4 (7,8,21,63)	2076	28.3% (18.7-38.9)
Stroke	3 (8,21,63)	1847	2.0% (0.0-6.8)
Heart disease	3 (55,63,73)	2792	9.5% (5.1-15.0)
Osteoarthritis	3 (8,21,55)	2537	11.4% (2.0-26.9)
SF-36 (mean)			
Samples without ID			
Physical component	3 (36,41,42)	239	44.6 (40.7-48.5)
Physical function	4 (32,36,41,68)	197	53.9 (34.0-73.7)
Role physical	4 (32,36,41,68)	197	72.6 (67.7-77.6)
Role emotional	4 (32,36,41,68)	197	82.7 (77.9-87.5)
Vitality	4 (32,36,41,68)	197	56.9 (53.3-60.5)
Mental health	4 (32,36,41,68)	197	73.8 (71.6-76.0)
Social functioning	4 (32,36,41,68)	197	80.6 (76.5-84.6)
Bodily pain	5 (32,36,41,44,68)	599	70.3 (63.5-77.2)
General health	4 (32,36,41,68)	197	70.6 (67.2-73.9)
Impairments			
Pain (%)			
Presence of any pain	16 (10,24,40,44,49,51,54,56,57,59,62,66,70,72,74,76)	1836	65.1% (55.1-74.5)
Fatigue (mean)			
Fatigue Severity Scale	5 (11,33,40,51,56)	352	4.1 (3.8-4.4)
Mobility of joints (%)			
Limited knee mobility	3 (22,51,70)	928	39.5% (21.1-59.5)
Limited hip mobility	3 (22,51,70)	928	33.4% (15.1-54.6)
Activities and participation			
Walking status			
Ambulatory (%) - subgroup analysis:			
Mixed samples	9 (7,10,46,65,67,71,73,74,77)	15839	57.9% (51.1-64.6)
Samples without ID	2 (43,72)	627	72.6% (58.8-84.5)
Ambulatory with assistive devices of those ambulatory (%)	5 (43,65,72-74)	689	31.9% (22.2-42.4)
Perceived walking decline (%)	4 (11,36,37,46)	293	56.2% (37.0-74.5)
6MWT (mean, in m)	4 (29,37,50,69)	127	387.9 (328.5-447.3)
Manual ability (%)			
No or little limitation (MACS level I or II)	3 (23,25,58)	485	65.4% (61.1-69.6)
Education (%)			
Tertiary education (all samples)	8 (28,38,45,48,65,72,73,77)	2635	18.2% (10.6-27.2)
Employment (%)			
Remunerative employment - subgroup analysis:			
Mixed samples	9 (12,23,38,48,65,67,73,74,77)	1845	39.2% (31.5-47.1)
Samples without ID	4 (11,27,41,72)	437	56.0% (31.7-78.9)
Living status (%)			
Living independently - subgroup analysis:			
Mixed samples	7 (23,38,47,67,71,73,77)	15927	29.3% (9.0-55.3)
Samples without ID	3 (11,41,72)	426	90.0% (83.8-94.8)

(continued on next page)

Table 2 (continued)

Outcome	Studies (n) (Reference nos.)	Cases in Analysis (N)	Overall Proportion or Mean (95% CI)
Activities and social roles (mean score Life-H)			
Activities domain			
Samples without ID	3 (^{26,32,41})	198	8.0 (7.5-8.4)
Social roles domain			
Samples without ID	3 (^{26,32,41})	198	8.2 (7.9-8.4)

Abbreviation: 6MWT, 6-minute walk test.

treatment of pain and fatigue are high priorities on a recent patient-centered research agenda⁹⁸ because literature on pain and fatigue management and lifestyle strategies for this population is scarce.^{32,99,100} Using insights from previous studies as a starting point, such as discussing pain and fatigue management with adults with CP and applying a multidimensional lifestyle intervention program, future studies should aim to determine the effectiveness of interventions reducing pain and fatigue.

We found that 1 in 3 adults with CP experience limited knee and hip mobility. However, variation of these proportions were very large, and few population-based studies were available that reported joint mobility in adults with CP. Systematic knowledge of joint mobility seems to be under-reported in current literature. Similarly, other impairments that frequently occur in individuals with CP, such as spasticity, hip displacement and spinal deformities, were not often reported on in population-representative samples or lacked standardized assessments.^{2,15,22}

Activities and participation

A majority of adults with CP were ambulatory (58%) and experienced little or no difficulty with manual ability (65%). However, the proportion of ambulant individuals is lower than that previously reported in children with CP (72%).¹⁵ This is likely explained by a decline in walking function as individuals with CP get older, which we found to be experienced by half of the ambulant adults with CP. The proportion of adults with CP who experienced little or no limitation with manual ability (65%) was very close to that reported in a population-based study of children with CP (64%).¹⁰¹ This suggests that although walking ability declines in individuals with CP, manual ability remains relatively stable in adulthood.

The overall proportions of adults with tertiary education (18%), having remunerative employment (39%), and living independently (29%) seem low but are very difficult to compare with reference values because of regional differences and age-specific participation rates, for example. Additionally, the variation between CP samples is also large, as shown by large statistical heterogeneity (92%-99%), a very large 95% CI for living situation, and substantially higher proportions of participation in samples without adults with ID. Importantly, in individuals without ID, proportions of living independently were very high (90%). Surprisingly, regarding tertiary education and remunerative employment there were no significant differences between mixed samples and samples without ID, which may indicate that adults with CP without ID may still be disadvantaged in these life areas. Although the sample from the study of Murphy et al used in the meta-analyses of the outcomes walking, education, employment status, and living situation was not reported to be selected based

on intellectual disability (ie, it was considered a mixed sample), the results were deviant.⁷⁷ This might be explained by a large proportion of the sample classified as dyskinetic (50%), which usually represent smaller parts of the population.² Individuals with dyskinetic CP may have different levels of participation than those with spastic CP, causing results of this sample to differ.

Overall scores on daily activities and social roles (Life-H) were close to the cutoff for experiencing difficulty.^{102,103} These scores are only estimated based on Dutch samples without ID, which may limit generalization. Similarly, overall domain scores for physical functioning and role limitations because of physical health (SF-36) were low compared with reference values reflecting limitations in activities and participation.¹⁰⁴

Heterogeneity in samples and studies

The high level of statistical heterogeneity, indicating substantial variation between study results, and the wide confidence intervals in some of the analyses may partly be explained by sample differences. These differences in samples reflect the heterogeneity of the population with CP, of which the definition is an umbrella term.¹ Previous studies found associations between CP severity and many of the described outcomes.^{8,12,14,95} Because of the available information, we were unable to disentangle differences in outcomes for other determinants (eg, level of motor function, age, or region) than mixed samples and samples without ID. We recommend future studies to systematically report overall results for subgroups of adults with CP to improve clinical interpretation. Moreover, we strongly recommend to further standardize assessments in future research and clinical care for adults with CP. Examples of presently ongoing standardization in this field are improving the uniformity of registries^{105,106} and extending them into the adult population,²³ developing core outcome sets for specific outcomes,¹⁰⁷ and creating an overarching ICF core set for adults with CP.⁶

Study limitations

The literature search we conducted was very comprehensive, with 386 English-language publications on adults with CP reporting outcomes of functioning identified by screening more than 6000 records from the database searches. However, some publication bias may be present because we omitted non-English language publications, which comprised less than 10% of the eligible publications. Another source of publication bias might be inconclusive trials that were not published. We expect this source of publication bias to be negligible because the outcomes we studied were non-comparative,¹⁰⁸ and intervention trials in adults with CP are very




A**B****C**


Fig 2 Forest plots of main outcomes of activities and participation. (A) Walking status; (B) manual ability; (C) education status; (D) employment status; (E) living situation.

D
Study**Population = Mixed**

Study	Events	Total
Michelsen 2005	279	778
Reddihough 2013	121	335
Huang 2013	64	279
Murphy 2000	71	101
Mesterman 2010	36	95
Furukawa 2001	36	81
Alriksson-Schmidt 2014	20	65
Bottos 2001	21	62
Benner 2017 2	22	49
Random effects model	1845	

Heterogeneity: $I^2 = 90\%$, $\tau^2 = 0.0124$, $P < .01$ **Population = No ID**

Study	Events	Total
Andersson 2001	146	204
Opheim 2009	43	145
van der Slot 2010	38	54
Nieuwenhuijsen 2009	17	34
Random effects model	437	

Heterogeneity: $I^2 = 96\%$, $\tau^2 = 0.0571$, $P < .01$ **Random effects model**Heterogeneity: $I^2 = 94\%$, $\tau^2 = 0.0251$, $P < .01$
Residual heterogeneity: $I^2 = 92\%$, $P < .01$ **E**
Study**Population = Mixed**

Study	Events	Total
Strauss et al ⁷¹ 2004	991	14806
Michelsen 2006	283	416
Reddihough 2013	70	335
Alriksson-Schmidt 2014	29	102
Murphy 2000	67	101
Mesterman 2010	16	95
Bottos 2001	9	72
Random effects model	15927	

Heterogeneity: $I^2 = 99\%$, $\tau^2 = 0.1249$, $P < .01$ **Population = No ID**

Study	Events	Total
Andersson 2001	190	221
Opheim 2009	140	149
Nieuwenhuijsen 2009	19	87
van der Slot 2010	50	56
Random effects model	513	


Heterogeneity: $I^2 = 68\%$, $\tau^2 = 0.0042$, $P = .04$ **Random effects model**Heterogeneity: $I^2 = 100\%$, $\tau^2 = 0.2182$, $P = 0$
Residual heterogeneity: $I^2 = 99\%$, $P < .01$ 

Fig 2 Continued.

scarce, addressing only 5% of the included studies.⁶ Notably, most of the included studies were conducted in North America, Western Europe, Asia, or Australia (see table 1), limiting generalizability of the results to other regions, for example, those with lower resources. Because of limited availability of population-based studies (eg, registry studies)¹⁵ in adults with CP compared with children, we were unable to restrict the selection of articles to population-based samples, causing substantial variability in severity of

motor and intellectual disability between samples (see table 1). However, the total of all study samples included is considered representative for the population of adults with CP with the majority male and with GMFCS I-III and including both mixed and no ID samples. The mean age of samples was 32 years, indicating the included articles mainly studied younger adults, and the results may not be applicable to older adults with CP. Note that we excluded nonrepresentative samples that were specifically selected

in original studies on having a certain impairment. Still, by including all representative study samples with at least 25 adults with CP who were relatively representative of (a subgroup of) the population with CP, we provided the best available knowledge on the most often studied outcomes in adults with CP. Another limitation of the study may be that we only focused on results that were reported as outcomes in the original studies, in line with the review on most studied outcomes of Benner et al,⁶ thus not including information that was originally reported on as sample characteristics, such as level of gross motor functioning, educational level, or employment status.

Conclusions

The present systematic review and meta-analysis provided state-of-the-art knowledge of the epidemiology of the most studied outcomes of health status, impairments, activities, and participation in adults with CP. Epilepsy and asthma are comorbidities that occur often in adults with CP. Overall, a majority of adults with CP experience pain, and on average adults with CP are fatigued. A majority of adults with CP are ambulant and have little or no difficulty in manual ability. A minority of the total population are employed or live independently, with much higher proportions living independently for individuals without ID. More uniformity in assessment and reporting is advised to enable more specified data synthesis on a wider range of outcomes.

Supplier

a. R 3.2.5; R Foundation for Statistical Computing.

Keywords

Adult; Cerebral palsy; Epidemiology; Health; Meta-analysis; Rehabilitation; Systematic review

Corresponding author

Marij E. Roebroek, PhD, Erasmus MC University Medical Center, Dept of Rehabilitation Medicine, PO Box 2040, 3000 CA Rotterdam, The Netherlands. *E-mail address:* m.roebroek@erasmusmc.nl.

Acknowledgment

We thank Wichor Bramer, PhD, for Rijndam Rehabilitation, Rotterdam his support in developing and conducting the literature search.

References

1. Rosenbaum P, Paneth N, Leviton A, et al. A report: the definition and classification of cerebral palsy April 2006. *Dev Med Child Neurol Suppl* 2007;109:8-14.
2. Odding E, Roebroek ME, Stam HJ. The epidemiology of cerebral palsy: incidence, impairments and risk factors. *Disabil Rehabil* 2006; 28:183-91.
3. Paneth N, Hong T, Korzeniewski S. The descriptive epidemiology of cerebral palsy. *Clin Perinatol* 2006;33:251-67.
4. Maenner MJ, Blumberg SJ, Kogan MD, Christensen D, Yeargin-Allsopp M, Schieve LA. Prevalence of cerebral palsy and intellectual disability among children identified in two U.S. National Surveys, 2011-2013. *Ann Epidemiol* 2016;26:222-6.
5. Access Economics. The economic impact of cerebral palsy in Australia in 2007. Canberra: Access Economics; 2008.
6. Benner JL, Noten S, Limsakul C, et al. Outcomes in adults with cerebral palsy: systematic review using the International Classification of Functioning, Disability and Health. *Dev Med Child Neurol* 2019;61:1153-61.
7. Fortuna RJ, Holub A, Turk MA, Meccarello J, Davidson PW. Health conditions, functional status and health care utilization in adults with cerebral palsy. *Fam Pract* 2018;35:661-70.
8. Whitney DG, Hurvitz EA, Ryan JM, et al. Noncommunicable disease and multimorbidity in young adults with cerebral palsy. *Clin Epidemiol* 2018;10:511-9.
9. Turk MA, Logan LR, Ansoanur F. Adults with childhood onset disabilities: a focused review of three conditions. *Curr Phys Med Rehabil Rep* 2013;1:72-87.
10. Benner JL, Hilberink SR, Veenis T, Stam HJ, van der Slot WM, Roebroek ME. Long-term deterioration of perceived health and functioning in adults with cerebral palsy. *Arch Phys Med Rehabil* 2017;98:2196-205.
11. Opheim A, Jahnsen R, Olsson E, Stanghelle JK. Walking function, pain, and fatigue in adults with cerebral palsy: a 7-year follow-up study. *Dev Med Child Neurol* 2009;51:381-8.
12. Benner JL, Hilberink SR, Veenis T, van der Slot WMA, Roebroek ME. Course of employment in adults with cerebral palsy over a 14-year period. *Dev Med Child Neurol* 2017;59:762-8.
13. Schmidt AK, Van Gorp M, Van Wely L, et al. Autonomy in participation in cerebral palsy from childhood to adulthood. *Dev Med Child Neurol* 2020;62:363-71.
14. van Gorp M, Van Wely L, Dallmeijer AJ, et al. Long-term course of difficulty in participation of individuals with cerebral palsy aged 16 to 34 years: a prospective cohort study. *Dev Med Child Neurol* 2019; 61:194-203.
15. Novak I, Hines M, Goldsmith S, Barclay R. Clinical prognostic messages from a systematic review on cerebral palsy. *Pediatrics* 2012;130:e1285-312.
16. Jonsson U, Eek MN, Sunnerhagen KS, Himmelmann K. Cerebral palsy prevalence, subtypes, and associated impairments: a population-based comparison study of adults and children. *Dev Med Child Neurol* 2019;61:1162-7.
17. WHO International. Classification of Functioning, Disability and Health. Geneva: ICF; 2001.
18. Ballester-Plane J, Laporta-Hoyos O, Macaya A, et al. Cognitive functioning in dyskinetic cerebral palsy: its relation to motor function, communication and epilepsy. *Eur J Paediatr Neurol* 2018;22:102-12.
19. Laporta-Hoyos O, Fiori S, Pannek K, et al. Brain lesion scores obtained using a simple semi-quantitative scale from MR imaging are associated with motor function, communication and cognition in dyskinetic cerebral palsy. *Neuroimage Clin* 2018;19:892-900.
20. Whitney DG, Hurvitz EA, Devlin MJ, et al. Age trajectories of musculoskeletal morbidities in adults with cerebral palsy. *Bone* 2018; 114:285-91.
21. Cremer N, Hurvitz EA, Peterson MD. Multimorbidity in middle-aged adults with cerebral palsy. *Am J Med* 2017;130:744.
22. Agustsson A, Sveinsson T, Pope P, Rodby-Bousquet E. Preferred posture in lying and its association with scoliosis and windswept hips in adults with cerebral palsy. *Disabil Rehabil* 2019;41:3198-202.
23. Alriksson-Schmidt A, Hagglund G, Rodby-Bousquet E, Westbom L. Follow-up of individuals with cerebral palsy through the transition

- years and description of adult life: the Swedish experience. *J Pediatr Rehabil Med* 2014;7:53-61.
24. Rodby-Bousquet E, Czuba T, Hagglund G, Westbom L. Postural asymmetries in young adults with cerebral palsy. *Dev Med Child Neurol* 2013;55:1009-15.
 25. van Gorp M, Roebroek ME, Swan Tan S, et al. Activity performance curves of individuals with cerebral palsy. *Pediatrics* 2018;142.
 26. van Meeteren J, Nieuwenhuijsen C, de Grund A, Stam HJ, Roebroek ME; Transition Research Group South West Netherlands. Using the manual ability classification system in young adults with cerebral palsy and normal intelligence. *Disabil Rehabil* 2010;32:1885-93.
 27. Nieuwenhuijsen C, Donkervoort M, Nieuwstraten W, Stam HJ, Roebroek ME; Transition Research Group South West Netherlands. Experienced problems of young adults with cerebral palsy: targets for rehabilitation care. *Arch Phys Med Rehabil* 2009;90:1891-7.
 28. van der Dussen L, Nieuwstraten W, Roebroek M, Stam HJ. Functional level of young adults with cerebral palsy. *Clin Rehabil* 2001;15:84-91.
 29. Kirk H, Geertsen SS, Lorentzen J, Krarup KB, Bandholm T, Nielsen JB. Explosive resistance training increases rate of force development in ankle dorsiflexors and gait function in adults with cerebral palsy. *J Strength Cond Res* 2016;30:2749-60.
 30. McPhee PG, Gorter JW, Cotie LM, Timmons BW, Bentley T, MacDonald MJ. Descriptive data on cardiovascular and metabolic risk factors in ambulatory and non-ambulatory adults with cerebral palsy. *Data Brief* 2015;5:967-70.
 31. McPhee PG, Gorter JW, Cotie LM, Timmons BW, Bentley T, MacDonald MJ. Associations of non-invasive measures of arterial structure and function, and traditional indicators of cardiovascular risk in adults with cerebral palsy. *Atherosclerosis* 2015;243:462-5.
 32. Slaman J, van den Berg-Emons HJ, van Meeteren J, et al. A lifestyle intervention improves fatigue, mental health and social support among adolescents and young adults with cerebral palsy: focus on mediating effects. *Clin Rehabil* 2015;29:717-27.
 33. Russchen HA, Slaman J, Stam HJ, et al. Focus on fatigue amongst young adults with spastic cerebral palsy. *J Neuroeng Rehabil* 2014;11:161.
 34. Slaman J, Roebroek M, van der Slot W, et al. Can a lifestyle intervention improve physical fitness in adolescents and young adults with spastic cerebral palsy? A randomized controlled trial. *Arch Phys Med Rehabil* 2014;95:1646-55.
 35. Ryan JM, Crowley VE, Hensey O, McGahey A, Gormley J. Waist circumference provides an indication of numerous cardiometabolic risk factors in adults with cerebral palsy. *Arch Phys Med Rehabil* 2014;95:1540-6.
 36. Morgan PE, Soh SE, McGinley JL. Health-related quality of life of ambulant adults with cerebral palsy and its association with falls and mobility decline: a preliminary cross sectional study. *Health Qual Life Outcomes* 2014;12:132.
 37. Morgan P, McGinley J. Performance of adults with cerebral palsy related to falls, balance and function: a preliminary report. *Dev Neurorehabil* 2013;16:113-20.
 38. Reddihough DS, Jiang B, Lanigan A, Reid SM, Walstab JE, Davis E. Social outcomes of young adults with cerebral palsy. *J Intellect Dev Disabil* 2013;38:215-22.
 39. van der Slot WM, Roebroek ME, Nieuwenhuijsen C, et al. Cardiovascular disease risk in adults with spastic bilateral cerebral palsy. *J Rehabil Med* 2013;45:866-72.
 40. van der Slot WM, Nieuwenhuijsen C, van Den Berg-Emons RJ, et al. Chronic pain, fatigue, and depressive symptoms in adults with spastic bilateral cerebral palsy. *Dev Med Child Neurol* 2012;54:836-42.
 41. van der Slot WM, Nieuwenhuijsen C, van den Berg-Emons RJ, et al. Participation and health-related quality of life in adults with spastic bilateral cerebral palsy and the role of self-efficacy. *J Rehabil Med* 2010;42:528-35.
 42. Opheim A, Jahnsen R, Olsson E, Stanghelle JK. Physical and mental components of health-related quality of life and musculoskeletal pain sites over seven years in adults with spastic cerebral palsy. *J Rehabil Med* 2011;43:382-7.
 43. Jahnsen R, Villien L, Egeland T, Stanghelle JK, Holm I. Locomotion skills in adults with cerebral palsy. *Clin Rehabil* 2004;18:309-16.
 44. Jahnsen R, Villien L, Aamodt G, Stanghelle JK, Holm I. Musculoskeletal pain in adults with cerebral palsy compared with the general population. *J Rehabil Med* 2004;36:78-84.
 45. Jahnsen R, Villien L, Stanghelle JK, Holm I. Coping potential and disability—sense of coherence in adults with cerebral palsy. *Disabil Rehabil* 2002;24:511-8.
 46. Maltais DB, Dumas F, Boucher N, Richards CL. Factors related to physical activity in adults with cerebral palsy may differ for walkers and nonwalkers. *Am J Phys Med Rehabil* 2010;89:584-97.
 47. Michelsen SI, Uldall P, Hansen T, Madsen M. Social integration of adults with cerebral palsy. *Dev Med Child Neurol* 2006;48:643-9.
 48. Michelsen SI, Uldall P, Kejs AM, Madsen M. Education and employment prospects in cerebral palsy. *Dev Med Child Neurol* 2005;47:511-7.
 49. Engel JM, Jensen MP, Hoffman AJ, Kartin D. Pain in persons with cerebral palsy: extension and cross validation. *Arch Phys Med Rehabil* 2003;84:1125-8.
 50. Gillet JG, Lichtwark GA, Boyd RN, Barber LA. Functional capacity in adults with cerebral palsy: lower limb muscle strength matters. *Arch Phys Med Rehabil* 2018;99:900-6.
 51. Lundh S, Nasic S, Riad J. Fatigue, quality of life and walking ability in adults with cerebral palsy. *Gait Posture* 2018;61:1-6.
 52. Park MW, Kim WS, Bang MS, et al. Needs for medical and rehabilitation services in adults with cerebral palsy in Korea. *Ann Rehabil Med* 2018;42:465-72.
 53. Peterson MD, Kamdar N, Hurvitz EA. Age-related trends in cardiometabolic disease among adults with cerebral palsy. *Dev Med Child Neurol* 2019;61:484-9.
 54. Sienko SE. An exploratory study investigating the multidimensional factors impacting the health and well-being of young adults with cerebral palsy. *Disabil Rehabil* 2018;40:660-6.
 55. Smith KJ, Peterson MD, O'Connell NE, et al. Risk of depression and anxiety in adults with cerebral palsy. *JAMA Neurol* 2019;76:294-300.
 56. Daunter AK, Kratz AL, Hurvitz EA. Long-term impact of childhood selective dorsal rhizotomy on pain, fatigue, and function: a case-control study. *Dev Med Child Neurol* 2017;59:1089-95.
 57. de Albuquerque Botura C, Ames FQ, de Albuquerque Botura AC, Bersani-Amado LE, Bardini AVSL, Cuman RKN. Pain symptoms in patients with severe cerebral palsy: prevalence among patients with higher degree of locomotor impairment. *Trop J Pharm Res* 2017;16:1431-6.
 58. Hayward K, Chen AY, Forbes E, Byrne R, Greenberg MB, Fowler EG. Reproductive healthcare experiences of women with cerebral palsy. *Disabil Health J* 2017;10:413-8.
 59. Park EY, Kim WH. Prevalence of secondary impairments of adults with cerebral palsy according to gross motor function classification system. *J Phys Ther Sci* 2017;29:266-9.
 60. Vukojević M, Cvitković T, Splavski B, Ostojić Z, Šumanović-Glamuzina D, Šimić J. Prevalence of intellectual disabilities and epilepsy in different forms of spastic cerebral palsy in adults. *Psychiatr Danub* 2017;29:111-7.
 61. Yıldız N, Akkoç Y, Ersöz M, et al. Cross-sectional study of urinary problems in adults with cerebral palsy: awareness and impact on the quality of life. *Neurol Sci* 2017;38:1193-203.
 62. Brunton L, Hall S, Passingham A, Wulff J, Delitala R. The prevalence, location, severity, and daily impact of pain reported by youth and young adults with cerebral palsy. *J Pediatr Rehabil Med* 2016;9:177-83.
 63. Peterson MD, Ryan JM, Hurvitz EA, Mahmoudi E. Chronic conditions in adults with cerebral palsy. *JAMA* 2015;314:2303-5.

64. Peterson MD, Haapala HJ, Chaddha A, Hurvitz EA. Abdominal obesity is an independent predictor of serum 25-hydroxyvitamin D deficiency in adults with cerebral palsy. *Nutr Metab (Lond)* 2014;11:22.
65. Huang IC, Wang YT, Chan F. Employment outcomes of adults with cerebral palsy in Taiwan. *Disabil Rehabil* 2013;35:228-35.
66. Hamrah Nedjad J, Jansson E, Bartonek Å. Physical activity in a Swedish youth and adult population with cerebral palsy. *Eur J Physiother* 2013;15:70-7.
67. Mesterman R, Leitner Y, Yifat R, et al. Cerebral palsy—long-term medical, functional, educational, and psychosocial outcomes. *J Child Neurol* 2010;25:36-42.
68. Gaskin CJ, Morris T. Physical activity, health-related quality of life, and psychosocial functioning of adults with cerebral palsy. *J Phys Act Health* 2008;5:146-57.
69. Andersson C, Asztalos L, Mattsson E. Six-minute walk test in adults with cerebral palsy. A study of reliability. *Clin Rehabil* 2006;20:488-95.
70. Sandström K, Alinder J, Öberg B. Descriptions of functioning and health and relations to a gross motor classification in adults with cerebral palsy. *Disabil Rehabil* 2004;26:1023-31.
71. Strauss D, Ojdana K, Shaville R, Rosenbloom L. Decline in function and life expectancy of older persons with cerebral palsy. *Neuro-Rehabilitation* 2004;19:69-78.
72. Andersson C, Mattsson E. Adults with cerebral palsy: a survey describing problems, needs, and resources, with special emphasis on locomotion. *Dev Med Child Neurol* 2001;43:76-82.
73. Bottos M, Feliciangeli A, Sciuto L, Gericke C, Vianello A. Functional status of adults with cerebral palsy and implications for treatment of children. *Dev Med Child Neurol* 2001;43:516-28.
74. Furukawa A, Iwatsuki H, Nishiyama M, Nii E, Uchida A. A study on the subjective well-being of adult patients with cerebral palsy. *J Phys Ther Sci* 2001;13:31-5.
75. Hodgkinson I, Jindrich ML, Duhaut P, Vadot JP, Metton G, Bérard C. Hip pain in 234 non-ambulatory adolescents and young adults with cerebral palsy: a cross-sectional multicentre study. *Dev Med Child Neurol* 2001;43:806-8.
76. Maruishi M, Mano Y, Sasaki T, Shinmyo N, Sato H, Ogawa T. Cerebral palsy in adults: independent effects of muscle strength and muscle tone. *Arch Phys Med Rehabil* 2001;82:637-41.
77. Murphy KPM, Molnar GE, Lankasky K. Employment and social issues in adults with cerebral palsy. *Arch Phys Med Rehabil* 2000;81: 807-11.
78. Aaronson NK, Muller M, Cohen PD, et al. Translation, validation, and norming of the Dutch language version of the SF-36 Health Survey in community and chronic disease populations. *J Clin Epidemiol* 1998;51:1055-68.
79. Krupp LB, LaRocca NG, Muir-Nash J, Steinberg AD. The fatigue severity scale. Application to patients with multiple sclerosis and systemic lupus erythematosus. *Arch Neurol* 1989;46:1121-3.
80. Eliasson AC, Krumlinde-Sundholm L, Rosblad B, et al. The Manual Ability Classification System (MACS) for children with cerebral palsy: scale development and evidence of validity and reliability. *Dev Med Child Neurol* 2006;48:549-54.
81. UNESCO Institute for Statistics. International Standard Classification of Education (ISCED) 2011. Montreal: UNESCO Institute for Statistics; 2012.
82. Lemmens J, van Engelen EISM, Post MWM, Beurskens AJHM, Wolters PMJC, de Witte LP. Reproducibility and validity of the Dutch Life Habits Questionnaire (LIFE-H 3.0) in older adults. *Clin Rehabil* 2007;21:853-62.
83. Collin C, Wade DT, Davies S, Horne V. The Barthel ADL Index: a reliability study. *Int Disabil Stud* 1988;10:61-3.
84. Linacre JM, Heinemann AW, Wright BD, Granger CV, Hamilton BB. The structure and stability of the Functional Independence Measure. *Arch Phys Med Rehabil* 1994;75:127-32.
85. Russell DJ, Rosenbaum PL, Cadman DT, Gowland C, Hardy S, Jarvis S. The gross motor function measure: a means to evaluate the effects of physical therapy. *Dev Med Child Neurol* 1989;31:341-52.
86. Graham HK, Harvey A, Rodda J, Nattrass GR, Pirpiris M. The Functional Mobility Scale (FMS). *J Pediatr Orthop* 2004;24:514-20.
87. Izumida S, Inoue S. Assessment of treatment for low back pain. *J Jpn Orthop Assoc* 1986;60:391-4.
88. Palisano RJ, Hanna SE, Rosenbaum PL, Russell DJ, Walter SD, Wood EP, et al. Validation of a model of gross motor function for children with cerebral palsy. *Phys Ther* 2000;80:974-85.
89. DerSimonian R, Laird N. Meta-analysis in clinical trials. *Control Clin Trials* 1986;7:177-88.
90. Freeman MF, Tukey JW. Transformations related to the angular and the square root. *The Ann Math Stat* 1950;21:607-11.
91. Miller JJ. The inverse of the Freeman-Tukey double arcsine transformation. *Am Stat* 1978;32:138.
92. Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ* 2003;327:557-60.
93. Himmelmann K, Beckung E, Hagberg G, Uvebrant P. Gross and fine motor function and accompanying impairments in cerebral palsy. *Dev Med Child Neurol* 2006;48:417-23.
94. To T, Stanojevic S, Moores G, et al. Global asthma prevalence in adults: findings from the cross-sectional world health survey. *BMC Public Health* 2012;12:204.
95. van der Slot W, Benner J, Brunton L, et al. Pain in adults with cerebral palsy: a systematic review and meta-analysis of individual participant data. *Ann Phys Med Rehabil* 2020 Feb 13 [Epub ahead of print].
96. Nahin RL. Estimates of pain prevalence and severity in adults: United States, 2012. *J Pain* 2015;16:769-80.
97. Valko PO, Bassetti CL, Bloch KE, Held U, Baumann CR. Validation of the fatigue severity scale in a Swiss cohort. *Sleep* 2008;31:1601-7.
98. Gross PH, Bailes AF, Horn SD, et al. Setting a patient-centered research agenda for cerebral palsy: a participatory action research initiative. *Dev Med Child Neurol* 2018;60:1278-84.
99. Vogtle LK. Pain in adults with cerebral palsy: impact and solutions. *Dev Med Child Neurol* 2009;51(Suppl 4):113-21.
100. Brunton LK, McPhee PG, Gorter JW. Self-reported factors contributing to fatigue and its management in adolescents and adults with cerebral palsy. *Disabil Rehabil* 2019;30(Jul):1-7.
101. Carnahan KD, Arner M, Hagglund G. Association between gross motor function (GMFCS) and manual ability (MACS) in children with cerebral palsy. A population-based study of 359 children. *BMC Musculoskelet Disord* 2007;8:50.
102. Langerak NG, Hillier SL, Verkoeijen PP, Peter JC, Fieggan AG, Vaughan CL. Level of activity and participation in adults with spastic diplegia 17-26 years after selective dorsal rhizotomy. *J Rehabil Med* 2011;43:330-7.
103. Donkervoort M, Roebroek M, Wiegerink D, van der Heijden-Maessen H, Stam H. Determinants of functioning of adolescents and young adults with cerebral palsy. *Disabil Rehabil* 2007;29: 453-63.
104. Jacobsen EL, Bye A, Aass N, et al. Norwegian reference values for the Short-Form Health Survey 36: development over time. *Qual Life Res* 2018;27:1201-12.
105. Surveillance of Cerebral Palsy in Europe. Surveillance of cerebral palsy in Europe: a collaboration of cerebral palsy surveys and registers. Surveillance of Cerebral Palsy in Europe (SCPE). *Dev Med Child Neurol* 2000;42:816-24.
106. Hurley DS, Sukal-Moulton T, Msall ME, Gaebler-Spira D, Krosschell KJ, Dewald JP. The cerebral palsy research registry: development and progress toward national collaboration in the United States. *J Child Neurol* 2011;26:1534-41.
107. McPhee PG, Benner JL, Balemans ACJ, et al. Multimorbidity risk assessment in adolescents and adults with cerebral palsy: a protocol for establishing a core outcome set for clinical research and practice. *Trials* 2019;20:176.
108. Maulik PK, Mascarenhas MN, Mathers CD, Dua T, Saxena S. Prevalence of intellectual disability: a meta-analysis of population-based studies. *Res Dev Disabil* 2011;32:419-36.