

# SARS-CoV-2 virus outbreak and the emergency public health measures in Bosnia and Herzegovina: January – July 2020

Mirsada Hukic<sup>1,2,#</sup>, Mirza Ponjavic<sup>3,4,#</sup>, Emin Tahirovic<sup>5,6,7</sup>, Almir Karabegovic<sup>3,8</sup>,  
Elvir Ferhatbegovic<sup>3,7</sup>, Maja Travar<sup>7,9,10</sup>, Fadila Serdarevic<sup>6,7,11,12\*</sup>

## ABSTRACT

Between March 5<sup>th</sup> and July 25<sup>th</sup>, 2020, the total number of SARS-CoV-2 confirmed cases in Bosnia and Herzegovina (BH) was 10,090, corresponding to a cumulative incidence rate of 285.7/100,000 population. Demographic and clinical information on all the cases along with exposure and contact information were collected using a standardized case report form. In suspected SARS-CoV-2 cases, respiratory specimens were collected and tested by real-time reverse-transcriptase polymerase chain reaction assay. The dynamic of the outbreak was summarized using epidemiological curves, instantaneous reproduction number  $R_t$  and interactive choropleth maps for geographical distribution and spread. The rate of hospitalization was 14.0% (790/5646) in the Federation of Bosnia and Herzegovina (FBH) and 6.2% (267/4299) in the Republic of Srpska (RS). The death rate was 2.2% (122/5646) in FBH and 3.6% in the RS (155/4299). After the authorities lifted mandatory quarantine restrictions, the instantaneous reproduction number increased from 1.13 on May 20<sup>th</sup> to 1.72 on May 31<sup>st</sup>. The outbreak concerns both entities, FBH and RS, and it is more pronounced in those aged 20-44 years. It is important to develop the communication and emergency plan for the SARS-CoV-2 outbreak in BH, including the mechanisms to allow the ongoing notification and updates at the national level.

KEYWORDS: SARS-CoV-2 outbreak; COVID-19 pandemic; emergency; public health

<sup>1</sup>Department of Medical Sciences, Academy of Sciences and Arts of Bosnia and Herzegovina, Sarajevo, Bosnia and Herzegovina

<sup>2</sup>Institute for Biomedical Diagnostic and Research NALAZ, Sarajevo, Bosnia and Herzegovina,

<sup>3</sup>GAUSS Centre for Geospatial Research Sarajevo, Sarajevo, Bosnia and Herzegovina

<sup>4</sup>Faculty of Engineering and Natural Sciences, International Burch University, Sarajevo, Bosnia and Herzegovina

<sup>5</sup>Faculty of Engineering and Natural Sciences, International University of Sarajevo, Sarajevo, Bosnia and Herzegovina

<sup>6</sup>South East European Network for Medical Research (SOVE), Sarajevo, Bosnia and Herzegovina

<sup>7</sup>Epidemic Location Intelligence System (ELIS) Project of Academy of Sciences and Arts of Bosnia and Herzegovina

<sup>8</sup>Faculty of Electrical Engineering, University of Sarajevo, Sarajevo, Bosnia and Herzegovina

<sup>9</sup>Faculty of Medicine, University of Banja Luka, Banja Luka, Bosnia and Herzegovina

<sup>10</sup>University Clinical Centre of the Republic of Srpska, Banja Luka, Bosnia and Herzegovina

<sup>11</sup>Erasmus Medical Center, Erasmus University, Rotterdam, Netherlands

<sup>12</sup>Faculty of Medicine, Sarajevo School of Science and Technology, Sarajevo, Bosnia and Herzegovina

\*Corresponding author: Fadila Serdarevic, Epidemic Location Intelligence System (ELIS) Project, Department of Medical Sciences, Academy of Sciences and Arts of Bosnia and Herzegovina, Bistrik 7, Sarajevo 71000, Bosnia and Herzegovina. Phone: +38733225485. E-mail: fserdarevic@gmail.com

#These authors equally contributed

DOI: <https://dx.doi.org/10.17305/bjbms.2020.5081>

Submitted: 23 August 2020/Accepted: 16 October 2020

Conflict of interests: The authors declare no conflict of interests.



©The Author(s) (2021). This work is licensed under a Creative Commons Attribution 4.0 International License

## INTRODUCTION

On March 5<sup>th</sup>, 2020, the two first probable cases of SARS-CoV-2 virus reported to the Institute of Public Health (IPH) of Republika Srpska (RS) in Bosnia and Herzegovina (BH) were confirmed positive by real-time reverse-transcription-polymerase chain reaction (RT-PCR) at the University Clinical Center Laboratory of the RS in Banja Luka. The cases were a 35-year-old male returning from Italy and his 14-year-old son. Both experienced mild symptoms of cough, fever, and sore throat. On March 21<sup>st</sup>, the first death occurred in BH. The patient was a 76-year-old woman without a travel history subsequently hospitalized for COVID-19 2 days earlier in Bihac, Federation of Bosnia and Herzegovina (FBH). Enhanced surveillance for COVID-19 cases began on April 25 in the RS and on April 19 in the FBH and revealed many additional cases [1]. This report describes the initial and ongoing SARS-CoV-2 virus outbreak in BH for the period from January to July 2020.

## MATERIALS AND METHODS

### Surveillance

In response to increased reports of SARS-CoV-2 virus infections in BH, from March 5<sup>th</sup> IPH in RS and FBH issued an

entity-wide health alert to health-care providers recommending a collection of a respiratory swab for SARS-CoV-2 virus testing using RT-PCR for persons with influenza-like illness (ILI) and travel history to affected areas. From April 10<sup>th</sup> in RS and from April 15<sup>th</sup> in FBH, testing for the SARS-CoV-2 virus was extended to all suspected and probable ILI cases using the WHO criteria, regardless of travel history [2]. Demographic and clinical information on all cases along with exposure and contact information was collected using a standardized case report form. The dynamics of the outbreak were summarized using epidemiological curves, instantaneous reproduction number  $R_t$  and interactive choropleth maps for geographical distribution and spread [3]. We calculated  $R_t$  using delayed reporting dates of positive confirmed cases. To estimate the dates of infection, we randomly generated the incubation and symptomatic period (1-6 days). The incubation period was governed by the Weibull distribution [4]. To estimate daily  $R_t$  and its credible interval pertaining to the previous week, we used EpiEstim R package [3] with the mean and the standard deviation of the serial interval [5].

## RESULTS

Between March 5<sup>th</sup> and July 25<sup>th</sup>, 2020, a total of 4299, 5646, and 145 laboratory-confirmed SARS-CoV-2 cases were reported to IPH of the RS, IPH of the FBH, and Department of Public Health of the District Brčko (DB) starting with 5<sup>th</sup>, 21<sup>st</sup>, and 27<sup>th</sup> of March respectively, with estimated infection dates between February 24<sup>th</sup> and July 13<sup>th</sup>, 2020 (Figure 1).

By July 25<sup>th</sup>, 94% of municipalities in FBiH (74/79), 92% of RS municipalities (57/62), and DB reported at least one case (Figure 2). Fifty-one percent of all cases were males.

As shown in Table 1, the majority of cases were 20-44 years old (38.5%). The rate of hospitalization was 14.0% (790/5646)

in FBH and 6.2% (267/4299) in RS. The death rate was 2.2% (122/5646) in FBH and 3.6% in the RS (155/4299).

## Prevention and control measures

Beginning March 15<sup>th</sup>, IPH in the RS and March 17<sup>th</sup> in the FBH issued a number of health alerts to the health care community about infection control, diagnostic testing, and reporting of cases. Early on in the COVID-19 outbreak, they implemented intense measures to facilitate physical distancing, including complete lock-down for children under age 18 years and elderly above age 65 years, curfew, border closings, home-isolation for residents coming from affected areas, obligatory face-masks wearing, and other restrictions in an attempt to slow the spread of the SARS-CoV-2 virus (Table 2) [6,7]. In less than a month, the daily number of new cases in BH leveled off. The instantaneous reproduction number dropped from 1.68 on March 15<sup>th</sup> to 1.00 on April 15<sup>th</sup> (Figure 1).

Until authorities lifted mandatory quarantine restrictions, by April 24<sup>th</sup> in the FBH and by May 12<sup>th</sup> in the RS, the number of infected persons was 786 and 985, respectively. Since May 20<sup>th</sup> (FBH) and May 21<sup>st</sup> (RS), mandatory self-isolation (14 days) at home upon entry into the country has been abolished. At the same time number of country-wide confirmed cases and instantaneous reproduction number started to increase again from 1.13 on May 20<sup>th</sup> to 1.72 on May 31<sup>st</sup> (Figure 1). Despite recommendations to avoid close contact and large congregations, large public gatherings occurred in the country (Table 2). As of July 25<sup>th</sup>, the total number of COVID-19 confirmed cases in BH was 10,090, corresponding to a cumulative incidence rate of 285.7/100,000 population (Figure 1).

Both entities IPH continued recommending mitigation measures, such as wearing face masks, physical distancing, widespread testing for suspected cases, as well as home

**TABLE 1.** Epidemiologic characteristics of the confirmed cases in Bosnia and Herzegovina

	Bosnia and Herzegovina (Total)		Distribution by the administrative unit					
	n	%	Federation of BH		Republika Srpska		District Brcko	
			n	%	n	%	n	%
Cases	10149	100.0	5646	55.6	4363	43.0	140	1.4
Gender								
Male	5142	50.7	2861	50.7	2209	50.6	72	51.4
Female	5007	49.3	2785	49.3	2154	49.4	68	48.6
Unknown	0	0.0	0	0.0	0	0.0	0	0.0
Age group (years)								
<5	204	2.0	96	1.7	107	2.5	1	0.7
5-19	564	5.6	424	7.5	127	2.9	13	9.3
20-44	3917	38.6	2339	41.4	1517	34.8	61	43.6
45-54	2013	19.8	977	17.3	1018	23.3	18	12.9
55-64	1776	17.5	955	16.9	801	18.4	20	14.3
65-74	1032	10.2	505	8.9	513	11.8	14	10.0
75-84	477	4.7	273	4.8	201	4.6	3	2.1
>84	139	1.4	56	1.0	79	1.8	4	2.9
Unknown	6	0.1	0	0.0	0	0.0	6	4.3




FIGURE 1. Epidemic curves and weekly Rt estimates based on inferred infection dates.




FIGURE 2. Cumulative number of laboratory-confirmed cases per 100,000 population.

**TABLE 2.** Restrictive, relaxing measures, and large public gatherings in Bosnia and Herzegovina

Date	Federation BiH	Republika Srpska	Type
11/3/2020		Classes moved online (primary ed.)	Restrictive
12/3/2020	Classes moved online (all academic levels)		Restrictive
16/3/2020		Emergency situation declared	Restrictive
17/3/2020	Large gatherings (>100) banned, Natural disaster proclaimed		Restrictive
19/3/2020	Regional and local public transportation discontinued		Restrictive
20/3/2020	Movement restriction for 18- i 65+		Restrictive
21/3/2020	Curfew 6 PM-5 AM, Physical distancing ordered by decree	Movement restriction for 65+, Curfew 8 PM-5 AM	Restrictive
22/3/2020	Physical distancing in public compulsory (1.5 m)	Classes moved online (high schools and universities), Regional public transportation discontinued	Restrictive
23/3/2020		State of emergency declared	Restrictive
29/3/2020	Curfew later start, 8 PM-5 AM		Relaxing
2/4/2020		Public transportation (<55 km) discontinued.	Restrictive
4/4/2020		Movement outside the place of residence banned	Restrictive
5/4/2020	Movement for 65+ between 8 AM and 12 PM allowed		Relaxing
6/4/2020		Movement outside the place of residence allowed	Relaxing
10/4/2020	Movement restriction for 65+ reinstated	Public gatherings limited to 3 persons, mask, and gloves compulsory	Restrictive
17/4/2020		Movement restriction during Easter weekend	Restrictive
20/4/2020		Movement restriction ceases	Relaxing
25/4/2020	Curfew and compulsory quarantine when entering the Federation discontinued		Relaxing
30/4/2020	Movement restriction during May Day	Movement restriction during May Day, Monetary fines 100-300 BAM	Restrictive
4/5/2020	Movement restriction ceases	Movement restriction ceases	Relaxing
11/5/2020	A wide range of businesses reopen	A wide range of businesses reopen	Relaxing
		High school graduation celebrations banned	Restrictive
14/5/2020		Movement restriction for 65+ discontinued	Relaxing
15/5/2020	Movement restriction for 65+ discontinued		Relaxing
15/5/2020	Antifascist movement protests (Sarajevo)		Large public gathering
25/5/2020		Pools, wellness, and spa centers allowed to reopen	Relaxing
29/5/2020	Platform for progress (political party organized) protests		Large public gathering
29/5/2020	State of natural disaster ceases		Relaxing
31/5/2020	All restrictive measures by Federal covid-19 crisis committee discontinued, A wide range of restaurants and bars reopen Public gatherings allowed up to 100 indoor, up to 300 outdoor		Relaxing
3/6/2020		Rules of operation for malls and stores adjusted	Restrictive
8/6/2020		Shortened working hours for bars, restaurants 6-23 hours; Sporting competitions and events banned	Restrictive
12/6/2020	Federal parliament transfers decisions about public health measures to cantonal public health bodies		
25/6/2020	Medjugorje (catholic pilgrimage site) 39 <sup>th</sup> anniversary of the apparition of Virgin Mary		Large public gathering
28/6/2020	Ajvatovica – (Muslim pilgrimage site) – an annual pilgrimage		Large public gathering
2/7/2020	Enhanced supervision over pH measures employment		Restrictive
6/7/2020		Sporting competitions and events allowed to continue without an audience present	Relaxing
7/7/2020		Public gatherings of up to 50 people allowed	
17/7/2020	SARS-COV-2 -Cov-2 epidemic officially proclaimed; enhanced sanitary surveillance at the border crossings and intensified control of the protective epidemiological measures on the ground		Restrictive
20/7/2020		Wearing face masks outside or maintaining 2 m physical distance compulsory	Restrictive

isolation for residents with acute febrile respiratory illness. They also continued disseminating educational messages regarding respiratory and hand hygiene through the media and their websites (Table 2).

## DISCUSSION

Rapidly establishing surveillance and public health response to the SARS-CoV-2 pandemic was particularly

challenging in BH due to the complex organization of the country's public health and health-care systems.

As a consequence of brutal conflict (1992-1995) that ended up with the Dayton peace agreement, BH has thirteen Ministries of Health (MoH), each with an IPH, for a population of less than 4 million – one for each entity, one for the District of Brčko, and one for each of the ten cantonal ministries in FBiH [8]. IPHs are responsible for communicable disease health monitoring, prevention, and control activities within their administrative jurisdiction. In RS, such regulation and management are under the jurisdiction of the MoH and Social Welfare. The central MoH of FBH coordinates cantonal health administrations at the entity level. The entity-level IPH compiles data and develops monthly/yearly reports for all notifiable diseases. There is limited communication of health data between entities. Communicable diseases data on BH level are compiled by the WHO based on the official entity level IPH monthly reports. This politically driven compartmentalization motivated the Academy of Sciences and Arts of Bosnia and Herzegovina to establish Epidemic Location Intelligence System under the patronage of the BH Presidency to track the SARS-CoV-2 pandemic at the national level.

Lower numbers of confirmed cases of SARS-CoV-2 infections compared to other areas of Europe were reported at the beginning of the outbreak [2]. Possible explanations for the relatively low confirmed case numbers in BH from March 16<sup>th</sup> until May 20<sup>th</sup> with average weekly  $R_t$  1.029, and cumulatively 2546 infected individuals can be potentially attributed to early measures, lock-down of the country, border closings, travel measures, home isolation, contact-tracing, and lack of testing. A potential explanation for the subsequent and the current increase in numbers of confirmed cases of SARS-CoV-2 infections from May 21<sup>st</sup> until July 25<sup>th</sup> with average weekly  $R_t$  for this period 1.19 and cumulatively 7569 infected individuals may be due to discontinuation of obligatory social measures, mass gatherings, and public events, enhanced hospital laboratory capacity and emerged of local transmission. Furthermore, the post-conflict complex governance structure in BH impedes communication and coordination, which in turn poses additional risk for disease spread. For example, when the authorities in the FBH abolished mandatory quarantine at the beginning of April without consultations with the authorities in the RS, it provoked strong reactions from RS authorities who named this “an insane and completely irresponsible move, professionally and epidemiologically unfounded” [9].

Among confirmed cases of SARS-CoV-2 infections in BH, the most affected age group was 20-44 years. These are working-age young people, usually working outside the home and socially most active hence most likely to have attended large gatherings. They can potentially transmit the virus to the elderly and other vulnerable groups later on [10]. The number

of reported deaths represents a small percentage, 2.8%, of total confirmed cases, either due to the current high transmission among young people or/and due to lag in transmission and reporting or simply due to expended testing, as more milder cases become identified [11].

## Public health recommendations

Responses for the current wave and preparations for future waves of SARS-CoV-2 influenza infections should be guided by populations that are most likely to be socially active and those at greatest risk for serious complications of SARS-CoV-2 infections. The lockdown of the country at the beginning of the outbreak was reflected in decreased instantaneous reproduction number. However, such a stringent measure is not easily sustainable, as it carries significant psychosocial, economic, and political consequences. Therefore, it is important to develop a communication and emergency plan for SARS-CoV-2 outbreak, including mechanisms to allow ongoing notification and updates at the national level, enable ongoing monitoring of transmission dynamic to facilitate timely adjustments of public health measures, and to communicate risk and event information to all communities including countering misinformation. Emphasis should continue on prompt testing of suspected cases and tracing of their contacts, and on universal prevention measures, including physical distancing, use of face masks, increased hand- and environmental hygiene, and avoidance of mass gatherings and crowded settings [7]. In addition, other strategies proposed recently for developing countries, such as zonal and dynamic blocs, can be considered [12,13]. Finally, coordination and collaboration with neighboring countries and the wider global community will be the key to reducing virus resurgence risks while minimizing travel and trade disruptions.

## ACKNOWLEDGMENTS

We would like to acknowledge Dr. Amra Uzicanin (CDC Atlanta, USA), Dr. Ambuj Kumar (USF, USA), and Tea Reljic (USF, USA) for their helpful comments to an earlier version of this manuscript. We would also like to acknowledge the COVID-19 B&H Epidemiologic Defense Group for providing data for this analysis: dr.Aida Pilav: Institute for Public Health of Sarajevo Canton, dr.Sead Karakas: Institute for Public Health of Central Bosnia Canton, dr.Ivan Vasilj: Institute for Public Health of West Herzegovina Canton, dr. Maida Mulic: Institute for Public Health of Tuzla Canton, dr. Nermin Avdic: Institute for Public Health of Herzegovina-Neretva Canton, dr.Zarina Mulabdic: Institute for Public Health of Una-Sana Canton, dr.Goran Pavic: Institute for Public Health of Posavina Canton, dr.Medina Bico: Institute

for Public Health of Bosnian-Podrinje Canton, dr. Diana Mamic, Institute for Public Health of Herzeg-Bosnian Canton, dr. Senad Huseinagic: Institute of Health and Food Safety of Zenica Canton, dr. Danica Lazic and dr. Anda Nikolic: Department of Health, Brcko District of Bosnia and Herzegovina

## REFERENCES

- [1] Arapović J, Skočibušić S. The first two months of the COVID-19 pandemic in Bosnia and Herzegovina: Single-center experience. *Bosn J Basic Med Sci.* 2020 Aug 3;20(3):396-400. <https://doi.org/10.17305/bjbms.2020.4838>.
- [2] World Health Organization. Global Surveillance for COVID-19 Caused by Human Infection with COVID-19 Virus: Interim Guidance. Geneva: World Health Organization; 2020.
- [3] Cori A, Ferguson NM, Fraser C, Cauchemez S. A new framework and software to estimate time-varying reproduction numbers during epidemics. *Am J Epidemiol* 2013;178(9):1505-12. <https://doi.org/10.1093/aje/kwt133>.
- [4] Backer JA, Klinkenberg D, Wallinga J. Incubation period of 2019 novel coronavirus (2019-nCoV) infections among travellers from Wuhan, China, 20-28 January 2020. *Euro Surveill* 2020;25(5):2000062. <https://doi.org/10.2807/1560-7917.es.2020.25.5.2000062>
- [5] Yang L, Dai J, Zhao J, Wang Y, Deng P, Wang J. Estimation of incubation period and serial interval of COVID-19: Analysis of 178 cases and 131 transmission chains in Hubei province, China. *Epidemiol Infect* 2020;148:e117. <https://doi.org/10.1017/S0950268820001338>
- [6] IJZRS. Korona virus COVID-19: Aktuelno: Institut za Javno Zdravstvo Republike Srpske; 2020. Available from: <https://www.phi.rs.ba/index.php?view=kategorija&id=45>. [Last accessed on 2020 Aug 10].
- [7] Paragraf Lex BA. Najnoviji PROPISI Bosne i Hercegovine Vežani za Sprečavanje Širenja Bolesti-covid19 (Korona Virus) Sarajevo, BH; 2020. Available from: <https://www.paragraf.ba/propisi-naredbe-bosne-i-hercegovine-za-sprečavanje-sirenja-korona-virusa-covid-19.html>. [Last accessed on 2020 Aug 10].
- [8] Organization for Security and Co-operation in Europe. The General Framework Agreement for Peace in Bosnia and Herzegovina. Dayton, OH: Organization for Security and Co-operation in Europe; 1995.
- [9] Vukić U. Nenad Stevandić: Ukidanje Karantina u FBiH Lud i Neodgovoran Potez. Banja Luka: Nezavisne Novine; 2020. Available from: <https://www.nezavisne.com/novosti/drustvo/Nenad-Stevandic-Ukidanje-karantina-u-FBiH-lud-i-neodgovoran-potez/596205>. Last accessed: 27 May 2020.
- [10] Golubev AG, Sidorenko AV. Theory and practice of aging upon COVID-19 pandemic. *Adv Gerontol* 2020;33(2):397-408.
- [11] Chang CS, Yeh YT, Chien TW, Lin JJ, Cheng BW, Kuo SC. The computation of case fatality rate for novel coronavirus (COVID-19) based on Bayes theorem: An observational study. *Medicine (Baltimore)* 2020;99(21):e19925. <https://doi.org/10.1097/md.00000000000019925>.
- [12] Chowdhury R, Luhar S, Khan N, Choudhury SR, Matin I, Franco OH. Long-term strategies to control COVID-19 in low and middle-income countries: An options overview of community-based, non-pharmacological interventions. *Eur J Epidemiol* 2020;35(8):743-8. <https://doi.org/10.1007/s10654-020-00660-1>.
- [13] Chowdhury R, Heng K, Shawon MS, Goh G, Okonofua D, Ochoa-Rosales C, et al. Dynamic interventions to control COVID-19 pandemic: A multivariate prediction modelling study comparing 16 worldwide countries. *Eur J Epidemiol* 2020;35(5):389-99. <https://doi.org/10.1007/s10654-020-00649-w>.

---

## Related articles published in BJBMS

1. [The first two months of the COVID-19 pandemic in Bosnia and Herzegovina: Single-center experience](#)  
Jurica Arapović et al., BJBMS, 2020
2. [SARS-CoV-2 infection of the nervous system: A review of the literature on neurological involvement in novel coronavirus disease-\(COVID-19\)](#)  
Alvin Oliver Payus et al., BJBMS, 2020
3. [Kawasaki-like disease and acute myocarditis in the SARS-CoV-2 pandemic - reports of three adolescents](#)  
Stasa Krasic et al. BJBMS, 2020
4. [Rapid, multimodal, critical care knowledge-sharing platform for COVID-19 pandemics](#)  
Amra Sakusic et al. BJBMS, 2020