

in T acute lymphoblastic leukemias improves understanding of early human T lymphoid lineage commitment. *Blood*. 2003;101:2693-2703.

10. Roumier C, Eclache V, Imbert M, et al. M0 AML, clinical and biologic features of

the disease, including AML1 genemutations: a report of 59 cases by the Groupe Francais d'Hematologie Cellulaire (GFHC) and the Groupe Francais de Cytogenetique Hematologique (GFCH). *Blood*. 2003;101:1277-1283.

## Response

### ***CEBPA* promoter hypermethylation in a subset of myeloid/T-lymphoid leukemias with a distinct gene expression profile**

With great interest we have read the letter by Terriou and coworkers, who have performed an extensive survey of *CEBPA* CpG promoter hypermethylation in T-ALL as well as in a selection of immature acute myeloid leukemia (AML) cases. This study relates to our previous work in which we identified hypermethylation of the proximal *CEBPA* promoter in a small subset of AMLs with low *CEBPA* mRNA.<sup>1</sup> It should be noted that the patient group we described was not primarily defined by *CEBPA* promoter hypermethylation. Instead, those cases were studied because they exhibited a unique gene expression profile. In addition to this most discriminating feature, the leukemic blasts coexpressed CD34 and T-cell antigens with myeloid markers, frequently carried *NOTCH1* mutations, and expressed the transforming gene *TRIB2*.<sup>1,2</sup> Using a predictive gene expression signature in an independent cohort of AML, we identified genetically and immunophenotypically similar cases with silenced *CEBPA*.

In the present study, Terriou et al found *CEBPA* promoter hypermethylation in 4/54 AML cases. These 4 cases all revealed one or more T-cell characteristics (cCD3, CD7, and/or *TCRG* rearrangement) and may therefore resemble the leukemias we described. In T-ALL, the investigators found *CEBPA* promoter hypermethylation more frequently, in 37/99 cases. Importantly, however, only a minority of those T-ALLs with methylated *CEBPA* coexpressed myeloid surface markers (CD13 and/or CD33) and CD34. The findings of Terriou et al therefore indicate that *CEBPA* proximal promoter hypermethylation in combination with an immature myeloid/T-lymphoid immunophenotype is generally rare

in acute leukemia, and can be found in a small percentage of AML as well as in a small fraction of T-ALL. It will be important to apply gene expression profiling to these cases to fully assess their relationship with the leukemias that we described previously.

Together, these observations highlight the challenges in classifying particular acute leukemias as AML, ALL, or a unique entity. We believe that studies as the one performed by Terriou et al will be instrumental in obtaining a better understanding of this topic. Gene expression profiling applied to these AML and T-ALL cohorts will clearly be of added value. Combining data from multiple research groups will be a requirement to address important questions regarding prognosis, and should help to evaluate according to which protocols these relatively small patient groups are best treated.

**Bas Wouters and Ruud Delwel**

*Conflict-of-interest disclosure: The authors declare no competing financial interests.*

*Correspondence: Bas Wouters or Ruud Delwel, Department of Hematology, Erasmus University Medical Center, PO Box 2040, 3000CA, Rotterdam, The Netherlands; e-mail: b.wouters@erasmusmc.nl or h.delwel@erasmusmc.nl.*

## References

1. Wouters BJ, Jorda MA, Keeshan K, et al. Distinct gene expression profiles of acute myeloid/T-lymphoid leukemia with silenced *CEBPA* and mutations in *NOTCH1*. *Blood*. 2007;110:3706-3714.
2. Keeshan K, He Y, Wouters BJ, et al. Tribbles homolog 2 inactivates C/EBPα and causes acute myelogenous leukemia. *Cancer Cell*. 2006;10:401-411.

## To the editor:

### **JAK2-V617F–triggered preemptive and salvage adoptive immunotherapy with donor-lymphocyte infusion in patients with myelofibrosis after allogeneic stem cell transplantation**

Primary myelofibrosis is a myeloproliferative disease, and results of conventional treatment remain unsatisfactory.<sup>1,2</sup> Allogeneic stem cell transplantation after dose-reduced conditioning has become a reasonable, curative treatment option.<sup>3,4</sup> Single case reports about successful donor lymphocyte infusion (DLI) for relapsed patients provided evidence of a graft-versus-myelofibrosis effect.<sup>5-7</sup> Here, we report on 17 patients with either myelofibrosis (n = 16) or secondary AML post myelofibrosis (n = 1) and a median age of 52 years (range, 32-63 years) who received DLI from related (n = 5) or unrelated (n = 12) donor, either for clinical relapse (salvage DLI; n = 9) or residual disease monitored by JAK2 mutation level in peripheral blood (preemptive DLI; n = 8). Details are summarized in Table 1. One patient (no. 9) received DLI twice: once for molecular residual disease and once for reappearance of molecular disease. Sixteen patients were JAK2-V617F–positive. The median time from transplantation to first DLI

was 269 days (range, 127-1570 days). The median percentage of JAK2V617 mutation level in peripheral blood before first DLI was 6.2% (range, 0.2%-72.8%) and significantly higher in patients with clinical relapse than with molecular relapse (24.7% vs 0.37%; *P* = .03). The median cell dose of the first DLI was 10<sup>6</sup> CD3<sup>+</sup> cells per kilogram of body weight (BW; range, 0.5-9 × 10<sup>6</sup> cells/kg BW). Two patients received as first DLI CD4-selected T cells (5 × 10<sup>6</sup> CD4<sup>+</sup> cells/kg BW). Second and subsequent half-log-increased DLI were given if no graft-versus-host disease (GVHD) and no significant response were observed. The median interval between the first and the second DLI was 103 days (range, 43-661 days). The response to DLI for relapsed patients was determined by the response criteria of the International Working Group,<sup>8</sup> and for residual disease by quantitative real time JAK2-V617F polymerase chain reaction (PCR) performed from genomic DNA from peripheral blood as recently described.<sup>9</sup> The sensitivity