

DERIVED STRENGTHS OF PREFERENCE RELATIONS ON COORDINATES *

Peter WAKKER

University of Nijmegen, 6500 HE Nijmegen, The Netherlands

Received 20 May 1988

Accepted 12 July 1988

A way is indicated to derive, from a preference relation on a Cartesian product, strength of preference relations on the coordinate sets. These strengths of preference relations are then used to reformulate several well-known properties of preference relations, and make their meaning more transparent. A new result for dynamic contexts is given.

1. Introduction

This paper examines the existence of several kinds of additively decomposable representing functions for preference relations on Cartesian products. This is done under the assumption that the coordinate sets are connected topological spaces, e.g., \mathbb{R}_+^n , or \mathbb{R} . The representing functions which we shall find will be cardinal.

Several authors have dealt with cardinal additively decomposable representing functions under the assumption that the coordinate sets are ‘mixture spaces’, the main intended examples of mixture spaces being sets of probability distributions. See, for instance, Anscombe and Aumann (1963), Fishburn (1965), Pollak (1967), Keeney and Raiffa (1976, Karni, Schmeidler and Vind (1983). Usually the axioms of von Neumann and Morgenstern (1953, ch. 3 and the appendix) are assumed, leading to linear representing functions. Cardinality is easily obtained in the presence of linearity because, if one linear function is a (strictly increasing) transformation of another linear function, then in fact it must be a linear transformation. Hence, if two linear functions induce the same preference relation, then they also induce the same strength of preference relation.

Matters are more complicated if the assumption of linearity is weakened to the assumption of continuity. To see this, let f and g be two continuous functions on a connected domain. Let $f = \varphi \cdot g$ for a transformation φ . For guaranteeing that φ is (positively) linear, the assumption of strict increasingness of φ no longer suffices. It must be strengthened to the more complicated assumption that φ preserves (‘first-order’) differences, as shown in Basu (1982). The characterizing properties for preference relations on connected topological spaces, met in literature, reflect this additional complication.

This paper handles the additional complication by indicating a way to derive, from a preference relation on a Cartesian product (‘revealed’), strength of preference relations on the coordinate sets. By means of these we formulate several properties and results, met in literature, for connected topological spaces. All reformulated versions will require that certain strengths of preferences,

* This paper was prepared while visiting the Department of Economics of the Tel-Aviv University.

derived from the preference relation, do not reveal contradictions. We think the reformulated versions are more appealing, in clarifying the first-order difference idea which is always underlying continuous cardinal utility.

Also we use the strength of preference relations to obtain a new result for dynamic contexts. Koopmans (1972) gave a characterization of additively decomposable representing functions with constant discount factors, mainly by means of a stationarity assumption. This requires an infinite set of (coordinates) points of time, \mathbb{N} in Koopmans' paper. Grodal (1978, Theorem 4) has given a result for continuous time. We obtain a result for individuals who deal with only a finite number of points of time, as considered for instance in Samuelson (1958). Zilcha (1988) is a recent survey, indicating the usefulness in intergenerational models of finiteness of the number of points of time for each individual.

2. Preferences and strengths of preferences

This section gives elementary definitions and notations. Further it indicates the way to derive the strength of preference relations from the preference relation.

Let $n \in \mathbb{N}$. Let $\mathcal{C}_1, \dots, \mathcal{C}_n$ be connected topological spaces. The set of alternatives is the Cartesian product $X_{i=1}^n \mathcal{C}_i$. It is endowed with the product topology, which is connected too.

Let \succsim be a binary relation on X , the preference relation of a decision maker. We write $x \preccurlyeq y$ if $y \succsim x$. As usual $\succ = \succsim \setminus \preccurlyeq$, and $\approx = \succsim \cap \preccurlyeq$. We write $x \prec y$ if $y \succ x$. Further, \succsim is continuous if $(x: x \succsim y)$ and $(x: x \preccurlyeq y)$ are closed sets for all $y \in X$. A weak order \succsim is complete ($x \succsim y$ or $y \succsim x$ for all $x, y \in X$) and transitive. A function $V: X \rightarrow \mathbb{R}$ represents \succsim if $[x \succsim y \Leftrightarrow V(x) \geq V(y)]$ for all $x, y \in X$. $V: X \rightarrow \mathbb{R}$ is additively decomposable if there exist $V_j: \mathcal{C}_j \rightarrow \mathbb{R}$, $j = 1, \dots, n$, such that $V: x \mapsto \sum_{j=1}^n V_j(x_j)$. Here, as usual, x_j is the j th coordinate of x .

We write $x_{-i}v_i$ for x (with x_i replaced by v_i), and for $i \neq j$, $(x_{-i,j}v_i, w_j)$ for x (with x_i replaced by v_i , and x_j by w_j).

Coordinate (or index) i is non-essential if $x_{-i}v_i \approx x$ for all x, v_i ; otherwise i is essential.

The following definitions give the main tools in this paper.

Definition 1. The quaternary relation \succ_i^{**} on \mathcal{C}_i , with $x_i y_i \succ_i^{**} v_i w_i$ meaning $(x_i, y_i, v_i, w_i) \in \succ_i^{**}$, is defined by

$$x_i y_i \succ_i^{**} v_i w_i \quad \text{if} \quad a_{-i} x_i \succsim b_{-i} y_i \quad \text{and}$$

$$a_{-i} v_i \prec b_{-i} w_i \quad \text{for some alternatives } a, b.$$

For an intuitive elucidation, suppose for simplicity that x_i is better than y_i , and v_i is better than w_i . Then $a_{-i} v_i \prec b_{-i} w_i$ suggests that the positive argument, to obtain v_i instead of w_i on coordinate i , does not outweigh the obviously negative argument for the left alternative against the right alternative, as yielded by the coordinates $j \neq i$. And $a_{-i} x_i \succsim b_{-i} y_i$ suggests that the positive argument, to obtain x_i instead of y_i , does outweigh the same (negative) argument. We interpret this to mean that it has been revealed from \succsim , that the strength of preference of x_i over y_i exceeds that of v_i over w_i .

A further definition:

Definition 2. The quaternary relation \succsim_i^* on \mathcal{C}_i , with $x_i y_i \succsim_i^* v_i w_i$ meaning $(x_i, y_i, v_i, w_i) \succsim_i^*$, is defined by

$$x_i y_i \succsim_i^* v_i w_i \text{ if } a_{-i} x_i \succ b_{-i} y_i \text{ and } a_{-i} v_i \preccurlyeq b_{-i} w_i \text{ for some alternatives } a, b.$$

The introduced binary relations in general do not have to satisfy properties such as completeness or transitivity. And \succ_i^{**} in no way has to be the asymmetric part of \succsim_i^* ; see Theorem 3.1.

3. Characterizations through strength of preferences

Throughout this section we assume that \succ is a continuous weak order on X . We shall list several properties of \succ , give their equivalent formulations in terms of \succsim_i^* and \succ_i^{**} , and give the main theorems in which the properties are used.

We say that \succ satisfies triple cancellation if

$$\left[\begin{array}{l} \text{For all } i, a, \dots, w_i, \\ a_{-i} x_i \preccurlyeq b_{-i} y_i \text{ and} \\ a_{-i} v_i \succ b_{-i} w_i \text{ and} \\ c_{-i} x_i \succ d_{-i} y_i \text{ imply} \\ c_{-i} v_i \succ d_{-i} w_i \end{array} \right] \text{ which is iff } \left[\begin{array}{l} \text{For all } i, x_i, \dots, w_i, \\ v_i w_i \succsim_i^* x_i y_i, \\ \text{then not} \\ x_i y_i \succ_i^{**} v_i w_i \end{array} \right]. \tag{1}$$

Theorem 1. Let at least two coordinates be essential. There exists a continuous additively decomposable representing function V for \succ , if and only if \succ satisfies triple cancellation. V is cardinal.

Proof. For the case of exactly two essential coordinates this can be derived from Chapter 6 of Krantz, Luce, Suppes and Tversky (1971), as indicated in Wakker (1986b, Theorem 4.4). For the case of three or more coordinates this follows straightforwardly from Theorem 2 below ($a_{-i} x_i \preccurlyeq a_{-i} x_i$ and $a_{-i} y_i \succ a_{-i} y_i$ give $x_i x_i \preccurlyeq_i^* y_i y_i$; (1) forbids $x_i x_i \succ_i^{**} y_i y_i$). \square

Analogues of the above theorem (and of the one below) for the case where coordinate sets are sets of probability distributions, are given in Fishburn (1965), Pollak (1967), and Keeney and Raiffa (1976, Theorem 6.4).

We say that \succ satisfies the **sure-thing principle** (other terms are strong separability, preferential/coordinate independence) if

$$\left[\begin{array}{l} \text{For all } i, a, b, x_i, y_i, \\ a_{-i} x_i \succ b_{-i} x_i \text{ implies} \\ a_{-i} y_i \succ b_{-i} y_i \end{array} \right] \text{ which is iff } \left[\begin{array}{l} \text{For no } i, x_i, y_i, \\ x_i x_i \succ_i^{**} y_i y_i \end{array} \right]. \tag{2}$$

Theorem 2. Let at least three coordinates be essential. There exists a continuous additively decomposable representing function V for \succ , if and only if \succ satisfies the sure-thing principle. This V is cardinal.

Proof. See Theorem 4.1 in Wakker (1986b), which generalizes Theorem 3 of Debreu (1960) by dropping the assumption of topological separability, and Theorem 6.14 of Krantz, Luce, Suppes and Tversky (1971) by also deriving continuity. \square

For the next property and theorem it is assumed that $\mathcal{C}_1 = \mathcal{C}_2 = \dots = \mathcal{C}_n = \mathcal{C}$ for some connected topological space \mathcal{C} . The present formulation by means of strength of preference has been indicated in section 3 of Wakker (1984a).

We say that \succsim satisfies **cardinal coordinate independence** (CCI) if

$$\left[\begin{array}{l} \text{For all } j, \text{ essential } i, \\ a, b, c, d \in X, \text{ and} \\ a, \beta, \gamma, \delta \in \mathcal{C}, \\ a_{-i}\alpha \preceq b_{-i}\beta \text{ and} \\ a_{-i}\gamma \succsim b_{-i}\delta \text{ and} \\ c_{-j}\alpha \succsim d_{-j}\beta \text{ imply} \\ c_{-j}\gamma \succsim d_{-j}\delta \end{array} \right] \text{ which is iff } \left[\begin{array}{l} \text{For all } j, \text{ essential } i, \\ \alpha, \beta, \gamma, \delta, \in \mathcal{C}: \\ \gamma\delta \succsim_i^* \alpha\beta \\ \text{then not} \\ \alpha\beta \succ_j^{**} \gamma\delta \end{array} \right]. \tag{3}$$

Note the difference between (3) and (1), apart from the unimportant detail of essentially of i . Formula (3) requires strength of preferences to be ‘non-contradictory’ across, possibly, different coordinates.

Theorem 3. Let $\mathcal{C}_i = \mathcal{C}$ for all i , and let at least two coordinates be essential. Then there exist non-negative p_1, \dots, p_n , summing to one, and a continuous $U: \mathcal{C} \rightarrow \mathbb{R}$, such that $x \mapsto \sum p_j U(x_j)$ represents \succsim , if and only if \succsim satisfies cardinal coordinate independence. U is cardinal, the p_1, \dots, p_n are unique.

Proof. See Wakker (1984a, Theorem 5.1). \square

When applied to the context of decision making under uncertainty, the above theorem gives a characterization of subjective expected utility maximization. It differs from Savage (1954) in dealing with a finite state space. Its analogue for the case where \mathcal{C} is a set of probability distributions as given in Anscombe and Aumann (1963). Anscombe and Aumann do not consider a Cartesian product of marginal distributions, but rather consider simultaneous distributions. Their Assumption 2 allows one to consider only the marginal distributions.

For the next property and theorem we shall assume that $\mathcal{C}_i = \mathbb{R}$ for every i . The following property reflects the idea of non-increasing marginal utility.

We say that \succsim satisfies the **concavity assumption** if

$$\left[\begin{array}{l} \text{For all } i, x, y, \\ \text{and real } \alpha, \beta, \epsilon \\ \text{with either } \alpha \geq \beta \\ \text{and } \epsilon \geq 0, \\ \text{or } \alpha \leq \beta \text{ and } \epsilon \leq 0, \\ x_{-i}\alpha \succsim y_{-i}\beta \text{ implies} \\ x_{-i}(\alpha - \epsilon) \succsim y_{-i}(\beta - \epsilon) \end{array} \right] \text{ which is iff } \left[\begin{array}{l} \text{For all } i \\ \text{and real } \alpha, \beta, \epsilon, \\ \text{with either } \alpha \geq \beta \\ \text{and } \epsilon \geq 0 \\ \text{or } \alpha \leq \beta \text{ and } \epsilon \leq 0, \\ \text{not } \alpha\beta \succ_i^{**} (\alpha - \epsilon)(\beta - \epsilon) \end{array} \right]. \tag{4}$$

Theorem 4. Let $\mathcal{C}_i = \mathbb{R}$ for all i , and let at least three coordinates be essential. Then there exists a concave additively decomposable representing function V for \succsim , if and only if \succsim satisfies the concavity assumption. This V is cardinal.

Proof. See Wakker (1986). \square

It is straightforwardly shown that in the above theorem ‘concave’ can be replaced by ‘strictly concave’, if in (4) for every $\epsilon \neq 0$ we have: not $\alpha\beta \succ_j^* (\alpha - \epsilon)(\beta - \epsilon)$. Yaari (1978) studied versions of (4) for $\alpha \neq \beta$ and $\epsilon \neq 0$.

In the following property and theorem, it is assumed, slightly more generally than Koopmans (1972), that $\mathcal{C}_1 = \dots = \mathcal{C}_n = \mathcal{C}$, for a connected topological space \mathcal{C} ; e.g., $\mathcal{C} = \mathbb{R}$, standing for money. Further we use the term ‘point of time’ for index. The property below is a reformulation of its version in Wakker (1984b).

We say that point of time j is CCI-related to point of time i if, for all $\alpha, \beta, \gamma, \delta$

$$\alpha\beta \succ_i^* \gamma\delta \Rightarrow \text{not } \gamma\delta \succ_j^{**} \alpha\beta. \tag{5}$$

Theorem 5. Let $n \geq 3$. Let all points of time $1, \dots, n$ be essential. Then the following two statements are equivalent for the continuous weak order \succ on \mathcal{C}^n :

- (i) There exists a continuous $U: \mathcal{C} \rightarrow \mathbb{R}$, and a λ in $(0, 1]$, such that $x \mapsto \sum_{j=1}^n \lambda^j U(x_j)$ represents \succ .
- (ii) Every point of time $j \geq 2$ is CCI-related to its predecessor $j - 1$; there exist $(\alpha, \dots, \alpha) \succ (\beta, \dots, \beta) \succ (\theta, \dots, \theta)$ such that, for every point of time $j < n$, $[(\theta, \dots, \theta)_{-j}\beta] = [(\theta, \dots, \theta)_{-(j+1)}\alpha]$; and $\gamma \succ_1^{**} \delta\delta$ for no $\gamma, \delta \in \mathcal{C}$.

Furthermore, the (‘discount factor’) λ in (i) is uniquely determined; U is cardinal.

Sketch of the proof. Since (i) \Rightarrow (ii) is straightforward, we assume (ii), and derive (i) and the ‘Furthermore’-statement. For every $j \geq 2$, and any x, γ, δ , $x_{-(j-1)}\delta \succ x_{-(j-1)}\delta$ and $x_{-(j-1)}\gamma \preccurlyeq x_{-(j-1)}\gamma$, hence by (5) $\gamma \succ_j^{**} \delta\delta$ for no $\gamma, \delta \in \mathcal{C}$. We may apply Theorem 2, to obtain a representing function $V: x \mapsto \sum V_j(x_j)$. We may set $V_j(\theta) = 0$ for all j .

Since every j is CCI-related to $j - 1$, there can be seen to exist λ_j such that $V_j = \lambda_j V_{j-1}$ for all j . This is much like in the proof of Theorem 2.1 in Wakker (1984a). In short, the idea is that, for every $j \geq 2$, and $\eta \in \mathcal{C}$, there exists an open neighbourhood of η on which V_j orders differences as V_{j-1} does it. Hence locally V_j can be seen to be a positive affine transformation of V_{j-1} . Then it must also be globally. Now $V_j(\theta) = 0 = V_{j-1}(\theta)$ implies the existence of λ_j as above.

Substitution of the preferences in (ii) concerning α, β , and θ , shows that, with $\lambda: = \lambda_2$, we have $\lambda_j = \lambda$ for all j ; and that $0 \leq \lambda \leq 1$. Since all points of time are essential, $\lambda = 0$ cannot be. Finally, set $U: = V_1/\lambda$.

The furthermore-statement follows from the cardinality of V in Theorem 2. \square

For an interpretation of (ii) above, say consequences are amounts of money, $\mathcal{C} = \mathbb{R}$. The CCI-relatedness condition says that revealed strengths of preferences concerning money on a point of time, should not contradict revealed strength of preferences on the previous point of time. For the condition concerning α, β , and θ , say θ is the status quo, receive \$0. Let α be some standardized amount of money, say \$1. The condition says that today’s equivalent β , for tomorrow’s receipt of \$1, should be the same on every today. This condition weakens Koopman’s stationary assumption.

One can replace $(0, 1]$ in (i) above by $(0, 1)$, if in (ii) one replaces $(\alpha, \dots, \alpha) \succ (\beta, \dots, \beta)$ by $(\alpha, \dots, \alpha) \succ (\beta, \dots, \beta)$.

4. Conclusion

It is desirable to formulate results, as much as possible, in terms of primitives of an adopted model. In decision theory the primitives are the set of alternatives, structure on this set, and the

preference relation of a decision maker. Derived notions (such as utility functions) should be adopted in the formulation of results, only if this achieves a significant intuitive gain. We hope that the derived strength of preference relations, as defined in this paper, do give a significant intuitive gain in the derivation of continuous cardinal utility.

References

- Anscombe, F.J. and R.J. Aumann, 1963, A definition of subjective probability, *Annals of Mathematical Statistics* 34, 199–205.
- Basu, K., 1982, Determinateness of the utility function: revisiting a controversy of the thirties, *Review of Economic Studies* 49, 307–311.
- Debreu, G., 1960, Topological methods in cardinal utility theory, in: K.J. Arrow, S. Karlin and P. Suppes, eds., *Mathematical methods in the social sciences* (Stanford University Press, Stanford, CA) 16–26.
- Fishburn, P.C., 1965, Independence in utility theory with whole product sets, *Operations Research* 13, 28–45.
- Grodal, B., 1978, Some further results on integral representation of utility functions (Institute of Economics, University of Copenhagen, Copenhagen). Forthcoming in: K. Vind, *Independent preferences*.
- Karni, E., D. Schmeidler and K. Vind, 1983, On state dependence preferences and subjective probabilities, *Econometrica* 51, 1021–1031.
- Keeney, R.L. and H. Raiffa, 1976, *Decisions with multiple objectives* (Wiley, New York).
- Koopmans, T.C., 1972, Representations of preference orderings with independent components of consumption; (and:) Representations of preference orderings over time, in: C.B. McGuire and R. Radner, eds., *Decision and organization* (North-Holland, Amsterdam) 57–100.
- Krantz, D.H., R.D. Luce, P. Suppes and A. Tversky, 1971, *Foundations of measurement, Additive and polynomial representations*, Vol. I (Academic Press, New York).
- Pollak, R.A., 1967, Additive von Neumann–Morgenstern utility functions, *Econometrica* 35, 485–494.
- Samuelson, P.A., 1958, An exact consumption–loan model of interest with or without the social contrivance of money, *The Journal of Political Economy* 66, 467–482.
- Savage, L.J., 1954, *The foundations of statistics* (Wiley, New York), Second ed., 1972 (Dover, New York).
- von Neumann, J. and O. Morgenstern, 1944, 1947, 1953, *Theory of games and economic behavior* (Princeton University Press, Princeton, NJ).
- Wakker, P.P., 1984a, Cardinal coordinate independence for expected utility, *Journal of Mathematical Psychology* 28, 110–117.
- Wakker, P.P., 1984b, Continuous expected utility for arbitrary state spaces, *Methods of Operations Research*, 50, 113–129.
- Wakker, P.P., 1986, Concave additively decomposable representing functions and risk aversion, in: L. Daboni, A. Montesano and M. Lines, eds., *Recent developments in the foundations of utility and risk theory* (Reidel, Dordrecht) 249–262.
- Zilcha, I., 1988, Intergenerational transfers, in: J. Eatwell and P. Newman, eds., *The new Palgrave: A dictionary of economic theory and doctrine* (Macmillan Press, Cambridge) forthcoming.