

An efficient multiplex genotyping approach for detecting the major worldwide human Y-chromosome haplogroups

Mannis van Oven · Arwin Ralf · Manfred Kayser

Received: 21 June 2011 / Accepted: 13 July 2011

© The Author(s). This article is published with open access at Springerlink.com 2011

Abstract The Y chromosome is paternally inherited and therefore serves as an evolutionary marker of patrilineal descent. Worldwide DNA variation within the non-recombining portion of the Y chromosome can be represented as a monophyletic phylogenetic tree in which the branches (haplogroups) are defined by at least one SNP. Previous human population genetics research has produced a wealth of knowledge about the worldwide distribution of Y-SNP haplogroups. Here, we apply previous and very recent knowledge on the Y-SNP phylogeny and Y-haplogroup distribution by introducing two multiplex genotyping assays that allow for the hierarchical detection of 28 Y-SNPs defining the major worldwide Y haplogroups. PCR amplicons were kept small to make the method sensitive and thereby applicable to DNA of limited amount and/or quality such as in forensic settings. These Y-SNP assays thus form a valuable tool for researchers in the fields of forensic genetics and genetic anthropology to infer a man's patrilineal bio-geographic ancestry from DNA.

Keywords Y chromosome · Y-SNP · Haplogroup · Patrilineal ancestry · Bio-geographic ancestry · Multiplex SNaPshot

Introduction

Knowledge about the bio-geographic ancestry revealed from crime-scene samples can be relevant for investigative intelli-

gence purposes in search for unknown sample donors who usually cannot be identified via conventional forensic STR profiling. DNA-based bio-geographic ancestry inference is also applied in genealogical and anthropological research for various purposes. The human Y chromosome is widely studied as an evolutionary marker of patrilineal descent. A well-established Y-chromosome phylogeny is available [6] and is continuously being expanded as novel SNPs are discovered. A wealth of data has been produced previously on the worldwide distribution and allele frequencies of numerous Y-SNPs and the respective Y haplogroups they define. Here, we take advantage of existing knowledge on the Y-SNP phylogeny and worldwide Y haplogroup distribution and introduce two Y-SNP multiplex assays, based on single-base primer extension (SNaPshot™) technology, for the detection of the major worldwide Y haplogroups. Together with well-known Y-SNPs, we have also included some relatively novel Y-SNPs such as M522 [4], M526 [4], P326 [8] and M412 [9], acknowledging most recent progress in Y-chromosome research.

Materials and methods

DNA samples

A subset of DNA samples from the HapMap 3 reference panel [1], belonging to various Y haplogroups, was obtained from the Coriell Institute for Medical Research (<http://www.coriell.org>).

Primer design

Primers were designed using Primer3Plus [11] with a T_m around 60°C for PCR primers and around 55°C for

M. van Oven (✉) · A. Ralf · M. Kayser
Department of Forensic Molecular Biology,
Erasmus MC University Medical Center Rotterdam,
3000 CA Rotterdam, The Netherlands
e-mail: m.vanoven@erasmusmc.nl

extension primers. Potential interactions between primers in the same multiplex were evaluated with the AutoDimer version 1.0 software [12]. In order to minimize allelic dropouts due to primer mismatches, we avoided as much as possible that primer-annealing sites overlapped with known Y-chromosome polymorphisms. Extension primers were varied in length through the addition of 5' non-homologous poly(GACT) tails to ensure electrophoretic separation of extended fragments.

PCR amplification

Multiplex PCR amplification was carried out in a reaction volume of 6 μ L, containing 1 \times GeneAmp PCR Gold Buffer (Applied Biosystems, CA, USA), 4.5 mM MgCl₂ (Applied Biosystems), 100 μ M of each dNTP (Roche, Mannheim, Germany), 0.35 units of AmpliTaq Gold DNA polymerase (Applied Biosystems), 1–2 ng of genomic DNA template, and PCR primers (desalted; Metabion, Martinsried, Germany) in concentrations as specified in Tables 1 and 2. The reactions were performed in a Dual 384-well GeneAmp PCR System 9700 (Applied Biosystems) using the following cycling conditions: 10 min at 95°C, followed by 30 cycles of 94°C for 15 s, 60°C for 45 s, and a final extension at 60°C for 5 min. PCR products were purified by adding 2 μ L ExoSAP-IT (USB Corporation, OH, USA) to 6 μ L PCR product, followed by incubation at 37°C for 30 min and 80°C for 15 min.

Single-base extension

Multiplex single-base primer extension was carried out in a reaction volume of 6 μ L, containing 1 μ L SNaPshot™ Ready Reaction Mix (Applied Biosystems), 1 μ L purified PCR product, and extension primers (HPLC-purified; Metabion, Martinsried, Germany) in concentrations as specified in Tables 1 and 2. The reactions were performed in a Dual 384-well GeneAmp PCR System 9700 (Applied Biosystems) using the following cycling conditions: 2 min at 96°C, followed by 25 cycles of 96°C for 10 s, 50°C for 5 s, and 60°C for 30 s. The reaction products were purified by adding 1 unit of Shrimp Alkaline Phosphatase (USB Corporation) to 6 μ L of extension product, followed by incubation at 37°C for 45 min and 75°C for 15 min.

Capillary electrophoresis

The extended fragments were separated and detected by capillary electrophoresis on a 3130xl Genetic Analyzer (Applied Biosystems) using POP-7 polymer. A mixture of 1 μ L purified extension product, 8.7 μ L Hi-Di formamide (Applied Biosystems) and 0.3 μ L GeneScan-120 LIZ

internal size standard (Applied Biosystems) was run with 10 s injection time at 1.2 kV and 500 s run time at 15.0 kV. Results were analysed using GeneMapper version 3.7 software (Applied Biosystems).

Results and discussion

Two genotyping multiplex assays were developed targeting a total of 28 Y-SNPs that define the major worldwide Y-chromosome haplogroups (Fig. 1). During the course of this work, a paper was published that reported a reorganization of the deepest clades of the Y-chromosome phylogeny, one of the consequences being that marker M91 no longer defines a monophyletic haplogroup A, but rather should be placed on the stem leading to the BCDEF (also referred to as BT) clade [5]. We have incorporated this change in our tables and figures to conform with the latest Y-chromosome topology. Furthermore, we took advantage of some recently discovered Y-SNPs (P326, M526, M522 and M412) that, as far as we know, were not included in previous Y genotyping systems [e.g. 2, 3, 10]. Of these novel SNPs, P326 (also known as L298) defines a new branch that joins haplogroups L and T into a single clade now called LT [8]. M526 is located downstream of marker M9 and encompasses haplogroups K1 to K4 as well as M to S [4]; the branch defined by M526 is now referred to as haplogroup K, and the former haplogroup K (defined by M9) is now relabelled as KLT. M522 (also known as L16 or S138) defines a new node within haplogroup F that encompasses haplogroups I, J and KLT [4] and is referred to as haplogroup IJKLT. M412 (also known as L51 or S167) defines a significant subhaplogroup within haplogroup R that is most abundant in western parts of Europe [9].

The 28 Y-SNPs were divided into two multiplexes such as to allow a hierarchical typing strategy. Multiplex 1 covers haplogroups BCDEF, B, C, DE, D, E, F, F3, G, H, IJKLT, I, J and KLT. If a sample is found to belong to the latter, it can subsequently be typed with multiplex 2 which covers haplogroups KLT, K, K1, K2, K3, K4, M, N, O, P, Q, R, R-M412 (also known as R1b1a2a1a), S and LT.

To maintain high sensitivity of the multiplexes, PCR amplicons were kept short with an average length of 103 bp (minimum, 46 bp; maximum, 178 bp). The recommended amount of template DNA for the PCR reactions is 1–2 ng, which gives satisfactory results when the DNA is of reasonable quality (Fig. 2). Although we did not further evaluate the sensitivity of the two multiplex assays, we expect that in many cases, lower amounts of template DNA will still yield informative genotypes.

Table 1 Genotyping details of Y-SNP multiplex 1

Locus	Mutation	PCR amplification		Single-base extension					
		Primer sequences (5'-3')	Conc. (μM)	Amplicon size (bp)	Primer sequence (5'-3') (5' aspecific tail in lowercase italics)	Conc. (μM)	Length (nt)	Orientation	Alleles (dye)
M91	ins T	F	CAAAAATCCCTTACATTCG	144/143	<i>g</i> CTACAGTAGTGAACCTGATTAATAAAAAA	0.300	28	R	a (yellow), i (green) ^a
		R	GCAGTGCCCTTCCAAATAAA						
M60	ins T	F	TCTTTACATTTCAAATGCATGACT	128/129	<i>ct(gact)</i> ₆ TAACCACCTGTGTGCCTGAT	0.600	45	R	a (yellow), i (green) ^a
		R	GAGAAGGTGGTGGTCAAGA						
M145	G->A	F	GCATACTTGCCTCCACGACT	96	<i>ct(gact)</i> _{3gac} TAGGCTAAGGCTGGCTCT	0.450	35	R	G (yellow), A (red)
		R	CCTCCCACTCCTTTTGGAT						
M174	T->C	F	TCTCCGTACACAGCAAAAATG	178	<i>ct(gact)</i> _{3g} ATACCTTCTGGAGTGCCTC	0.100	41	F	T (red), C (yellow)
		R	AGGAGAAGGACAAAGACCCATC						
M96	G->C	F	TGAGCTGTGATGTGTAACCTTGG	117	<i>act(gact)</i> _{10gac} TGGAAAAACAGGTCTCTCATAATA	0.200	69	F	G (blue), C (yellow)
		R	CACCCACTTTGTGCTTTGT						
M216	C->T	F	CCTCAACCAGTTTTATGAAGCTA	102	<i>ct(gact)</i> _{10g} CTGGTAGTATGTATACCTGTTGAAT	0.075	53	R	C (blue), T (green)
		R	TTCTAAATCTGAATCTGACACTGC						
M89	C->T	F	CAGCTTCCCTGGATTCAGCTC	105	<i>ct(gact)</i> _{13ga} AACTCAGGCCAAAGTGAGAGAT	0.300	77	R	C (blue), T (green)
		R	CACCTTGGGTCCAAGATCAC						
M282	A->G	F	TGTGCAACCTCAACTTTGCTT	106	<i>t(gact)</i> ₁₅ GAAAAGCAAAATCTCAATATGATAA	1.000	85	F	A (green), G (blue)
		R	TGTGATCAACTTCTTCCCTCA						
P257	G->A	F	ACCCCTCAGTCTCTCCGAT	71	<i>(gact)</i> _{3g} ATATCCCACTGCAITTTCTG	0.300	57	F	G (blue), A (green)
		R	TCATCTCCAACCCCAATCT						
M69	T->C	F	GGAGGCTGTTTACACTCCTGA	87	<i>(gact)</i> _{10g} GGCTGTTTACACTCCTTGAAA	0.150	61	F	T (red), C (yellow)
		R	TCTCCCTTAGCTCTCCTGTT						
M522	G->A	F	TCCAAITCCCATGTCCTCTC	109	<i>t(gact)</i> ₁₁ CTACTACGCCCTCTCTTTGTCC	0.075	65	F	G (blue), A (green)
		R	CAGTGCAGAAAATCACGGTAGA						
M258	T->C	F	TTCAGGATTTGTCAAGGATGG	108	<i>t(gact)</i> _{3gac} GGGATTCCTCAAGTTCCCA	0.300	33	R	T (green), C (blue)
		R	GCTATGACTAAGAGGGATTCCAA						
M304	A->C	F	TTGTAAACAAAACAGTATGTGGGATTT	88	<i>act(gact)</i> _{11ga} TTATACCAAAAATATCACCAGTTTGT	0.300	73	R	A (red), C (blue)
		R	CGTCTTAACCAAAAATATCACCAGTT						
M9	C->G	F	CTGCAAAGAAACGGCCTAAG	90	<i>t(gact)</i> _{7g} CCGCCTAAGATGGTTGAATCGTTT	0.100	49	F	C (yellow), G (blue)
		R	AACTAAGTATGTAAGACATGAA						

^a *a* ancestral, *i* insertion

Table 2 Genotyping details of Y-SNP multiplex 2

Locus	Mutation	PCR amplification		Single-base extension					
		Primer sequences (5'-3')	Conc. (μM)	Amplicon size (bp)	Primer sequence (5'-3') (5' aspecific tail in lowercase italics)	Conc. (μM)	Length (nt)	Orientation	Alleles (dye)
M9	C->G	F CTGCAAAAGAAACGGCCTAAG R AACTAAGTATGTAAGACATGGAACGTTTG	0.300 0.300	90	<i>t(gact)</i> _{9g} CCGCCCTAAAGATGGTTGAAT	0.100	49	F	C (yellow), G (blue)
M526	A->C	F TAGAGGCAGGGTGTGCTCT R TACTTTGGGAGGCTGCTGT	0.300 0.300	100	<i>ct(gact)</i> _{10ga} TGTCAATCAGGCTGAATCATAC	0.450	65	F	A (green), C (yellow)
M147	ins T	F CCTGAATAAGCTGGTGAAGAAA R GGAGACCCCTGCTCTGAAAGAA	0.500 0.500	114/115	<i>ct(gact)</i> _{11ga} CCTGTCTCTGAAAGAAAAAA	1.000	69	R	a (yellow), i (green) ^a
P308	C->T	F GCTACCAATACCCCAAGA R CCTGGAATATGGACGAAAT	0.050 0.050	108	<i>gactgac</i> GAAATGATTAAGTAGGCCTTCT	0.150	31	R	C (blue), T (green)
P79	T->C	F TTGCTTAGTATAATGCTTTTCATGCTC R AAATGAGGCTAATCAATGGAACA	0.500 0.500	101	<i>ct(gact)</i> _{9g} TGCTCAATCGCAATCTTTG	1.000	37	F	T (red), C (yellow)
P261	G->A	F TCCTAGAAGGTAACCCACTACCC R TGTGCATATGTTATCCCAATGT	0.500 0.500	93	<i>t(gact)</i> _{7gac} TTTTGTGTTTTATTAATGAAATGCTA	1.000	57	R	G (yellow), A (red)
P256	G->A	F TCTTGGTTTTCCCATGACC R CATCTCCCAACTTGTCTGTGC	0.200 0.200	91	<i>t(gact)</i> _{13ga} TGCCCTACACTAGATAGAAAAGG	0.150	77	F	G (blue), A (green)
M231	G->A	F AACACAATTACTGTTTCTACTGCTTTC R TTCACATCATCCAGTACAGCA	0.300 0.300	119	<i>act(gact)</i> _{3g} CGAATCTTCCCCCAAT	0.450	33	R	G (yellow), A (red)
M175	5 bp del	F CCCAAATCAACTCAACTCCAG R TTCTACTGATACCTTTGTTCTGTTCA	0.300 0.300	101/96	<i>t(gact)</i> ₁₀ CACAATGCCCTTCTCACTTCTC	0.600	61	F	a (red), d (green) ^a
M45	G->A	F CATCGGGGTGTGGACTTTA R CCTCAGAAAGGAGCTTTTTGC	0.400 0.400	109	<i>act(gact)</i> _{6g} AATTTGGCAGTGAAAAAAT ATAGATA	0.750	53	F	G (blue), A (green)
M242	C->T	F AAAAAAGGTGACCAAGGTGCT R AAAAAACACGTTAAGACCAATGC	0.400 0.400	46	<i>ct(gact)</i> _{9g} CGTTAAGACCAATGCCAA	0.100	45	R	C (blue), T (green)
M207	A->G	F GGGGCAAAATGTAAGTCAAGC R TCACTTCAACCTCTTGTGGAA	0.300 0.300	83	<i>(gact)</i> _{14g} AATGTAAGTCAAGCAAGA AATTTA	0.300	81	F	A (green), G (blue)
M412	G->A	F GGCACCTCCCGTCACTT R GGTGAAGTGGACCCATCCA	0.300 0.300	114	<i>ct(gact)</i> ₁₆ GGGTACAATCTGATGAGGC	0.300	85	F	G (blue), A (green)
P202	T->A	F AAACCTCCCAAGTTTGTGGTTC R TGATCCCTTAATAATAGCAAGACC	0.300 0.300	125	<i>ct(gact)</i> _{12ga} CCAGTTTGTGGTCTTTTGTGA	0.300	73	F	T (red), A (green)
P326	T->C	F TTCAGATATCAGGCCGCTT R GAGCTGTCAAGCCTGCCTAAG	0.200 0.200	61	<i>t(gact)</i> ₃ CCTAAGCAGAGGAAAAATA GTACAG	0.150	37	R	T (green), C (blue)

^a a ancestral, i insertion, d deletion




Fig. 1 Y-SNP marker phylogeny, inferred haplogroups and their geographic distributions as covered by the two Y-SNP multiplex assays introduced here. The phylogeny shown is a truncation of the


entire Y-chromosome tree. Due to the fact that some haplogroups have so far been observed only sporadically, their regions of occurrence are less certain and are therefore shown in *parentheses*

The multiplexes were optimized on a Genetic Analyzer using POP-7 polymer. We noticed in the past that the type of POP polymer has some influence on the relative electrophoretic mobilities as well as peak intensities of the extended fragments. Therefore, re-adjustment of 5' tail lengths as well as reaction concentrations of the extension primers might be necessary when employing a POP polymer that is different from the one used here.

Conclusion

The multiplex assays presented here form a convenient tool for detecting the major worldwide Y haplogroups, hence giving a first idea about the patrilineal bio-geographic ancestry of men, being of relevance in forensic investigation and anthropological research. Notably, for most of the haplogroups covered here, more detailed phylogenetic resolution can be obtained by

Multiplex 1



Multiplex 2




Fig. 2 Typical electropherograms obtained with the two Y-SNP multiplex assays introduced here, using DNA samples belonging to a range of Y haplogroups. For each peak, the detected allele is

indicated in concordance with Tables 1 and 2. As is convention, the yellow dye is shown as *black* for better contrast

genotyping additional Y-SNPs. Hence, we foresee that additional multiplex assays, targeting more downstream Y-SNPs and dedicated to the dissection of particular (sub) haplogroups, will form useful additions to the global assays presented here. For more complete reconstruction of a person's

overall bio-geographic ancestry, we recommend that Y-chromosome markers are combined with ancestry-informative markers from mitochondrial DNA and autosomal DNA, as already achievable with efficient multiplex tools offering resolution on a continental level [e.g. 7, 13].

Acknowledgements We thank Qasim Ayub and Chris Tyler-Smith for help in providing access to DNA samples belonging to various Y haplogroups. This work was supported in parts by funding provided by the Netherlands Forensic Institute (NFI) and by the Netherlands Genomics Initiative (NGI)/Netherlands Organization for Scientific Research (NWO) within the framework of the Forensic Genomics Consortium Netherlands (FGCN).

Open Access This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

References

1. Altshuler DM, Gibbs RA, Peltonen L, International HapMap 3 Consortium et al (2010) Integrating common and rare genetic variation in diverse human populations. *Nature* 467(7311):52–58
2. Berniell-Lee G, Sandoval K, Mendizabal I, Bosch E, Comas D (2007) SNPlexing the human Y-chromosome: a single-assay system for major haplogroup screening. *Electrophoresis* 28(18):3201–3206
3. Brión M, Sanchez JJ, Balogh K, Thacker C, Blanco-Vérea A, Børsting C, Stradmann-Bellinghausen B, Bogus M, Syndercombe-Court D, Schneider PM, Carracedo A, Morling N (2005) Introduction of a single nucleotide polymorphism-based “Major Y-chromosome haplogroup typing kit” suitable for predicting the geographical origin of male lineages. *Electrophoresis* 26(23):4411–4420
4. Chiaroni J, Underhill PA, Cavalli-Sforza LL (2009) Y chromosome diversity, human expansion, drift, and cultural evolution. *Proc Natl Acad Sci USA* 106(48):20174–20179
5. Cruciani F, Trombetta B, Massaia A, Destro-Biol G, Sellitto D, Scozzari R (2011) A revised root for the human Y chromosomal phylogenetic tree: the origin of patrilineal diversity in Africa. *Am J Hum Genet* 88(6):1–5
6. Karafet TM, Mendez FL, Meilerman MB, Underhill PA, Zegura SL, Hammer MF (2008) New binary polymorphisms reshape and increase resolution of the human Y chromosomal haplogroup tree. *Genome Res* 18(5):830–838
7. Lao O, Vallone PM, Coble MD, Diegoli TM, van Oven M, van der Gaag KJ, Pijpe J, de Knijff P, Kayser M (2010) Evaluating self-declared ancestry of U.S. Americans with autosomal, Y-chromosomal and mitochondrial DNA. *Hum Mutat* 31(12):E1875–E1893
8. Mendez FL, Karafet TM, Krahn T, Ostrer H, Soodyall H, Hammer MF (2011) Increased resolution of Y chromosome haplogroup T defines relationships among populations of the Near East, Europe, and Africa. *Hum Biol* 83(1):39–53
9. Myres NM, Rootsi S, Lin AA, Järve M, King RJ, Kutuev I, Cabrera VM, Khusnutdinova EK, Pshenichnov A, Yunusbayev B, Balanovsky O, Balanovska E, Rudan P, Baldovic M, Herrera RJ, Chiaroni J, Di Cristofaro J, Vilems R, Kivisild T, Underhill PA (2011) A major Y-chromosome haplogroup R1b Holocene era founder effect in Central and Western Europe. *Eur J Hum Genet* 19(1):95–101
10. Onofri V, Alessandrini F, Turchi C, Pesaresi M, Buscemi L, Tagliabracci A (2006) Development of multiplex PCRs for evolutionary and forensic applications of 37 human Y chromosome SNPs. *Forensic Sci Int* 157(1):23–35
11. Untergasser A, Nijveen H, Rao X, Bisseling T, Geurts R, Leunissen JA (2007) Primer3Plus, an enhanced web interface to Primer3. *Nucleic Acids Res* 35:W71–W74
12. Vallone PM, Butler JM (2004) AutoDimer: a screening tool for primer-dimer and hairpin structures. *Biotechniques* 37(2):226–231
13. van Oven M, Vermeulen M, Kayser M (2011) Multiplex genotyping system for efficient inference of matrilineal genetic ancestry with continental resolution. *Investig Genet* 2:6