

REVIEW

Review of health-related quality of life data in multiple myeloma patients treated with novel agents

P Sonneveld^{1,7}, SG Verelst^{1,7}, P Lewis², V Gray-Schopfer³, A Hutchings⁴, A Nixon⁵ and MT Petrucci⁶

In multiple myeloma (MM), health-related quality of life (HRQoL) data is becoming increasingly important, owing to improved survival outcomes and the impact of treatment-related toxicity on HRQoL. Researchers are more frequently including HRQoL assessments in clinical trials, but analysis and reporting of this data has not been consistent. A systematic literature review assessed the effect of novel agents (thalidomide, bortezomib and lenalidomide) on HRQoL in MM patients, and evaluated the subsequent reporting of these HRQoL results. A relatively small body of literature addresses HRQoL data in MM patients treated with novel MM therapeutic agents: 9 manuscripts and 15 conference proceedings. The literature demonstrates the complementary value of HRQoL when assessing clinical response, progression, overall survival and toxicity. However, weaknesses and inconsistencies in analysis and presentation of HRQoL data were observed, often complicating interpretation of the impact of treatment on HRQoL in MM. Further evaluation of HRQoL in MM patients treated with novel agents is required in larger cohorts, and ideally in head-to-head comparative studies. Additionally, the development of standardised MM-specific best practice guidelines in HRQoL data collection and analysis is recommended. These would ensure that future data are more useful in guiding predictive models and clinical decisions.

Leukemia (2013) 27, 1959–1969; doi:10.1038/leu.2013.185

Keywords: multiple myeloma; quality of life; thalidomide; bortezomib; lenalidomide

INTRODUCTION

Multiple myeloma (MM), a clonal proliferation of plasma cells, is an incurable disease in which patients often have pronounced symptoms and substantially reduced health-related quality of life (HRQoL).¹ Eighty percent of patients experience skeletal destruction,^{2,3} ~73% will have anaemia at diagnosis⁴ and ~30% of patients present with renal insufficiency.⁵ Impaired immune function is also an important characteristic of the disease that leads to severe infections.^{6,7} Treatment of MM has improved substantially in recent years, leading to prolonged overall survival upon introduction of high-dose chemotherapy combined with autologous stem cell transplantation (ASCT) and use of agents such as thalidomide, bortezomib and lenalidomide.^{8,9} Despite therapeutic advances, survival prognosis remains poor, with a 5-year relative survival rate of 35–37% in newly diagnosed multiple myeloma (NDMM) patients, although substantial improvements have been observed in patients up to 59 years of age. Survival rates and prognoses decrease with patient age (≥ 70 years).^{10–13} Bortezomib monotherapy and lenalidomide in combination with dexamethasone are both licensed for the treatment of MM patients who have undergone at least one prior therapy. Both therapies have shown statistically and clinically significant improvements over dexamethasone alone in terms of response rates, time to progression and overall survival.^{14–18} High-dose chemotherapy with ASCT is the standard of care for transplant-eligible, NDMM patients, typically those younger than 65 years of age.^{19,20} Novel agents, such as thalidomide, lenalidomide and bortezomib, figure prominently in the treatment of these NDMM

patients, and are being used in combination with vincristine, adriamycin and/or dexamethasone/low-dose dexamethasone.^{21–23} These novel agents have also shown efficacy as maintenance treatment post-ASCT, although no drug is currently licensed in this indication.^{22,24–27}

Novel agents have also changed MM management in elderly patients not eligible for ASCT.²⁸ Combinations of melphalan, prednisone and thalidomide (MPT)^{29–31} and of bortezomib, melphalan and prednisone (VMP)^{32,33} have shown improved progression-free survival and overall survival compared with melphalan and prednisone (MP) alone. Both MPT and VMP combinations are licensed in Europe for treatment of non-ASCT-eligible NDMM patients and are the current standards of care for elderly patients.²⁰ Data also support the role of the combination of melphalan, prednisone and lenalidomide followed by lenalidomide maintenance (MPR-R) in the treatment of elderly non-ASCT-eligible patients,^{34,35} although it is not currently licensed.

While many of these novel agents have demonstrated improved survival rates, they are also associated with some adverse events (AEs), which can impact on patient HRQoL.^{36,37} In a prospective study of 154 MM patients in the United Kingdom and Germany, the impact of specific MM symptoms and AEs were correlated with HRQoL scores after adjusting for the level of disease severity. While severe bone pain and being severely symptomatic had the most deleterious effect on patient HRQoL, patients who were receiving MM treatment also reported lower HRQoL, related to treatment toxicity.³⁸ It is therefore important to

¹Department of Hematology, Erasmus University Medical Center Rotterdam, Rotterdam, The Netherlands; ²Celgene GmbH, Munich, Germany; ³OmniScience SA, Geneva, Switzerland; ⁴Global Market Access Solutions, London, UK; ⁵Oxford Outcomes, an ICON plc. company, Oxford, UK and ⁶Sapienza University of Rome, Rome, Italy. Correspondence: SG Verelst, Department of Hematology, Erasmus University Medical Centre Rotterdam, Room Na817, PO Box 2040, 3000 CA Rotterdam, The Netherlands. E-mail: s.verelst@erasmusmc.nl

⁷These authors contributed equally to this work.

Received 8 March 2013; revised 11 June 2013; accepted 12 June 2013; accepted article preview online 20 June 2013; advance online publication, 16 July 2013

fully understand the effects of treatments on patients' HRQoL. The increasing number of trials evaluating HRQoL in haematological malignancies is a testament to the growing importance of patient-reported outcomes.^{39–42}

The aims of MM treatment are to control disease, prolong survival and maximise patient wellbeing. HRQoL instruments can be incorporated into clinical studies in order to get a more comprehensive evaluation of treatment outcomes. However, the value of HRQoL data in guiding clinical practice depends upon the quality and comparability of these data.

Careful consideration should be given to the study design, including HRQoL instrument selection, a statistical analysis plan and the reporting of results.^{43,44} Indeed, accumulating evidence indicate that published clinical trials assessing HRQoL have failed to meet good scientific standards of reporting, and internationally agreed upon standards have been called for.^{45,46}

HRQoL data are now routinely captured in studies of new treatments for MM. The objective of this publication was to review available HRQoL data for the newer MM treatments and to critically evaluate the standards of HRQoL data collection, analysis and reporting. It is hoped that our review will help guide the standardisation of HRQoL data collection, analysis and reporting, to work towards either internationally agreed upon guidelines or incorporation into future MM studies.

MATERIALS AND METHODS

A systematic literature review was performed to identify relevant articles pertaining to the treatment-related HRQoL impact of thalidomide, bortezomib or lenalidomide in MM patients. The following electronic bibliographic and treatment guideline databases were searched: EMBASE, PubMed, National Guideline Clearinghouse Cochrane Database of Systematic Reviews, ClinicalTrials.gov and ClinicalStudyResults.org clinical trial registers (January 2000 through 31 December 2012). Supplementary searches included oncology and haematology conference proceedings (European Hematology Association (EHA), American Society of Clinical Oncology (ASCO), American Society of Hematology (ASH) and International Myeloma Working Group (IMW)), grey literature and reference lists of key papers. The search was restricted to documentation published in human subjects. All interventional study designs involving patients with MM were included, with the exception of case studies. Greater emphasis was placed on randomised controlled trial (RCT) data. Review articles were excluded.

The literature search identified 420 potential publications. After the abstracts for each study were reviewed against the inclusion/exclusion criteria by two independent contributors, 9 manuscripts and 15 conference proceedings (7 ASH, 5 EHA, 2 ASCO and 1 IMW) were considered relevant. Reasons for exclusion included insufficient HRQoL data and non-relevant clinical intervention. The relatively limited amount of retrieved literature was anticipated, given that these therapies are fairly recent and are supported by a limited number of clinical trials that have derived HRQoL analysis. The most frequently used HRQoL instruments in the MM studies included in the current review are briefly outlined in Box 1 and Table 1.

Studies reporting on HRQoL in MM patients treated with thalidomide

Two clinical trials evaluating MPT vs MP,^{47–49} and one study evaluating dexamethasone combined with thalidomide (TD) or bortezomib (VD)⁵⁰ were retrieved. Key HRQoL results from these trials are summarised in Table 2.

HRQoL data from the HOVON49 phase III study. The HOVON49 randomised, multi-centre, open-label, phase III trial compared MP ($n = 168$) with MPT ($n = 165$) followed by thalidomide maintenance in elderly (> 65 years) NDMM patients.⁵¹ The MPT regimen has become a standard treatment in this population, based on data from five RCTs.^{29,30,48,51–54} Eight treatment cycles were planned in the HOVON49 study. Patients who completed the planned MPT cycles received thalidomide maintenance therapy until progression, whereas patients in the MP arm received no maintenance. After disease progression (PD) or no response, salvage therapy was given according to the physician's choice. HRQoL was evaluated as a secondary end point in the study using the QLQ-C30 and QLQ-MY24 questionnaires

Box 1. HRQoL in MM and overview of commonly used HRQoL instruments

- **QLQ-C30:** 30-item, self-administered HRQoL patient questionnaire is designed for use in cancer patients. It has a one-week recall period and is composed of multi-item and single scales, including five functional scales (Physical, Role, Emotional, Social, and Cognitive Functioning), three symptom scales (Fatigue, Nausea/Vomiting and Pain) and a Global Health Status (GHS)/Quality of Life (QoL) scale. Single items include Dyspnoea, Insomnia, Appetite Loss, Constipation, Diarrhoea and Financial Difficulties. The QLQ-C30 has demonstrated good psychometric properties in solid tumours^{85,86} and reliability and validity in MM patients^{87–90}. The Minimal Important Difference (MID), defined as the smallest change in an HRQoL score considered important to patients that would lead the patient or clinician to consider a change in therapy,⁹¹ has been estimated for QLQ-C30 in MM patients.⁵⁶
- **QLQ-MY24 and QLQ-MY20:** QLQ-MY24 was developed as an addition to QLQ-C30 for specific use in MM.^{92,93} The QLQ-MY24 was refined to 20 items (QLQ-MY20). It has a 1-week recall period and addresses four important domains in MM: Disease Symptoms, Side Effects of Treatment, Body Image and Future Perspectives. The psychometrics, including reliability and validity of QLQ-MY20, in MM have been published.⁹²
- **QLQ-CIPN20:** the reliability, validity and responsiveness of the 20-item QLQ-CIPN20 instrument for patient-reported chemotherapy-induced peripheral neuropathy (CIPN) is currently being investigated in a wide range of oncology patient populations. The instrument has three subscales: a sensory, motor and autonomic subscale.⁹⁴
- **FACT-Multiple Myeloma:** a 14-item disease-specific FACT-MM HRQoL measure has been developed and has demonstrated acceptable psychometric properties.^{95,96}
- **FACIT-Fatigue:** bolt-on module to the base FACT-General (FACT-G) HRQoL questionnaire, a 27-item instrument measuring Well being (Physical, Social/Family, Emotional and Functional), with a recall period of 1 week.⁹⁷ The MID has been estimated and verified in a group of mixed diagnosis cancer patients.⁹⁸ No reliability, validity or other psychometric properties of the FACIT-Fatigue specific to MM have been described.
- **FACT-NTx:** another FACT-G bolt-on module with additional neurotoxicity parameters. It has demonstrated good psychometric properties in women with ovarian cancer,⁹⁹ but has only been used in a limited number of MM studies, similarly to FACIT-Fatigue.
- **EQ-5D:** standardised generic HRQoL questionnaire that can be converted into a 'health utility' score questionnaire used to measure health outcomes. The MID has recently been described for EQ-5D in MM patients.¹⁰⁰

(Table 1). HRQoL was measured at five pre-determined time points during the course of treatment.⁴⁷

Both treatment arms resulted in improved overall Global Health Status (GHS)/Global QoL, Fatigue, Side Effects of Treatment, Pain, Insomnia and Appetite Loss scores, although differences in favour of MPT were observed for these latter three scores. However, MPT was associated with a significant increase in paraesthesia from post-induction onwards, consistent with a cumulative dose-dependent effect of thalidomide.^{51,55} The higher incidence of constipation and paraesthesia with MPT vs MP was

Table 1. HRQoL instruments used in MM studies

Trial (Reference)	HRQoL instruments	Timing of HRQoL assessments
<i>Thalidomide</i> HOVON49 ⁴⁷	QLQ-C30, QLQ-MY24	Measurements for QoL were collected at study entry, after cycle 3 (~3 months after the start of cycle 1), after cycle 8 (~9 months after the start of cycle 1), at 12 and 18 months after the start of cycle 1
Waage <i>et al.</i> ; Guldbrandsen <i>et al.</i> ^{48,49} Hjorth <i>et al.</i> ⁵⁰	QLQ-C30 QLQ-C30	The questionnaires were completed at inclusion and later posted to patients every third month throughout the study Questionnaires were completed before randomisation (before start of treatment) and mailed to the patients after 6 and 12 weeks, after 6 months and every 6 months thereafter until the end of the study
<i>Bortezomib</i> APEX ^{60,65}	QLQ-C30, FACT-Ntx	Questionnaires were administered at baseline and at weeks 6, 12, 18, 24, 30, 36 and 42
SUMMIT ^{58,59,63,64}	QLQ-C30, QLQ-MY24, FACIT-Fatigue, FACT-Ntx	Patients completed the questionnaires during screening, on day 1 of cycles 3, 5 and 7 of treatment, as well as at the end of the study
VISTA ^{62,66,67}	QLQ-C30	Patients completed the questionnaire at screening, on day 1 of each cycle during the treatment phase and every 8 weeks until progression during follow-up
UPFRONT ⁶⁸⁻⁷¹	QLQ-C30	Patients completed the questionnaire before dosing on day 1 of cycle 1 (baseline), before dosing on day 1 of every odd-numbered cycle, at the end-of-treatment visit and every 12 weeks thereafter
<i>Lenalidomide</i> MM-015 ⁷³⁻⁷⁶	QLQ-C30, QLQ-MY20	Questionnaires were completed at baseline, at the beginning of every third cycle (cycles 4, 7, 10, 13 and 16), at study discontinuation and every 6 months in long-term follow-up
MM-018 ⁷⁷⁻⁷⁹	QLQ-C30, QLQ MY20	Questionnaires were completed at baseline and at week 24

Abbreviations: HRQoL, health-related quality of life; MM, multiple myeloma.

not reflected in overall HRQoL. No unfavourable overall difference in self-reported side effects between the two arms was observed during the study protocol. Verelst *et al.* explored whether the improvement in HRQoL from baseline seen was clinically significant. The MM-specific minimally important difference (MID) for the QLQ-C30 was defined as a difference of 6–17 points, as estimated by Kvam *et al.*⁵⁶ MID is defined as the smallest change in an HRQoL score considered important to patients that would lead the patient or clinician to consider a change in therapy. It was concluded that a clinically significant difference was observed for GHS/QoL, Role Functioning, Emotional Functioning, Social Functioning, Fatigue and Pain at the end of post-induction (18 months, during the period following the first week after induction treatment until the start of the next treatment off protocol) in favour of MPT-treated patients compared with MP. This prospective study showed that the higher frequency of toxicity associated with MPT did not translate into a negative effect on HRQoL and that patients on MPT have a better outlook. The authors concluded that MPT improved clinical outcome with no reduction in HRQoL (Table 2).

Critical review of HOVON49 HRQoL data. The use of a prospective design allowed the evaluation of HRQoL at different time points and treatment stages, with repeated measurements, owing to a linear mixed statistical model, which took into account correlation between measurements from the same patients. The model allowed a clear distinction between differences in the two randomisation arms at baseline and those possibly caused by additional thalidomide treatment. Thus, differences present at baseline did not generate significant interaction with time.

The HRQoL questionnaires were first completed by patients before treatment but after randomisation. The HOVON49 study authors hypothesised that anticipation of receiving beneficial treatment may have biased HRQoL reporting. As the study was open-labelled, patients in the MPT arm may score better at baseline. Indeed, at baseline, the MPT cohort had statistically significantly higher HRQoL scores for QLQ-C30 Emotional Functioning and GHS/QoL subscales, and for QLQ-MY24 Future Perspectives and Social Support subscales.

The quality of the data may be diminished by the open-label study design, which risks incorporating bias, and is further limited by the fact that not all HOVON49 trial patients participated in the HRQoL survey. Compliance rates with completing questionnaires at different study time points were not reported. Furthermore, although patients were evaluated at pre-determined treatment time points, not all patients completed

questionnaires at the same time. Finally, although clinically meaningful MID thresholds for MM were applied,⁵⁶ within-trial and domain-specific distribution-based MID estimates would have provided additional insight into the clinically meaningful changes.⁴³

HRQoL data from the Nordic Myeloma Study Group (NMSG). Waage *et al.* conducted a double-blind, placebo-controlled study in untreated elderly NDMM patients randomised to receive MPT ($n=182$) or MP ($n=175$) (Table 2). HRQoL was assessed as a secondary end point. Generalised estimating equations were applied, making full use of repeated quarterly measures (Table 1) and allowing for within-patient correlations over successive time points. Analyses were carried out using the observed values of QLQ-C30 scores, including baseline (pre-randomisation) QLQ-C30 scores as covariates.^{48,49}

In both treatment arms, HRQoL improved after treatment initiation. Little difference was detected between treatment arms, although significant differences were observed in favour of MP in Physical Functioning ($P=0.025$) and Social Functioning ($P=0.013$). There was a marked increase in the Constipation score among patients in the MPT arm ($P<0.001$), and a corresponding tendency to an increase (HRQoL worsening) in the Diarrhoea score in MP patients ($P=0.002$). Compliance with completing the QLQ-C30 questionnaire was 82% in the MPT arm and 90% in the MP arm at 3 months, and 50% and 62% at 12 months, respectively.

Critical review of NMSG HRQoL data. The main advantage of this HRQoL data is the study's double-blind, placebo-controlled design, which eliminates reporting bias. Multiple imputation of missing data values showed no evidence of bias in the comparison of treatments. However, the HRQoL data were substantially weakened in this double-blind trial by poor compliance in questionnaire completion as the study progressed.

HRQoL data from the NMSG (Hjorth *et al.*). Thalidomide- and bortezomib-naive patients with melphalan-refractory myeloma were randomly assigned to low-dose TD ($n=67$) vs VD ($n=64$) in an open phase III randomised multi-centre trial conducted by Hjorth *et al.* HRQoL was assessed as a secondary end point, measured by the QLQ-C30 questionnaire (Tables 1 and 2). The questionnaire was completed by 96% of patients still alive at 6 weeks, 90% at 12 weeks and by 76% patients at 6 months. No HRQoL improvement over time was observed for either treatment group. No between-group differences were noted, except that

Table 2. Key results from thalidomide clinical trials reporting HRQoL data

Trial/references	Trial design	Key HRQoL results
Thalidomide HOVON49 ⁴⁷	Prospective HRQoL study to assess the impact of thalidomide on HRQoL. Standard MP ($n = 168$) vs MPT followed by Thal maintenance ($n = 165$) in elderly (>65 years) NDMM patients	Physical Function and Constipation (significant) showed an improvement during induction in favour of the MP arm. During Thal maintenance, paraesthesia was significantly higher in the MPT arm, and a trend towards improved Pain, Insomnia, Appetite Loss and the QLQ-MY24 item sick scores was observed. The GHS/Global HRQoL scale showed a significant time trend towards more favourable mean values during protocol treatment without differences between MP and MPT. For the QLQ-C30 subscales, Emotional Functioning and Future Perspectives, a difference in favour of the MPT arm from the start of treatment was observed, with no significant 'time \times arm' interaction, indicating a persistent better patient perspective with MPT treatment. Overall, HRQoL outcomes improved equally in both arms, apart from markedly increased Constipation in the MPT arm.
Waage <i>et al.</i> ; Guldbrandsen <i>et al.</i> ^{48,49}	Phase III randomised, double-blind, placebo-controlled study in untreated elderly NDMM patients to compare MPT ($n = 182$) vs MP ($n = 175$)	No differences were noted for Physical Functioning, Pain and GHS/QoL. Fatigue and Sleep Disturbances were more prevalent in the VD group. The Fatigue score for the VD group was worse at 12 weeks, with a score difference of 10 ($P = 0.04$). The difference from the time of randomisation in score for Sleep Disturbances in the VD group reached statistical significance at 12 weeks ($P < 0.01$).
Hjorth <i>et al.</i> ⁵⁰	Open phase III randomised multi-centre trial to compare TD ($n = 67$) vs VD ($n = 64$) in melphalan-refractory myeloma patients	

Abbreviations: GHS, Global Health Status; HRQoL, health-related quality of life; MP, melphalan and prednisone; MPT, melphalan, prednisone and thalidomide; NDMM, newly diagnosed multiple myeloma; Thal, thalidomide; TD, thalidomide and dexamethasone; VD, bortezomib and dexamethasone.

the Fatigue score was worse at 12 weeks ($P = 0.04$) in the VD group. A higher Sleep Disturbances score was also noted in the VD group at 6 ($P = 0.06$) and 12 weeks ($P < 0.01$), potentially related to neurotoxicity.

*Critical review of NMSG HRQoL data (Hjorth *et al.*).* The trial was prematurely closed because of low accrual and was therefore weakened by the low number of recruited patients. However, it remains pertinent, as no other randomised data comparing thalidomide with bortezomib are available.

All domains of the QLQ-C30 were collected and reported for all time points, and all patients were included in the analysis in accordance with intention to treat principle. However, there was relatively little description of how HRQoL data were analysed, and it was not possible to infer how missing data were treated.

Clinically meaningful thresholds were used in interpreting HRQoL results, but a single MID of ≥ 10 points was used across all domains.⁵⁷ Compliance in questionnaire completion was high. The HRQoL questionnaires were first completed before randomisation, thus eliminating bias in HRQoL reporting at that time point. As in the HOVON49 trial, the quality of the data is diminished by the open-label design.

Studies reporting on HRQoL in MM patients treated with bortezomib

Three publications reported HRQoL results for bortezomib treatment in the relapsed/refractory multiple myeloma (RRMM) setting,^{58–60} covering the APEX phase III and SUMMIT phase II trials.^{17,61} The phase III VISTA trial reported HRQoL results for bortezomib treatment in elderly NDMM patients.⁶² These publications were also discussed in earlier conference proceedings,^{63–67} which also reported on the UPFRONT clinical trial in NDMM patients.^{68–71} Key HRQoL results from these trials are summarised in Table 3.

HRQoL data from the APEX phase III trial. The APEX study was a randomised, open-label trial comparing bortezomib ($n = 296$) with high-dose dexamethasone ($n = 302$) in patients with relapsed MM, evaluated with the QLQ-C30 and the FACT-Ntx questionnaires (Table 1). Assessment of HRQoL was included as a pre-specified exploratory efficacy objective.

The APEX trial was stopped early owing to a 29% vs 52% progression rate in favour of bortezomib. In the bortezomib arm, 9% completed all protocol-specified treatment, while in the dexamethasone arm, 5% completed treatment.⁶⁰ HRQoL assessments were discontinued when patients stopped protocol treatment, leading to a high amount of missing data.

At baseline, mean QLQ-C30 scores were significantly better for bortezomib vs dexamethasone in Emotional Functioning, Fatigue, Sleep and Diarrhoea. Baseline FACT-Ntx scores were comparable across groups. HRQoL scores during the 42 weeks of the trial were analysed using generalised estimating equation analysis of covariance.⁷² QLQ-C30 analysis found significantly better HRQoL in the bortezomib group vs dexamethasone, although a declining trend in mean GHS score was observed in both arms (Table 3). The component scores for Physical, Role, Cognitive and Emotional Functioning, and the symptom scores for Dyspnoea and Sleep were significantly better for the bortezomib group. For the overall FACT-Ntx score, statistically significant differences favouring the bortezomib arm were reported, when missing data due to patient death were imputed as worst possible score (zero), but this difference became nonsignificant when treated as withdrawals/missing data.

Critical review of APEX HRQoL data. The APEX trial was an open-label randomised study. This landmark trial reported an HRQoL benefit of bortezomib over dexamethasone. There were some differences in the two treatment arms at baseline: the bortezomib group reported better functioning and fewer symptoms than the dexamethasone group. These differences may be owing to chance, as the treatment arm was randomly assigned, HRQoL assessments were made before therapy started and the two groups were balanced for clinical characteristics. In responding patients, HRQoL changes from baseline were similar for most domains, except for better sleep and more neurotoxicity in bortezomib-treated patients, and less nausea and anorexia in the high-dose dexamethasone-treated group, consistent with clinical experience. The analysis may have been more robust if the HRQoL changes over time had been taken into account. The number of patients excluded from the HRQoL analyses was disclosed ($n = 45$, because a baseline assessment was missing or because only a baseline assessment was completed). The study also acknowledged

Table 3. Key results from bortezomib clinical trials reporting HRQoL data

Trial/references	Trial design	Key HRQoL results
Bortezomib APEX ^{60,65}	Prospective, open-label, randomised, phase III trial of bortezomib (<i>n</i> = 296) vs Dex (<i>n</i> = 302) in patients with relapsed MM	Bortezomib was associated with significantly better HRQoL compared with Dex, consistent with better clinical outcomes, although a declining trend in mean GHS score was observed in both arms Patients receiving bortezomib demonstrated significantly better mean GHS over the study compared with patients receiving Dex, as well as significantly better Physical Health, Role, Cognitive and Emotional Functioning scores, lower Dyspnoea and Sleep symptom scores, and better FACT-Ntx questionnaire scores
SUMMIT ^{58,59,63,64}	Open-label, multi-centre, phase II trial of bortezomib in patients with refractory MM (<i>n</i> = 202, 193 evaluated)	Change in HRQoL scores showed statistically significant differences between response groups with HRQoL improvement in patients with CR or PR, mostly stable scores in patients with minor response or no change Fifteen HRQoL parameters were significant in predicting mortality when univariate logistic regression was used. When using multivariate regression with stepwise selection to predict survival, only Fatigue and physical subscores were significant predictors of survival
VISTA ^{62,66,67}	Randomised, open-label, multi-centre, phase III study to compare VMP (<i>n</i> = 344) vs MP (<i>n</i> = 338) in NDMM patients	Clinically meaningful, transient HRQoL deterioration was observed with VMP vs MP during early treatment (up to cycle 4), followed by HRQoL improvements in VMP-treated patients for all domains, relative to baseline/MP from cycle 5 onwards Overall, mean scores improved in both arms by end-of-treatment vs baseline. Among responding patients, mean scores improved from time of response to end-of-treatment assessment. Multivariate analysis showed a significant impact of response duration or CR on GHS/Global HRQoL, Pain and Appetite Loss. Analyses by bortezomib dose intensity indicated better HRQoL in patients receiving lower dose intensity
UPFRONT ^{68–71}	Randomised, open-label, multi-centre phase IIIb trial to compare bortezomib with (i) Dex, (ii) Thal and Dex, or (iii) VMP, followed by bortezomib maintenance in NDMM patients (100 patients per arm)	A trend to decreased HRQoL score was observed in all treatment groups during induction, followed by an increase or stabilisation by the end of treatment There were no differences between treatment arms during induction. Moderate improvements were seen during maintenance, except for Nausea/Vomiting and Diarrhoea

Abbreviations: CR, complete response; Dex, dexamethasone; GHS, Global Health Status; HRQoL, health-related quality of life; MM, multiple myeloma; MP, melphalan and prednisone; NDMM, newly diagnosed multiple myeloma; PR, partial response; Thal, thalidomide; VMP, melphalan, prednisone and bortezomib.

that missing data increased with time, owing to AEs that led to discontinuation, PD, premature termination of the dexamethasone arm of the study and death. Although the number of patients completing treatment cycles was reported, the study did not specify compliance rates.

The study assessed the change in HRQoL scores over time by comparing the change in scores according to clinical response between baseline and the best clinical response (best endpoint).⁶⁰ The choice of 'best endpoint' for HRQoL analysis rather than a combination of data points throughout the trial could be considered selective. This approach may introduce bias, as AEs or symptoms occurring at other time points would be discounted.

Furthermore, the statistical analysis plan for the evaluation of the HRQoL data pre-specified four analytical methods, including two that included and two that excluded multiple data imputations, each combined with deaths assigned a zero value or deaths treated as withdrawals. Details on GHS/QoL data were primarily reported based on the method with no multiple data imputation, and deaths assigned a zero value, reporting an HRQoL benefit for bortezomib. No statistically significant differences in GHS/QoL between bortezomib and dexamethasone were observed with multiple data imputations and deaths treated as withdrawals.⁶⁰ The way in which deaths were treated had an important impact on data interpretation. As acknowledged by the authors, this approach likely biased the results in favour of bortezomib, given the significant survival benefit seen in the trial.¹⁶

HRQoL data from the SUMMIT phase II trial. SUMMIT was an open-label, single-arm trial of bortezomib in 202 patients with RRMM after at least two

previous treatments. The study design was described previously.⁶¹ During the first two cycles, all patients received bortezomib, and dexamethasone could be added for patients with stable disease or PD after four cycles. The study reported rates of 96–97% HRQoL completion at baseline and 76–77% at close-out.

The trial analysed HRQoL results (Tables 1 and 3) reported by clinical response in order to predict the prognostic value of HRQoL for survival in MM.^{58,59}

For the total patient population with available clinical response information (*n* = 151) and available HRQoL data (*n* = 144), there was a positive change between baseline and best end point. Changes in HRQoL scores showed statistically significant differences between response groups with HRQoL improvement in responding patients (with complete response (CR) or partial response), overall stable scores in patients with minor response or stable disease, and overall decreased scores in PD patients.⁵⁸

The value of baseline HRQoL data in predicting mortality during treatment was analysed through univariate and multivariate logistic regression and by partial least squares regression. Fifteen baseline HRQoL parameters were significant in predicting mortality during treatment when univariate logistic regression was used, but only the QLQ-C30 fatigue and physical subscores were significant predictors of survival in a subsequent multivariate regression (Table 3).⁵⁹

Critical review of SUMMIT HRQoL data. As a treatment comparison arm was not included, the impact of treatment on HRQoL changes is difficult to

interpret. The study did not present the mean HRQoL change from baseline by cycle or domain. The specified analytical end point assessed the change in HRQoL scores over time by comparing the change in scores according to clinical response between baseline and the best clinical response to treatment.⁵⁸ This choice of 'best end point' for HRQoL analysis rather than pre-determined data points throughout the trial is selective and was not justified in the reporting.

HRQoL data from the VISTA phase III trial. The phase III VISTA trial was a randomised, open-label, multi-centre study performed to assess VMP ($n = 344$) vs MP ($n = 338$) treatment on overall survival and other clinical benefits in elderly patients with previously untreated MM and patients not eligible for SCT.^{62,66,67} HRQoL was an exploratory end point of the VISTA trial. Patients were followed up over 54 weeks (nine 6-week cycles) and post treatment. The QLQ-C30 questionnaire was completed at screening, on day 1 of each treatment cycle and every 8 weeks until progression during follow-up (Table 1). A sustained HRQoL improvement was defined as a change in score of at least 5 points for at least two consecutive cycles after best response, as described by Dubois *et al.*⁵⁸ During early treatment cycles, an overall deterioration in HRQoL was observed in VMP patients, both vs baseline and MP-treated patients. At the cycle 4 assessment, mean differences between the VMP and MP arms were deemed clinically meaningful (≥ 5 points) and statistically significant ($P < 0.05$) for all domain scores, except for Cognitive Functioning, Nausea/Vomiting, and Dyspnoea. From cycle 5 onwards, a general increase in all HRQoL domain scores was reported in VMP-treated patients vs baseline and MP-treated patients (Table 3).⁶²

The VISTA trial also evaluated the impact of clinical response on HRQoL across both treatment arms. Mean scores improved overall in responding patients from time of response to end-of-treatment assessment, especially in patients achieving CR. Multivariate analysis showed a significant impact of duration of response/CR on improving GHS/QoL, Pain, Appetite Loss and Diarrhoea scores ($P \leq 0.03$ for all). Sustained HRQoL improvements of ≥ 5 points were seen following achievement of response.⁶²

The impact of bortezomib dose on HRQoL was also evaluated in VMP-treated patients at the time of, and one and two cycles before, both best response and the end-of-treatment visit. Patients receiving a lower dose intensity of bortezomib (< 5.6 mg/m²/cycle) for at least two cycles before achieving overall response or before their end-of-treatment visit generally reported better HRQoL vs the higher dose intensity group.⁶²

Critical review of VISTA HRQoL data. The open-label VISTA study is to date the largest randomised, multi-centre study in previously untreated, transplant-ineligible MM patients reporting on HRQoL, with HRQoL collected as an exploratory end point. HRQoL analyses were restricted to data collected from baseline to the end-of-treatment visit, owing to low questionnaire completion rates post-treatment. The results, however, demonstrated that HRQoL is not compromised in the long term with VMP vs MP.

While the number of patients for whom HRQoL data were available and comparable across the study, no between-group comparisons of patient and disease characteristics or compliance rates were reported. This is important, given that the authors did no imputation for missing data. Patients may drop out of the study because of poor response or side effects, which may result in overestimations of HRQoL scores.

Additional analysis investigated differences in HRQoL within treatment arms using multivariate linear regression. The improvement in some HRQoL domains identified after 'best response' may be influenced by the fact that HRQoL scores for VMP patients were lowest at the point of CR, due to toxicity issues.

Exploratory, *post hoc* analyses by dose intensity provide some evidence regarding the impact on HRQoL of bortezomib dose intensity. The VISTA study authors acknowledged that the study was not designed to compare HRQoL during the periods of twice-weekly and once-weekly bortezomib dosing, so the evidence on higher HRQoL with lower dosages of bortezomib is preliminary. Further studies will be required to confirm a statistical relationship between bortezomib dose intensity and HRQoL.

HRQoL data from the UPFRONT phase IIIb trial. In an ongoing randomised, open-label, multi-centre clinical trial that compared the efficacy and safety of three bortezomib-based regimens in untreated, transplant-ineligible NDMM patients, Niesvizky *et al.* described HRQoL data as a primary objective from a total of 300 patients (100 patients per arm) who completed the QLQ-C30 questionnaire. The phase IIIb UPFRONT study compared the safety and efficacy of VD with thalidomide and dexamethasone (VTD), and with melphalan and prednisone (VMP), followed by bortezomib maintenance therapy (Table 3).⁶⁸⁻⁷¹

Scores improved in all three treatment arms, except for Physical Functioning, Role Functioning and GHS/QoL, which worsened in the VTD arm.⁶⁸⁻⁷¹ The observed data, linear mixed model estimates and sensitivity analyses all showed a common trend to a transient decrease in HRQoL during VD, VTD and VMP induction, followed by a subsequent trend to improvement/stabilisation in HRQoL during single-agent bortezomib maintenance.

A significant worsening (reduction) ($P < 0.05$) in mean GHS/QoL score at cycle 7 from baseline was reported in the VTD and VMP arms (linear mixed effect model). A sensitivity analysis used last observation carried forward for patients with missing data, showing a significant worsening from baseline at cycle 7 in all bortezomib-based treatment arms ($P < 0.05$). Symptom scores changed very little during induction with all bortezomib-based regimens, with moderate HRQoL improvements seen during maintenance (except for Nausea, Vomiting and Diarrhoea).

Niesvizky *et al.* concluded that the trend to declining HRQoL during induction may reflect the onset of treatment-associated toxicity. Subsequent HRQoL improvement may reflect the positive impact of achieving a response. The transient decline in HRQoL observed in this study is similar to the trend previously reported in the VISTA study.⁶⁶

Critical review of UPFRONT HRQoL data. The UPFRONT data were presented in abstract/poster format, which limits the scope for in-depth critical evaluation. The trial design was open-labelled, but given the fact that all patients received the investigational product, the potential for enhanced response in patients who know that they are receiving an investigational therapy can be excluded.

In terms of compliance, HRQoL assessments were available at baseline and at least one post-baseline time point for 80% (VD), 67% (VTD) and 80% (VMP) of patients. The information on compliance rates is fairly unspecific, as it does not provide sufficient information on overall compliance per treatment cycle, and in particular during the maintenance phase.

The UPFRONT study authors concluded that post-induction improvements/stabilisation in HRQoL may reflect the beneficial impact of achieving a response and the limited toxicity profile associated with weekly bortezomib maintenance. However, unhealthier patients may not have completed HRQoL questionnaires at later stages of treatment, for example, owing to neurotoxicity, leading to a potential bias in reporting.

For patients who died within the HRQoL evaluation period, missing HRQoL assessments were assigned a score of zero, representing the worst possible HRQoL score. While potentially exaggerating the HRQoL of treatments that reduce mortality, this aspect is unlikely to have been a major issue in this trial, given the similar survival rates observed.

Studies reporting on HRQoL in MM patients treated with lenalidomide

Two studies reporting HRQoL data in MM patients treated with lenalidomide were identified in the search, including comparisons of MPR-R vs MPR vs MP in NDMM patients above the age of 65 (MM-015 trial),⁷³⁻⁷⁶ and lenalidomide plus high-dose dexamethasone in RRMM patients (MM-018 trial).⁷⁷⁻⁷⁹ Key HRQoL results from these trials are summarised in Table 4.

HRQoL data from the MM-015 phase III trial. The MM-015 phase III trial was a multi-centre, randomised, double-blind, placebo-controlled, three-arm parallel-group study to determine the efficacy and safety of MPR-R ($n = 152$) vs MPR ($n = 153$) and MP ($n = 154$) in NDMM subjects who are 65 years of age or older (Table 4). This study investigated HRQoL as a secondary outcome at baseline and the beginning of every third cycle, and at time of progression (PD) or discontinuation for reasons other than progression (DC). HRQoL was assessed using the QLQ-C30 and QLQ-MY20 for the following pre-selected HRQoL domains: GHS/QoL, Physical Functioning, Fatigue and Pain (QLQ-C30), and Side Effects of Treatment and Disease Symptoms (QLQ-MY20) (Table 1).

In all three study arms, HRQoL improvement was observed from baseline to study end for all specified HRQoL domains. Statistically significant HRQoL improvements ($P < 0.05$) from baseline to cycle 10 (end of induction phase) were observed in each domain for MPR-treated patients, with the exception of Side Effects of Treatment (nonsignificant for all treatment arms). Improvements in HRQoL in patients aged 65-75 years were comparable, or slightly greater, to those in all patients aged ≥ 65 years.

Comparably favourable HRQoL results could not be replicated for the subset of patients aged ≥ 75 years, owing to the limited number of patients.

Table 4. Key results from lenalidomide clinical trials reporting HRQoL data

Trial/references	Trial design	Key HRQoL results
Lenalidomide MM-015 ⁷³⁻⁷⁶	Phase III, multi-centre, randomised, double-blind, placebo-controlled, three-arm parallel-group study (MPR-R <i>n</i> = 152, MPR <i>n</i> = 153 or MP <i>n</i> = 154) in NDMM patients ≥65 years	Clinically meaningful improvements in HRQoL were more frequently observed in patients receiving MPR-R than those receiving MP The differences in HRQoL were most marked in terms of Physical Functioning, and overall results were consistent in patients aged 65 to 75 years A higher percentage of MPR-R patients achieved the MID in six pre-selected HRQoL domains than those receiving MP. Improved or stabilising longitudinal HRQoL trends were observed for most other HRQoL domains
MM-018 ⁷⁷⁻⁷⁹	Phase III, multi-centre, single-arm, open-label, expanded-access study in RRMM subjects (<i>N</i> = 587) treated with Len + high-dose Dex in 4-week cycles. Subjects followed until PD or DC	In the Spanish cohort of the study, GHS/Global QoL, Fatigue, Emotional Functioning, Physical Functioning, Role Functioning, Social Functioning, Cognitive Functioning and Pain improved >5 points Preservation of HRQoL correlated with response to treatment in terms of Role Functioning, Emotional Functioning, Social Functioning and Pain scores

Abbreviations: DC, discontinuation; Dex, dexamethasone; HRQoL, health-related quality of life; GHS, Global Health Status; Len, lenalidomide; MID, minimal important difference; MM, multiple myeloma; MP, melphalan and prednisone; MPR(-R), melphalan, prednisone and lenalidomide (and maintenance lenalidomide); NDMM, newly diagnosed multiple myeloma; PD, progressive disease.

Changes in HRQoL from cycle 10 to cycle 16 were not statistically significant ($P > 0.05$) in any HRQoL domain, suggesting that the majority of HRQoL improvement occurred during the induction phase and stabilised during maintenance.⁷⁶ With regards to changes from baseline at cycle 16, there were statistically significant changes in the MPR-R group for all domains (again with the exception of Side Effects of Treatment), while statistically significant changes were not consistently observed for MPR and MP patients.⁷⁶

A mixed model repeated measures analysis for the QLQ-C30 and QLQ-MY20 scores indicated a significant difference in changes in Physical Functioning scores across treatment groups: scores improved significantly ($P < 0.05$) from baseline in MPR-R and MPR patients but worsened in MP patients.⁷⁶

Comparisons of HRQoL scores using trial- and domain-specific MIDs were also conducted and time points at which mean changes from baseline for each arm exceeded the MID were presented.^{75,76} In this study, the standard error of measurement was used to establish MID.^{80,81} The MPR-R group mean change from baseline exceeded the MID more frequently than was observed in the other two treatment groups. MID HRQoL improvements were observed as early as cycle 4 for Pain. MIDs were applied to investigate the percentage of patients who had a change from baseline exceeding the MID at cycle 10 and 16 for MPR-R and MP treatment groups. A higher number of patients in the MPR-R group exceeded the MID for all domains.

An additional analysis presented results from a mixed-effects multiple regression model that estimated which clinical parameters were associated with statistically significant and clinically meaningful improvements in HRQoL in the MPR-R and MP treatment arms. More patients achieved very good partial response or better (\geq VGPR) when receiving continuous MPR-R treatment than those receiving MP. \geq VGPR was shown to improve GHS/QoL in a clinically meaningful and statistically significant way, suggesting that clinical responses with MPR-R treatment were not related to HRQoL impairments due to treatment-related toxicity. PD was also shown to negatively impact GHS/QoL (-8.34 ; $P < 0.001$), with MPR-R significantly reducing the risk of PD vs MP. Continuous MPR-R may therefore delay PD and help improve and maintain HRQoL.⁷⁴

Critical review of MM-015 HRQoL data. The percentage of patients completing questionnaires until cycle 16 was consistently above a threshold of 76%, with compliance rates above 65% at PD/DC. The percentage of compliant subjects was not significantly different between treatment arms at any of the visits, except at cycle 7 for the QLQ-MY20 questionnaire ($P = 0.036$).⁷⁶

Mean HRQoL domain scores were presented for each treatment at each measurement time point, along with longitudinal differences between treatment arms.

Significant emphasis in the presentation of the findings, as reflected by most reported comparative analyses between treatment arms, was placed on 'responder' analyses (percentage of patients achieving a clinically

meaningful HRQoL response in each arm), using MID as the definition of clinically meaningful response.

Patients randomised to MPR-R had worse HRQoL scores compared with the other arms, but the difference was statistically significant only for Physical Functioning ($P = 0.014$). Sensitivity analyses and inclusion of 'time \times arm' interactions accounted for baseline differences. Mixed models were used to estimate the treatment effect on HRQoL over time, adjusted and unadjusted for baseline HRQoL scores.

The study used repeated measures mixed-effects modelling to account for missing variables and described the extent of missing data overall and by treatment arm. There were no statistically significant differences in demographics and disease-related characteristics between the three treatment arms at baseline, cycle 10 or cycle 16, suggesting that there was no significant difference between treatment arms in patients who dropped out or were non-compliant.

HRQoL data from the MM-018 phase III trial. In the MM-018 phase III single-arm, open-label study, lenalidomide plus high-dose dexamethasone was administered to 587 RRMM patients in the United Kingdom, Spain and Ireland, to assess the safety of this regimen and its impact on HRQoL (Table 4). Secondary outcome HRQoL assessments were conducted at baseline and after 24 weeks of treatment using the QLQ-C30 and QLQ-MY20 questionnaires (Table 1).

QLQ-C30 revealed no significant median change (>5 points MID) from baseline in 14 of 15 domains for patients completing questionnaires at baseline and 24 weeks. Median Fatigue increased in the United Kingdom/Ireland population (score 11.1). QLQ-MY20 revealed no significant median change from baseline of all scores except an improvement in Future Perspective in Spanish patients (median 11.1), for patients completing questionnaires at baseline and 24 weeks.⁷⁹ Alegre *et al.*^{77,78} reported further HRQoL data from 63 patients enrolled in the Spanish cohort. At week 24, 42 patients were available for HRQoL assessment. In addition to the reported improvement in Future Perspective, a nonsignificant impairment in the Physical Functioning domain of the QLQ-C30 functional scores was also observed (<5 points MID). The majority of patients who experienced HRQoL changes according to QLQ-C30 and QLQ-MY20 scores had clinically meaningful improvements in HRQoL, regardless of response (20/42 patients achieved either a CR or VGPR during treatment). Despite comedication with high-dose dexamethasone, pre-existing neuropathy in $>50\%$ patients, prior MM treatment, and late disease course, patients were able to maintain median QoL scores over 24 weeks.⁷⁷⁻⁷⁹

Critical review of MM-018 HRQoL data. The open-label, single-arm study design did not allow for reliable inference on the extent to which

treatment truly impacts upon HRQoL. The data were collected at two end points (baseline and week 24) and are therefore likely to be incomplete for some patients (for example, 42/63 patients completed HRQoL at 24 weeks in the Spanish subset). Unlike the APEX trial for bortezomib, no information on missing data or withdrawals was reported. This may constitute a major weakness especially among an RRMM population where missing data are more likely to introduce potential bias.

CONCLUSIONS

To date, there has been a relatively small body of HRQoL data published on novel MM treatments. Available HRQoL data do not allow for comparisons of HRQoL impact across MM treatments, owing to differences in patient population, lack of comparative trials, differences in study designs and in methodology applied for the specific HRQoL analysis.

Patient groups differ between trials, for example, in terms of age distribution and pathology (NDMM or RRMM) being particularly diverse, which may impact on HRQoL. Baseline HRQoL values were significantly different between treatment groups in the thalidomide HOVON49 and the bortezomib APEX trials. In cases where baseline HRQoL results are statistically significantly different between treatment arms, it is important to conduct sensitivity analyses in order to control for baseline differences in HRQoL, as was done in the MM-015 trial.

Study design may also impact on interpretation of HRQoL outcomes. Few studies were double-blind RCTs, such as the lenalidomide MM-015 and thalidomide Waage *et al.* trials, making inference for all other trials more difficult. Unblinded studies (for example, the thalidomide HOVON49 and all retrieved bortezomib trials) may increase the potential for an enhanced response in patients who are aware they are receiving an investigational treatment. Even though a number of studies were open-labelled (for example, thalidomide HOVON49 and Hjorth *et al.* trials, and bortezomib APEX, VISTA and UPFRONT trials), their value in terms

of inference is superior to single-arm trials (for example, bortezomib SUMMIT and lenalidomide MM-018 trials). Furthermore, some studies have only been reported as conference proceedings (UPFRONT and extended analyses on clinical parameters affecting HRQoL in MM-015).

Differences in interpreting clinically meaningful MID thresholds for MM were observed across retrieved studies. However, there is no unanimous agreement on what constitutes clinically meaningful MID changes per HRQoL domain, allowing for several different definitions of MIDs as well as methods of analysis.

All retrieved studies explored HRQoL as pre-specified secondary end points, except for the bortezomib UPFRONT study, in which measurement of HRQoL changes is described as a primary objective. RCTs measuring HRQoL as a primary outcome have been shown to display higher concordance on pre-specified quality measures.⁸² Consistent use of the well-validated QLQ-C30 and QLQ-MY20 questionnaires facilitates comparisons between treatments. Of note, for individual treatment combinations containing novel therapies, patterns in longitudinal HRQoL trends are generally consistent across the majority of HRQoL domains analysed. There are similarities between studies, regarding timing of HRQoL assessments (Table 1), with common time points at the beginning of treatment cycles. There have, however, been inconsistent approaches to the analysis of HRQoL data across studies in MM, and not all studies have reported the observed HRQoL at each time point for all arms in the study, which would constitute the most straightforward and transparent way to present findings.

Differences in dealing with and reporting missing data were observed across the retrieved studies. The thalidomide trials retrieved in this analysis incorporated all data points at which HRQoL was assessed and presented the observed results at all time points. APEX and UPFRONT trials applied missing data imputations but did not present these findings in detail. In contrast, missing data imputations were not carried out in the

Table 5. Guidance in collecting and analysing HRQoL in MM patients treated with novel agents based on the current analysis

Instruments

- Internationally validated questionnaires, to be used in their entirety
- Questionnaires measuring the impact of treatment toxicity
- Prospective design

Study design

- Intention to treat principle
- Preferably randomised double-blind trial
- If study design is a randomised, open-label trial, baseline questionnaire to be completed before randomisation

Assessment time points:

- At baseline and at different pre-determined treatment time points
- At the end of treatment, ideally including a differentiation between disease progression and discontinuation
- Regular HRQoL assessments following end of treatment, if possible

Reporting and analysis

Compliance reporting:

- Compliance rates regarding questionnaire completion at each assessment time point and per study arm
- Statistical between-group comparisons at individual measurement time points, assessing data interpretability independent of absolute compliance levels
- Between-group comparisons of individual patient categories (study drop-outs vs non-compliant vs compliant patients) in terms of patient and disease characteristics and inclusion of treatment interaction terms

Types of HRQoL assessment:

- Longitudinal and cross-sectional analysis reporting of HRQoL
- Reporting of individual HRQoL scores at each time point and for each study arm
- Illustration of mean HRQoL changes from baseline over time via repeated measures analysis
- Illustration of both statistical significance and clinical meaningfulness (e.g., with MIDs)
 - Determination of MID should be based on a combination of statistical reasoning and clinical judgement,⁴³ including both trial- and domain-specific analyses
 - Linear mixed model per treatment arm across all measurement time points
 - Sensitivity analysis controlling for baseline HRQoL and baseline differences in key patient and disease characteristics

Abbreviations: HRQoL, health-related quality of life; MID, minimal important difference; MM, multiple myeloma.

thalidomide Hjorth *et al.* trial, in the VISTA and SUMMIT trials for bortezomib, or the MM-015 and MM-018 trials for lenalidomide. However, results from MM-015 were strengthened through mixed model repeated measures analyses. Mixed model repeated measures analyses across time points and paired analyses of data both between treatment arms at individual time points and longitudinally within treatment arms across two time points are methods of assuring that observed changes are not attributable to the changing nature of the sample across time.

Differences in compliance rates between treatment arms, an important pre-requisite for cross-sectional HRQoL data comparisons, were reported in the thalidomide Waage *et al.* and in the lenalidomide MM-015 studies, but were not discussed in the other trials.

Finally, in HRQoL analyses such as APEX, which assume a zero HRQoL score for patients who have died, results may have favoured bortezomib owing to the significant survival benefit established in the study.

As some treatment options have prolonged survival in MM patients, and owing to the impact of treatment-related toxicity on HRQoL, HRQoL data have become increasingly relevant key performance indicators. In the absence of differences in treatment efficacy, the choice of initial treatment should be based on HRQoL, among other patient-related factors. Quality-adjusted survival analyses that integrate HRQoL considerations may be important, particularly in treatments that do not show significant survival advantages.⁸³ Guidelines for best practice in collecting and analysing HRQoL in MM would ensure that future data are more useful in informing clinical decisions, whereby more consistent reporting of HRQoL data will improve the understanding of the HRQoL impact of different MM treatments. For those assessing HRQoL in MM studies, our review provides guidance on good practices and standardisation for HRQoL data collection, analysis and reporting (Table 5). The proposed incorporation of HRQoL as a clinically relevant end point in MM drug registration dossiers and in RCTs stresses the need for validated instruments and specific questionnaires, for instance to measure the impact of toxicities such as peripheral neuropathy.^{37,84} Future HRQoL investigations in MM patients would gain value if head-to-head comparative studies were carried out.

CONFLICT OF INTEREST

Sonneveld P: Advisory Boards and Research support: Janssen, Celgene, Onyx, Millennium; Verelst S: no disclosures; Lewis P: Celgene employee, stock options; Gray-Schopfer V: Consultancy fees as external contractual partner to Global Market Access Solutions (GMAS); Hutchings A: Consultancy fees to GMAS from Celgene; Nixon A: Consultancy fees from Celgene to Oxford Outcomes for MM-015 HRQoL analysis and publications; Petrucci MT Honoraria: Celgene, Janssen-Cilag.

REFERENCES

- Johnsen AT, Tholstrup D, Petersen MA, Pedersen L, Groenvold M. Health related quality of life in a nationally representative sample of haematological patients. *Eur J Haematol* 2009; **83**: 139–148.
- Siris ES, Sherman WH, Baquiran DC, Schlatterer JP, Osserman EF, Canfield RE. Effects of dichloromethylene diphosphonate on skeletal mobilization of calcium in multiple myeloma. *N Engl J Med* 1980; **302**: 310–315.
- Terpos E, Berenson J, Cook RJ, Lipton A, Coleman RE. Prognostic variables for survival and skeletal complications in patients with multiple myeloma osteolytic bone disease. *Leukemia* 2010; **24**: 1043–1049.
- Palumbo A, Anderson K. Multiple myeloma. *N Engl J Med* 2011; **364**: 1046–1060.
- Dimopoulos MA, Roussou M, Gkotzamanidou M, Nikitas N, Psimenou E, Mparmparoussi D *et al.* The role of novel agents on the reversibility of renal impairment in newly diagnosed symptomatic patients with multiple myeloma. *Leukemia* 2013; **27**: 423–429.
- Pratt G, Goodyear O, Moss P. Immunodeficiency and immunotherapy in multiple myeloma. *Br J Haematol* 2007; **138**: 563–579.
- Van Valckenborgh E, Schoupe E, Movahedi K, De Bruyne E, Menu E, De Baetselier P *et al.* Multiple myeloma induces the immunosuppressive

capacity of distinct myeloid-derived suppressor cell subpopulations in the bone marrow. *Leukemia* 2012; **26**: 2424–2428.

- Kumar S, Rajkumar SV, Dispenzieri A, Lacy MQ, Hayman SR, Buadi FK *et al.* Improving survival in multiple myeloma: impact of novel therapies. Proceedings of the 49th Annual Meeting of the American Society of Hematology (ASH), Atlanta, GA; USA. *Blood* 2007; **110**: Abstract 3954.
- Laubach JP, Mahindra A, Mitsiades CS, Schlossman RL, Munshi NC, Ghobrial IM *et al.* The use of novel agents in the treatment of relapsed and refractory multiple myeloma. *Leukemia* 2009; **23**: 2222–2232.
- Brenner H, Gondos A, Pulte D. Recent major improvement in long-term survival of younger patients with multiple myeloma. *Blood* 2008; **111**: 2521–2526.
- Kristinsson SY, Landgren O, Dickman PW, Derolf AR, Bjorkholm M. Patterns of survival in multiple myeloma: a population-based study of patients diagnosed in Sweden from 1973 to 2003. *J Clin Oncol* 2007; **25**: 1993–1999.
- Verelst SGR, Blommestein HM, Karim-Kos HE, Huijgens PC, Sonneveld P. Trends in incidence and survival of multiple myeloma in the Netherlands in the last two decades. Results from a national population based study. Proceedings of the 53rd Annual Meeting of the American Society of Hematology (ASH), San Diego, CA; USA. *Blood* 2011; **118**: Abstract 5071.
- Verelst S, Karim-Kos H, Blommestein H, Sonneveld P. Are we making progress? Survival in plasma cell malignancies in the era of novel treatments a population based study of 17.790 patients in The Netherlands. Proceedings of the 17th Congress of the European Hematology Association (EHA), Amsterdam, The Netherlands. *Haematologica* 2012; **97**: 242.
- Dimopoulos M, Spencer A, Attal M, Prince HM, Harousseau JL, Dmoszynska A *et al.* Lenalidomide plus dexamethasone for relapsed or refractory multiple myeloma. *N Engl J Med* 2007; **357**: 2123–2132.
- Dimopoulos MA, Chen C, Spencer A, Niesvizky R, Attal M, Stadtmauer EA *et al.* Long-term follow-up on overall survival from the MM-009 and MM-010 phase III trials of lenalidomide plus dexamethasone in patients with relapsed or refractory multiple myeloma. *Leukemia* 2009; **23**: 2147–2152.
- Richardson PG, Sonneveld P, Schuster M, Irwin D, Stadtmauer E, Facon T *et al.* Extended follow-up of a phase 3 trial in relapsed multiple myeloma: final time-to-event results of the APEX trial. *Blood* 2007; **110**: 3557–3560.
- Richardson PG, Sonneveld P, Schuster MW, Irwin D, Stadtmauer EA, Facon T *et al.* Bortezomib or high-dose dexamethasone for relapsed multiple myeloma. *N Engl J Med* 2005; **352**: 2487–2498.
- Weber DM, Chen C, Niesvizky R, Wang M, Belch A, Stadtmauer EA *et al.* Lenalidomide plus dexamethasone for relapsed multiple myeloma in North America. *N Engl J Med* 2007; **357**: 2133–2142.
- Larocca A, Palumbo A. Evolving paradigms in the treatment of newly diagnosed multiple myeloma. *J Natl Compr Canc Netw* 2011; **9**: 1186–1196.
- Palumbo A, Attal M, Roussel M. Shifts in the therapeutic paradigm for patients newly diagnosed with multiple myeloma: maintenance therapy and overall survival. *Clin Cancer Res* 2011; **17**: 1253–1263.
- Rajkumar SV, Jacobus S, Callander NS, Fonseca R, Vesole DH, Williams ME *et al.* Lenalidomide plus high-dose dexamethasone versus lenalidomide plus low-dose dexamethasone as initial therapy for newly diagnosed multiple myeloma: an open-label randomised controlled trial. *Lancet Oncol* 2010; **11**: 29–37.
- Sonneveld P, Schmidt-Wolf I, van der Holt B, el Jarari L, Bertsch U, Salwender H *et al.* HOVON-65/GMMG-HD4 randomized phase III trial comparing bortezomib, doxorubicin, dexamethasone (PAD) vs VAD followed by high-dose melphalan (HDM) and maintenance with bortezomib or thalidomide in patients with newly diagnosed multiple myeloma (MM). Proceedings of the 52nd Annual Meeting of the American Society of Hematology (ASH), Orlando, FL; USA. *Blood* 2010; **116**: Abstract 40.
- Palumbo A, Rajkumar SV. Treatment of newly diagnosed myeloma. *Leukemia* 2009; **23**: 449–456.
- Attal M, Lauwers-Cances V, Marit G, Caillot D, Moreau P, Facon T *et al.* Lenalidomide maintenance after stem-cell transplantation for multiple myeloma. *N Engl J Med* 2012; **366**: 1782–1791.
- McCarthy PL, Owzar K, Hofmeister CC, Hurd DD, Hassoun H, Richardson PG *et al.* Lenalidomide after stem-cell transplantation for multiple myeloma. *N Engl J Med* 2012; **366**: 1770–1781.
- Hahn-Ast C, von Lilienfeld-Toal M, van Heteren P, Mückter S, Brossart P, Glasmacher A. Improved progression-free and overall survival with thalidomide maintenance therapy after autologous stem cell transplantation in multiple myeloma: a meta-analysis of five randomized trials. Proceedings of the 16th Congress of the European Hematology Association (EHA), London, United Kingdom. *Haematologica* 2011; **96**: 367 Abstract 0884.
- Dispenzieri A, Jacobus S, Vesole DH, Callander N, Fonseca R, Greipp PR. Primary therapy with single agent bortezomib as induction, maintenance and re-induction in patients with high-risk myeloma: results of the ECOG E2A02 trial. *Leukemia* 2010; **24**: 1406–1411.

- 28 Bang SM, Kyle RA, Rajkumar SV, Kumar S. Treatment patterns and outcomes in elderly patients with multiple myeloma. *Leukemia* 2013; **27**: 971–974.
- 29 Facon T, Mary JY, Hulin C, Benboubker L, Attal M, Pegourie B et al. Melphalan and prednisone plus thalidomide versus melphalan and prednisone alone or reduced-intensity autologous stem cell transplantation in elderly patients with multiple myeloma (IFM 99-06): a randomised trial. *Lancet* 2007; **370**: 1209–1218.
- 30 Hulin C, Facon T, Rodon P, Pegourie B, Benboubker L, Doyen C et al. Efficacy of melphalan and prednisone plus thalidomide in patients older than 75 years with newly diagnosed multiple myeloma: IFM 01/01 trial. *J Clin Oncol* 2009; **27**: 3664–3670.
- 31 Kapoor P, Rajkumar SV, Dispenzieri A, Gertz MA, Lacy MQ, Dingli D et al. Melphalan and prednisone versus melphalan, prednisone and thalidomide for elderly and/or transplant ineligible patients with multiple myeloma: a meta-analysis. *Leukemia* 2011; **25**: 689–696.
- 32 Mateos MV, Richardson PG, Schlag R, Khuageva NK, Dimopoulos MA, Shpilberg O et al. Bortezomib plus melphalan and prednisone compared with melphalan and prednisone in previously untreated multiple myeloma: updated follow-up and impact of subsequent therapy in the phase III VISTA trial. *J Clin Oncol* 2010; **28**: 2259–2266.
- 33 San Miguel JF, Schlag R, Khuageva NK, Dimopoulos MA, Shpilberg O, Kropff M et al. Bortezomib plus melphalan and prednisone for initial treatment of multiple myeloma. *N Engl J Med* 2008; **359**: 906–917.
- 34 Palumbo A, Hajek R, Delforge M, Kropff M, Petrucci MT, Catalano J et al. Continuous lenalidomide treatment for newly diagnosed multiple myeloma. *N Engl J Med* 2012; **366**: 1759–1769.
- 35 Palumbo A, Adam Z, Kropff M, Foa R, Catalano J, Gisslinger H et al. A phase 3 study evaluating the efficacy and safety of lenalidomide (len) combined with melphalan and prednisone followed by continuous lenalidomide maintenance (MPR-R) in patients (pts) ≥65 years (yrs) with newly diagnosed multiple myeloma (NDMM): updated results for pts aged 65-75 yrs enrolled in MM-015. Proceedings of the 53rd Annual Meeting of the American Society of Hematology (ASH), San Diego, CA; USA. *Blood* 2011; **118**: Abstract 475.
- 36 Palumbo A, Mateos MV, Brinchen S, San Miguel JF. Practical management of adverse events in multiple myeloma: can therapy be attenuated in older patients? *Blood Rev* 2011; **25**: 181–191.
- 37 Richardson PG, Delforge M, Beksac M, Wen P, Jongen JL, Sezer O et al. Management of treatment-emergent peripheral neuropathy in multiple myeloma. *Leukemia* 2012; **26**: 595–608.
- 38 Jordan K, Ishak JK, Lewis P, Proskorovsky I, Payne KA, Lordan N et al. Determinants of global QOL and physical and social functionality in multiple myeloma. Proceedings of the 52nd Annual Meeting of the American Society of Hematology (ASH), Orlando, FL; USA. *Blood* 2010; **116**: Abstract 934.
- 39 Schumacher A, Kessler L, Buchner T, Wewers D, van de Loo J. Quality of life in adult patients with acute myeloid leukemia receiving intensive and prolonged chemotherapy—a longitudinal study. *Leukemia* 1998; **12**: 586–592.
- 40 Efficace F, Baccarani M, Breccia M, Cottone F, Alimena G, Deliliers GL et al. Chronic fatigue is the most important factor limiting health-related quality of life of chronic myeloid leukemia patients treated with imatinib. *Leukemia* 2013; **27**: 1511–1519.
- 41 Aliibhai SM, Leach M, Kowgier ME, Tomlinson GA, Brandwein JM, Minden MD. Fatigue in older adults with acute myeloid leukemia: predictors and associations with quality of life and functional status. *Leukemia* 2007; **21**: 845–848.
- 42 Koenigsman M, Koehler M, Franke A, Frommer J. Acute leukaemia in adults: researching the patient's perspective. *Leukemia* 2006; **20**: 206–207.
- 43 European Medicines Agency. Reflection paper on the regulatory guidance for the use of health-related quality of life (HRQL) measures in the evaluation of medicinal products: EMEA/CHMP/EWP/139391/2004 2004.
- 44 US Department of Health and Human Services Food and Drug Administration Guidance for Industry. Patient-Reported Outcome Measures: Use in Medical Product Development to Support Labeling Claims 2009.
- 45 Bottomley A, Efficace F, Fayers PM. Standards are needed for quality of life clinical trials. *BMJ* 2002; **324**: 1156.
- 46 Lee CW, Chi KN. The standard of reporting of health-related quality of life in clinical cancer trials. *J Clin Epidemiol* 2000; **53**: 451–458.
- 47 Vereist SG, Termorshuizen F, Uyl-de Groot CA, Schaafsma MR, Ammerlaan AH, Wittebol S et al. Effect of thalidomide with melphalan and prednisone on health-related quality of life (HRQoL) in elderly patients with newly diagnosed multiple myeloma: a prospective analysis in a randomized trial. *Ann Hematol* 2011; **90**: 1427–1439.
- 48 Waage A, Gimsing P, Fayers P, Abildgaard N, Ahlberg L, Bjorkstrand B et al. Melphalan and prednisone plus thalidomide or placebo in elderly patients with multiple myeloma. *Blood* 2010; **116**: 1405–1412.
- 49 Gulbrandsen N, Waage A, Gimsing P, Turesson I, Juliusson G, Abildgaard N et al. A randomised placebo controlled study with melphalan/prednisone vs melphalan/prednisone/thalidomide: quality of life and toxicity. Proceedings of the 13th Congress of the European Hematology Association (EHA), Copenhagen, Denmark. *Haematologica* 2008; **93**: 84 (Abstract 0209).
- 50 Hjorth M, Hjertner O, Knudsen LM, Gulbrandsen N, Holmberg E, Pedersen PT et al. Thalidomide and dexamethasone vs. bortezomib and dexamethasone for melphalan refractory myeloma: a randomized study. *Eur J Haematol* 2012; **88**: 485–496.
- 51 Wijermans P, Schaafsma M, Termorshuizen F, Ammerlaan R, Wittebol S, Sinnige H et al. Phase III study of the value of thalidomide added to melphalan plus prednisone in elderly patients with newly diagnosed multiple myeloma: the HOVON 49 study. *J Clin Oncol* 2010; **28**: 3160–3166.
- 52 Fayers PM, Palumbo A, Hulin C, Waage A, Wijermans P, Beksac M et al. Thalidomide for previously untreated elderly patients with multiple myeloma: meta-analysis of 1685 individual patient data from 6 randomized clinical trials. *Blood* 2011; **118**: 1239–1247.
- 53 Palumbo A, Brinchen S, Caravita T, Merla E, Capparella V, Callea V et al. Oral melphalan and prednisone chemotherapy plus thalidomide compared with melphalan and prednisone alone in elderly patients with multiple myeloma: randomised controlled trial. *Lancet* 2006; **367**: 825–831.
- 54 Palumbo A, Brinchen S, Liberati AM, Caravita T, Falcone A, Callea V et al. Oral melphalan, prednisone, and thalidomide in elderly patients with multiple myeloma: updated results of a randomized controlled trial. *Blood* 2008; **112**: 3107–3114.
- 55 Mohty B, El-Cheikh J, Yakoub-Agha I, Moreau P, Housseau JL, Mohty M. Peripheral neuropathy and new treatments for multiple myeloma: background and practical recommendations. *Haematologica* 2010; **95**: 311–319.
- 56 Kvam AK, Fayers P, Wisloff F. What changes in health-related quality of life matter to multiple myeloma patients? A prospective study. *Eur J Haematol* 2010; **84**: 345–353.
- 57 Osoba D, Rodrigues G, Myles J, Zee B, Pater J. Interpreting the significance of changes in health-related quality-of-life scores. *J Clin Oncol* 1998; **16**: 139–144.
- 58 Dubois D, Dhawan R, van de Velde H, Esseltine D, Gupta S, Viala M et al. Descriptive and prognostic value of patient-reported outcomes: the bortezomib experience in relapsed and refractory multiple myeloma. *J Clin Oncol* 2006; **24**: 976–982.
- 59 Viala M, Bhakar AL, de la Loge C, van de Velde H, Esseltine D, Chang M et al. Patient-reported outcomes helped predict survival in multiple myeloma using partial least squares analysis. *J Clin Epidemiol* 2007; **60**: 670–679.
- 60 Lee SJ, Richardson PG, Sonneveld P, Schuster MW, Irwin D, San Miguel JF et al. Bortezomib is associated with better health-related quality of life than high-dose dexamethasone in patients with relapsed multiple myeloma: results from the APEX study. *Br J Haematol* 2008; **143**: 511–519.
- 61 Richardson PG, Barlogie B, Berenson J, Singhal S, Jagannath S, Irwin D et al. A phase 2 study of bortezomib in relapsed, refractory myeloma. *N Engl J Med* 2003; **348**: 2609–2617.
- 62 Delforge M, Dhawan R, Robinson Jr D, Meunier J, Regnault A, Esseltine DL et al. Health-related quality of life in elderly, newly diagnosed multiple myeloma patients treated with VMP versus MP: Results from the VISTA trial. *Eur J Haematol* 2012; **89**: 16–27.
- 63 Dubois D, Ravinder D, van de Velde HA, Esseltine D, Gupta S, Viala M et al. Use of patient-reported outcomes (PROs) to assess prognosis, toxicity, and response: The bortezomib experience in relapsed and refractory multiple myeloma (MM). Proceedings of the 46th Annual Meeting of the American Society of Hematology (ASH), San Diego, CA; USA. *Blood* 2004; **104**: Abstract 3468.
- 64 Lee S, Richardson PG, Barlogie B, Berenson JR, Singhal S, Jagannath S et al. For the SUMMIT Investigators. Quality-of-life (QOL) and clinical benefit assessment in patients with relapsed and refractory multiple myeloma (MM) treated with bortezomib. 2003 ASCO Annual Meeting Proceedings. *J Clin Oncol* 2003; **22**: Abstract 2339.
- 65 Lee SJ, Richardson PG, Sonneveld P, Schuster M, Irwin D, Massaro J et al. Health-related quality of life (HRQL) associated with bortezomib compared with high-dose dexamethasone in relapsed multiple myeloma (MM): results from APEX study. 2005 ASCO Annual Meeting Proceedings. *J Clin Oncol* 2005; **23**: Abstract 6535.
- 66 Dhawan R, Meunier J, Regnault A, Rosa K, Robinson D, Cakana A et al. Impact of complete response on health-related quality of life in newly diagnosed multiple myeloma patients: results from the VISTA trial. XII International Myeloma Workshop (IMW), Washington DC; USA. *Clin Lymphoma Myeloma* 2009; **9**: S58 (Abstract 352).
- 67 Dhawan R, Robinson D, Meunier J, Regnault A, Rosa K, Cakana AZ et al. Sustained health-related quality of life (HRQoL) improvement in newly diagnosed multiple myeloma patients treated with bortezomib/melphalan/prednisone versus melphalan/prednisone: results from the VISTA trial. Proceedings of the 51st Annual Meeting of the American Society of Hematology (ASH), New Orleans, LA; USA. *Blood* 2009; **114**: Abstract 1881.

- 68 Niesvizky R, Flinn IW, Rifkin R, Gabrail N, Charu V, Clowney B *et al*. Patient-reported quality of life (QoL) in elderly, newly diagnosed multiple myeloma (MM) patients receiving bortezomib-based combinations: results from all randomized patients in the community-based, phase 3b UPFRONT study. Proceedings of the 53rd Annual Meeting of the American Society of Hematology (ASH), San Diego, CA; USA. *Blood* 2011; **118**: Abstract 1864.
- 69 Niesvizky R, Flinn I, Rifkin R, Gabrail N, Charu V, Clowney B *et al*. Patient-reported quality of life (QOL) in previously untreated, elderly multiple myeloma (MM) patients treated with bortezomib-based regimens: results from the phase 3b UPFRONT study. Proceedings of the 16th Congress of the European Hematology Association (EHA), London, United Kingdom. *Haematologica* 2011; **96**: 123 (Abstract 0298).
- 70 Niesvizky R, Flinn IW, Rifkin RM, Gabrail NY, Veena Charu V, Clowney B *et al*. Phase 3b UPFRONT study: safety and efficacy of weekly bortezomib maintenance therapy after bortezomib-based induction regimens in elderly, newly diagnosed multiple myeloma patients. Proceedings of the 52nd Annual Meeting of the American Society of Hematology (ASH), Orlando, FL; USA. *Blood* 2010; **116**: Abstract 619.
- 71 Niesvizky R, Reeves J, Flinn I, Gabrail N, Rifkin R, Charu V *et al*. Phase 3b UPFRONT study: interim results from a community-based prospective randomized trial evaluating three bortezomib-based regimens in elderly, newly diagnosed multiple myeloma patients. Proceedings of the 15th Congress of the European Hematology Association (EHA), Barcelona, Spain. *Haematologica* 2010; **95**: 144 (Abstract 0358).
- 72 Zeger SL, Liang KY, Albert PS. Models for longitudinal data: a generalized estimating equation approach. *Biometrics* 1988; **44**: 1049–1060.
- 73 Dimopoulos M, Delforge M, Hajek R, Kropff M, Petrucci MT, Lewis P *et al*. Melphalan, prednisone and lenalidomide followed by lenalidomide maintenance improves health-related quality of life (HRQoL), with newly diagnosed multiple myeloma (NDMM) patients ≥ 65 years benefiting from delays in disease progression. Proceedings of the 53rd Annual Meeting of the American Society of Hematology (ASH), San Diego, CA; USA. *Blood* 2011; **118**: Abstract 3157.
- 74 Dimopoulos M, Palumbo A, Hajek R, Kropff M, Petrucci MT, Lewis P *et al*. Melphalan, prednisone and lenalidomide followed by lenalidomide maintenance displays treatment characteristics favourable to global quality of life in newly diagnosed multiple myeloma (NDMM) patients ≥ 65 years. Proceedings of the 53rd Annual Meeting of the American Society of Hematology (ASH), San Diego, CA; USA. *Blood* 2011; **118**: Abstract 3988.
- 75 Dimopoulos MA, Delforge M, Hajek R, Kropff M, Petrucci MT, Lewis P *et al*. Lenalidomide plus melphalan and prednisone followed by lenalidomide maintenance provides favourable efficacy and health-related quality-of-life in newly diagnosed multiple myeloma patients ≥ 65 years. Proceedings of the 16th Congress of the European Hematology Association (EHA), London, United Kingdom. *Haematologica* 2011; **96**: 365 (Abstract 0880).
- 76 Dimopoulos MA, Delforge M, Hajek R, Kropff M, Petrucci MT, Lewis P *et al*. Lenalidomide, melphalan, and prednisone, followed by lenalidomide maintenance, improves health-related quality of life in newly diagnosed multiple myeloma patients aged 65 years or older: results of a randomized phase III trial. *Haematologica* 2013; **98**: 784–788.
- 77 Alegre A, Oriol-Rocafiguera A, Garcia-Larana J, Mateos MV, Sureda A, Chamorro CM *et al*. Efficacy, safety and quality-of-life associated with lenalidomide plus dexamethasone for the treatment of relapsed or refractory multiple myeloma: the Spanish experience. *Leuk Lymphoma* 2012; **53**: 1714–1721.
- 78 Alegre A, Oriol-Rocafiguera A, Garcia-Larana J, Mateos MV, Sureda A, Martinez Chamorro C *et al*. Safety, efficacy, and quality-of-life study of lenalidomide plus dexamethasone in previously treated patients with multiple myeloma: the Spanish experience. Proceedings of the 52nd Annual Meeting of the American Society of Hematology (ASH), Orlando, FL; USA. *Blood* 2010; **116**: Abstract 3045.
- 79 Yong K, Alegre Amor A, Browne P, Cavenagh J, Dodds T, Greil R *et al*. A multi-center, single-arm, open-label safety and quality of life study of lenalidomide plus dexamethasone in previously treated patients with multiple myeloma. Proceedings of the 15th Congress of the European Hematology Association (EHA), Barcelona, Spain. *Haematologica* 2010; **95**: 392 (Abstract 0944).
- 80 Wyrwich KW, Nienaber NA, Tierney WM, Wolinsky FD. Linking clinical relevance and statistical significance in evaluating intra-individual changes in health-related quality of life. *Med Care* 1999; **37**: 469–478.
- 81 Wyrwich KW, Tierney WM, Wolinsky FD. Further evidence supporting an SEM-based criterion for identifying meaningful intra-individual changes in health-related quality of life. *J Clin Epidemiol* 1999; **52**: 861–873.
- 82 Brundage M, Bass B, Davidson J, Queenan J, Bezjak A, Ringash J *et al*. Patterns of reporting health-related quality of life outcomes in randomized clinical trials: implications for clinicians and quality of life researchers. *Qual Life Res* 2011; **20**: 653–664.
- 83 Levy V, Porcher R, Leblond V, Femand JP, Cazin B, Maloisel F *et al*. Evaluating treatment strategies in advanced Waldenström macroglobulinemia: use of quality-adjusted survival analysis. *Leukemia* 2001; **15**: 1466–1470.
- 84 Anderson KC, Kyle RA, Rajkumar SV, Stewart AK, Weber D, Richardson P. Clinically relevant end points and new drug approvals for myeloma. *Leukemia* 2008; **22**: 231–239.
- 85 Aaronson NK, Ahmedzai S, Bergman B, Bullinger M, Cull A, Duez NJ *et al*. The European Organization for Research and Treatment of Cancer QLQ-C30: a quality-of-life instrument for use in international clinical trials in oncology. *J Natl Cancer Inst* 1993; **85**: 365–376.
- 86 Osoba D, Zee B, Pater J, Warr D, Kaizer L, Latreille J. Psychometric properties and responsiveness of the EORTC quality of Life Questionnaire (QLQ-C30) in patients with breast, ovarian and lung cancer. *Qual Life Res* 1994; **3**: 353–364.
- 87 Gulbrandsen N, Wisloff F, Brinch L, Carlson K, Dahl IM, Gimsing P *et al*. Health-related quality of life in multiple myeloma patients receiving high-dose chemotherapy with autologous blood stem-cell support. *Med Oncol* 2001; **18**: 65–77.
- 88 Uyl-de Groot CA, Buijt I, Gloudemans IJ, Ossenkoppele GJ, Berg HP, Huijgens PC. Health related quality of life in patients with multiple myeloma undergoing a double transplantation. *Eur J Haematol* 2005; **74**: 136–143.
- 89 Wisloff F, Eika S, Hippe E, Hjorth M, Holmberg E, Kaasa S *et al*. Measurement of health-related quality of life in multiple myeloma. Nordic Myeloma Study Group. *Br J Haematol* 1996; **92**: 604–613.
- 90 Wisloff F, Hjorth M. Health-related quality of life assessed before and during chemotherapy predicts for survival in multiple myeloma. Nordic Myeloma Study Group. *Br J Haematol* 1997; **97**: 29–37.
- 91 Schunemann HJ, Guyatt GH. Commentary—goodbye M(C)ID! Hello MID, where do you come from? *Health Serv Res* 2005; **40**: 593–597.
- 92 Cocks K, Cohen D, Wisloff F, Sezer O, Lee S, Hippe E *et al*. An international field study of the reliability and validity of a disease-specific questionnaire module (the QLQ-MY20) in assessing the quality of life of patients with multiple myeloma. *Eur J Cancer* 2007; **43**: 1670–1678.
- 93 Stead ML, Brown JM, Velikova G, Kaasa S, Wisloff F, Child JA *et al*. Development of an EORTC questionnaire module to be used in health-related quality-of-life assessment for patients with multiple myeloma. European Organization for Research and Treatment of Cancer Study Group on Quality of Life. *Br J Haematol* 1999; **104**: 605–611.
- 94 Lavoie Smith EM, Barton DL, Qin R, Steen PD, Aaronson NK, Loprinzi CL. Assessing patient-reported peripheral neuropathy: the reliability and validity of the European Organization for Research and Treatment of Cancer QLQ-CIPN20 Questionnaire. *Qual Life Res* 2013, e-pub ahead of print 2 April 2013 doi:10.1007/s11136-013-0379-8.
- 95 Wagner LI, Robinson Jr. D, Weiss M, Katz M, Greipp P, Fonseca R *et al*. Content development for the functional assessment of cancer therapy-multiple myeloma (FACT-MM): use of qualitative and quantitative methods for scale construction. *J Pain Symptom Manage* 2012; **43**: 1094–1104.
- 96 Weiss M, Jacobus S, Wagner LI, Cella D, Katz MS, Rajkumar V *et al*. Development of the Functional Assessment of Cancer Therapy-Multiple Myeloma (FACT-MM) scale and validation in the Eastern Cooperative Oncology Group Trial E1A05. Proceedings of the 53rd Annual Meeting of the American Society of Hematology (ASH), San Diego, CA; USA. *Blood* 2011; **118**: Abstract 4184.
- 97 Cella DF, Tulsky DS, Gray G, Sarafian B, Linn E, Bonomi A *et al*. The Functional Assessment of Cancer Therapy scale: development and validation of the general measure. *J Clin Oncol* 1993; **11**: 570–579.
- 98 Cella D, Eton DT, Lai JS, Peterman AH, Merkel DE. Combining anchor and distribution-based methods to derive minimal clinically important differences on the Functional Assessment of Cancer Therapy (FACT) anemia and fatigue scales. *J Pain Symptom Manage* 2002; **24**: 547–561.
- 99 Calhoun EA, Welshman EE, Chang CH, Lurain JR, Fishman DA, Hunt TL *et al*. Psychometric evaluation of the Functional Assessment of Cancer Therapy/Gynecologic Oncology Group-Neurotoxicity (Fact/GOG-Ntx) questionnaire for patients receiving systemic chemotherapy. *Int J Gynecol Cancer* 2003; **13**: 741–748.
- 100 Kvam AK, Fayers PM, Wisloff F. Responsiveness and minimal important score differences in quality-of-life questionnaires: a comparison of the EORTC QLQ-C30 cancer-specific questionnaire to the generic utility questionnaires EQ-5D and 15D in patients with multiple myeloma. *Eur J Haematol* 2011; **87**: 330–337.



This work is licensed under a Creative Commons Attribution 3.0 Unported License. To view a copy of this license, visit <http://creativecommons.org/licenses/by/3.0/>