



---

Nosocomial and Community-Acquired Staphylococcus aureus Bacteremias from 1980 to 1993:  
Impact of Intravascular Devices and Methicillin Resistance

Author(s): James P. Steinberg, Catherine C. Clark, Betsy O. Hackman

Source: *Clinical Infectious Diseases*, Vol. 23, No. 2 (Aug., 1996), pp. 255-259

Published by: The University of Chicago Press

Stable URL: <http://www.jstor.org/stable/4459597>

Accessed: 03/06/2009 06:20

---

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at <http://www.jstor.org/page/info/about/policies/terms.jsp>. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Please contact the publisher regarding any further use of this work. Publisher contact information may be obtained at <http://www.jstor.org/action/showPublisher?publisherCode=ucpress>.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

JSTOR is a not-for-profit organization founded in 1995 to build trusted digital archives for scholarship. We work with the scholarly community to preserve their work and the materials they rely upon, and to build a common research platform that promotes the discovery and use of these resources. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).



The University of Chicago Press is collaborating with JSTOR to digitize, preserve and extend access to *Clinical Infectious Diseases*.

# Nosocomial and Community-Acquired *Staphylococcus aureus* Bacteremias from 1980 to 1993: Impact of Intravascular Devices and Methicillin Resistance

James P. Steinberg, Catherine C. Clark,  
and Betsy O. Hackman

From the Division of Infectious Diseases, Department of Medicine,  
Emory University School of Medicine, and the Infection Control  
Department, Crawford Long Hospital of Emory University,  
Atlanta, Georgia

The rate of nosocomial bacteremia due to *Staphylococcus aureus* has increased over the past decade, but trends in community-acquired *S. aureus* bacteremia are less certain. This hospital-based observational study compares nosocomial and community-acquired *S. aureus* bacteremias during 1980–1983 and 1990–1993. The rate of nosocomial *S. aureus* bacteremia increased from 0.75 to 2.80 cases per 1,000 discharges, while the rate of community-acquired *S. aureus* bacteremia increased from 0.84 to 2.43 cases per 1,000 discharges. The number of nosocomial device-related bacteremias increased eightfold; 56% of *S. aureus* bacteremias were associated with devices during 1990–1993. Intravascular devices were associated with no community-acquired *S. aureus* bacteremias during 1980–1983 but with 22% during 1990–1993. Methicillin-resistant *S. aureus* (MRSA) seldom caused bacteremia during 1980–1983. From 1990 to 1993, MRSA caused 32% and 18.5% of nosocomial and community-acquired *S. aureus* bacteremias, respectively. The rates of both community-acquired and nosocomial *S. aureus* bacteremias have increased significantly since 1980. In addition to their role in nosocomial infections, MRSA and intravascular device-related *S. aureus* bacteremias are emerging problems in the nonhospital setting.

In the 1980s, gram-positive organisms, including *Staphylococcus aureus*, reemerged as the leading causes of nosocomial bacteremia [1, 2]. From 1980 to 1989, the National Nosocomial Infections Surveillance System (NNIS) of the Centers for Disease Control and Prevention (CDC) reported increases in the rates of primary bacteremia due to *S. aureus* of 122% to 283% [1]. During the late 1980s, *S. aureus* caused 16% of nosocomial bacteremias reported to the NNIS, second only to coagulase-negative staphylococci [3]. An increase in the rate of *S. aureus* bacteremias also has been observed abroad. In Denmark, the annual incidence rate of *S. aureus* bacteremia increased from 2.7 cases per 100,000 population in 1960 to 19.2 cases per 100,000 population in 1990, with nosocomial bacteremias accounting for most of this increase [4].

---

See the editorial by Darouiche and Musher  
on pages 260–1.

---

A larger population of immunocompromised individuals and increased use of intravascular devices are among the factors causing this resurgence of staphylococcal bacteremias. In addition, the emergence of methicillin-resistant *S. aureus* (MRSA) contributes to the upward trend in *S. aureus* infections. Changes in community-acquired *S. aureus* bacteremia are less well char-

acterized. The objectives of this study are to describe trends in nosocomial and community-acquired *S. aureus* bacteremias since 1980 and to examine risk factors for *S. aureus* bacteremia including the impact of intravascular devices.

## Methods

Crawford Long Hospital of Emory University is a 500-bed acute care hospital in downtown Atlanta. Community-based physicians and full-time university faculty utilize the facility, which serves as both a community hospital and a referral center.

All bloodstream infections were identified by the infection control department as part of routine surveillance. For this study, we compared bloodstream infections due to *S. aureus* during 1980–1983 with those during 1990–1993. Clinical information, including the presence of underlying diseases and source of bacteremia, was obtained by chart review. Nosocomial bacteremia was defined by a bloodstream isolate obtained  $\geq 48$  hours after hospital admission, while community-acquired bacteremia was defined by a bloodstream isolate obtained within 48 hours of hospital admission. However, blood isolates obtained within 48 hours of readmission were classified as nosocomial if the infection was determined by the infection control practitioner to be incubating at the time of a recent (<30 days) hospital discharge.

Blood isolates of *S. aureus* were considered to be contaminants if only one blood culture set yielded the organisms, if the clinicians judged the organisms to be contaminants, and if antibiotic therapy directed against the organisms was not administered.

Intravascular device-related *S. aureus* bacteremia was defined as bacteremia unrelated to infection at an extravascular


---

Received 9 November 1995; revised 20 February 1996.

Reprints or correspondence: Dr. James P. Steinberg, Division of Infectious Diseases, 20 Linden Avenue, Suite 101, Atlanta, Georgia 30308.

Clinical Infectious Diseases 1996;23:255–9

© 1996 by The University of Chicago. All rights reserved.  
1058-4838/96/2302-0007\$02.00



**Figure 1.** Rates of nosocomial and community-acquired bacteremias due to methicillin-resistant *Staphylococcus aureus* (MRSA), methicillin-susceptible *S. aureus* (MSSA), and other bacteria (excluding mycobacteria) during two study periods.

site with semiquantitative culture of a catheter tip yielding  $\geq 15$  colonies or with infection at a local catheter site that was due to the same organism. When the catheter was not removed or the tip was not cultured, the infection was considered to be device-related if the infection control practitioner, the hospital epidemiologist, or the involved clinicians determined a device to be the likely source.

During the 1980–1983 study period, the hospital microbiology laboratory used the BACTEC 460 radiometric system (Becton Dickinson, Sparks, MD); for each set of blood cultures, a 7.5- to 10-mL blood sample was inoculated into aerobic, anaerobic, and hypertonic media. During the 1990–1993 study period, the BACTEC 660 nonradiometric system (Becton Dickinson) was used; a blood sample of 20 mL was inoculated into aerobic and anaerobic vials.

During both study periods, the hospital's policy called for replacement of peripheral intravenous catheters after 72 hours. There was no similar standard governing the duration of the placement of central venous catheters. From 1980–1983, there was limited use of central venous catheters outside of the intensive care units, and implanted ports and tunneled cuffed catheters were either not commercially available or not used.

The Mantel-Haenszel  $\chi^2$  test was used to determine statistical significance.

## Results

The rate of bacteremia doubled during the second 4-year period, with the rate of nosocomial bacteremia increasing from 4.59 to 9.44 cases per 1,000 discharges and the rate of community-acquired bacteremia increasing from 6.2 to 13.46 per 1,000 discharges (figure 1). The rates of nosocomial and community-acquired *S. aureus* bacteremias also increased from 0.75 to 2.80 cases per 1,000 discharges and from 0.84 to 2.43 cases per 1,000 discharges, respectively.

Table 1 shows the numbers of *S. aureus* and other bacteremias for the study periods. The proportion of nosocomial bacteremia caused by *S. aureus* increased from 16.3% to 29.6% ( $P < .001$ ), and the proportion of community-acquired bacteremia caused by *S. aureus* increased from 13.5% to 18.1% ( $P < .05$ ). The annual average number of blood cultures performed also doubled from 7,758 to 15,725, while the annual number of hospital discharges decreased slightly from 22,117 to 20,540 over the same periods. Two *S. aureus* isolates (1.4%) recovered from blood cultures were considered contaminants during the first period compared with 14 (3.1%) during the second period.

## Sources of Bacteremia and Underlying Diseases

The sources for all *S. aureus* bacteremias during the study periods are shown in table 2. The number of intravascular device-associated nosocomial *S. aureus* bacteremias increased eightfold from 16 (25% of total) during 1980–1983 to 128 (56% of total) during 1990–1993. This increase in the number of device-associated bacteremias accounts for 70% of the increase in the number of nosocomial *S. aureus* bloodstream infections. Fifty-two percent of the device-associated bacteremias from the 1990–1993 period were confirmed by cultures of catheter tips or specimens from device sites that yielded *S. aureus*.

No single source accounted for the increase in the number of community-acquired *S. aureus* bacteremias (table 2). Intravascular devices, not documented as a source of community-acquired bacteremia in the 1980–1983 period, were associated with 43 community-acquired bacteremias (22%) in the 1990–1993 period. These infections resulted primarily from long-term indwelling catheters (see next section). Skin and soft-tissue infections caused 44 community-acquired bacteremias (22%) in the 1990–1993 period. Of these 44 infections, 16 (nine due to MRSA) were from infected decubitus ulcers in nursing home patients, and 11 were from lower extremity infections in patients with diabetes or vascular insufficiency. Bacteremias related to hemodialysis shunts and fistulas were common in both study periods.

Two patients with community-acquired bacteremia in the 1980–1983 period and four patients in the 1990–1993 period were intravenous drug users. Comparing 1990–1993 with 1980–1983, nursing home residence (16% vs. 4%, respectively), AIDS (10% vs. 0, respectively), and sickle cell anemia (8% vs. 0, respectively) were significantly ( $P < .05$ ) more common, whereas diabetes (27% vs. 22%, respectively), renal failure (28% vs. 35%, respectively), and cancer (9% vs. 16%, respectively) were not. The mean age of patients with community-acquired bacteremia was 55 years in the 1980–1983 period and 57.6 years in the 1990–1993 period. Although nursing home residence was more common in the 1990–1993 period, a greater percentage of patients were 65 years of age or older in the earlier period (45% vs. 39%, respectively). Sixty-five percent of patients with community-acquired *S. aureus* bacter-

**Table 1.** Numbers of nosocomial and community-acquired bacteremias during 1980–1983 and 1990–1993.

Type of bacteremia, study period	No. of bacteremias due to indicated pathogen				Total
	<i>Staphylococcus aureus</i> (total)	MRSA	Other gram-positive organisms	Gram-negative organisms	
Nosocomial					
1980–1983	16.5	0.25	25.5	65	101.5
1990–1993	57.5	18	84	67	194
Community-acquired					
1980–1983	18.5	0.25	33	88	137
1990–1993	50	9	93	142.5	276.5

NOTE. Data represent average numbers of bacteremias per year during the 4-year study period. MRSA = methicillin-resistant *S. aureus*.

emia during the second study period were hospitalized within 1 year before the bacteremia, and 52% were hospitalized within 90 days of the bacteremia.

**Types of Intravascular Devices**

The types of intravascular devices associated with *S. aureus* bloodstream infections are listed in table 3. From 1990 to 1993, central venous catheters and peripheral intravascular catheters were associated with 31% and 18%, respectively, of the nosocomial device-related infections. These bacteremias occurred in patients with a wide variety of underlying medical and surgical

problems. Implanted ports were the source of 25% of the nosocomial bacteremias; 18 (56%) of the port-related infections occurred in patients with sickle cell disease, and 12 (38%) occurred in oncology patients. The 18 bacteremias in patients with sickle cell disease occurred in 12 patients. Six patients each had two bacteremias associated with implanted ports; the same device was implicated as the source in three of these six patients.

Sources of community-acquired device-related bacteremias during 1990–1993 included implanted ports (49%), hemodialysis catheters (33%), tunneled cuffed catheters (9%), and peripheral intravenous catheters (5%). The 43 bacteremias occurred in 39 patients. Underlying diseases in these patients included chronic renal failure (36%), AIDS (23%), sickle cell anemia (21%), and cancer (8%).

**Table 2.** Sources of *Staphylococcus aureus* bacteremia during the two study periods.

Source	No. of bacteremias due to MRSA/ total no. due to <i>S. aureus</i>	
	1980–1983	1990–1993
Nosocomial		
Intravascular devices	1/16	30/128*
Unknown†	0/23	12/54
Surgical wound infection	0/10	15/19
Skin/decubitus ulcers	0/3	6/9
Respiratory tract	0/5	6/11
Other	0/8	4/9
Total	1/65	73/230
Community-acquired		
Intravascular devices	0/0	8/43*
Unknown†	0/27	7/52
Dialysis shunt or fistula	1/18	5/28
Skin/decubitus ulcers	0/7	13/44‡
Bone/joint	0/4	0/11
Intravenous drug use	0/2	0/4
Respiratory tract	0/10*	0/4
Other	0/6	4/14
Total	1/74	37/200

\* Significant difference, *P* < .01.

† Includes cases where multiple possible sources were identified.

‡ Significant difference, *P* < .05.

**MRSA Bacteremia**

Only one nosocomial MRSA bacteremia occurred in the 1980–1983 period. During the 1990–1993 period, MRSA caused 73 nosocomial *S. aureus* bacteremias (32%). Sources of these bacteremias are listed in table 2.

Thirty-seven community-acquired MRSA bacteremias occurred from 1990 to 1993, whereas only one occurred a decade earlier. These 37 bacteremias occurred in 35 patients, all of whom had ongoing contact with health care settings. The patient had been hospitalized at our institution within the previous year in 24 (65%) of the cases; in 19 cases (51%), the patient had been hospitalized within the previous 90 days. Fourteen (40%) of the patients with community-acquired MRSA bacteremia resided in nursing homes. Forty-five percent (14 of 31) of *S. aureus* bacteremias occurring in nursing home residents were caused by MRSA.

Underlying diseases in patients with community-acquired MRSA bacteremia included diabetes (29%), chronic renal failure (23%), cancer (14%), AIDS (9%), and sickle cell anemia (3%). Nursing home residence, but not any of these underlying diseases, was associated with a significantly higher risk of

**Table 3.** Types of intravascular devices associated with *Staphylococcus aureus* bacteremia during two study periods.

Type of infection, study period	No. of <i>S. aureus</i> infections associated with indicated catheter					
	Peripheral intravenous	Central venous*	Tunneled†‡	Ports†§	Hemodialysis	Other
Nosocomial						
1980–1983	5	7	0	0	0	4
1990–1993	23	40	8	32	3	22
Community-acquired						
1980–1983	0	0	0	0	0	0
1990–1993	2	1	4	21	14	1

\* Includes single-, double-, and triple-lumen catheters.

† Tunneled cuffed catheters, primarily Hickman catheters.

‡ Not in use during 1980–1983.

§ Implanted ports.

|| Includes arterial catheters, Swan-Ganz catheters, umbilical catheters, and femoral vein catheters.

MRSA infection ( $P < .001$ ). The mean age of patients with community-acquired MRSA bacteremia was 67.7 years; 58% were 65 years of age or older.

## Discussion

This study shows a significant increase in the number of community-acquired as well as nosocomial *S. aureus* bacteremias at our institution from 1980–1983 to 1990–1993. The increase in the number of nosocomial *S. aureus* bacteremias is consistent with data generated by NNIS and other investigators [1, 2, 4, 5]. However, there are little comparable data on trends in community-acquired *S. aureus* bacteremia. Intravascular devices are the major source of nosocomial *S. aureus* bacteremia and are emerging as important sources of community-acquired bacteremia. Similarly, MRSA has emerged as a cause of bacteremia in patients outside the hospital setting.

Intravascular devices, implicated in no community-acquired bacteremias during the first study period, were associated with 22% of community-acquired *S. aureus* bacteremias from 1990 to 1993. We believe that this increase likely reflects the shifting of acute medical care to the outpatient setting and the increasing use of long-term devices in patients with chronic disorders, such as AIDS, cancer, and sickle cell disease. Although outpatient infusion therapy has been associated with a low rate of infectious complications [6], given current economic pressures, this low rate is likely to apply to an increasing population. In addition, the low reported rate of infections may not apply to all populations. Thus, a further increase in the total number of cases of community-acquired device-related infections may occur.

MRSA was responsible for 32% of nosocomial *S. aureus* bacteremias during the 1990–1993 period, a proportion consistent with NNIS data for hospitals of comparable size [7]. MRSA also caused 18.5% of community-acquired *S. aureus* bacteremias from 1990 to 1993. Most of these infections oc-

curred in patients who had been hospitalized recently or were nursing home residents, and all patients with community-acquired MRSA bacteremia had regular contact with health care settings.

Our definition of nosocomial infection was adapted from definitions of the CDC that were designed for the NNIS hospitals [8]; all other infections were termed *community-acquired*. For patients with chronic medical illnesses who may require periodic hospitalizations, home health care, and, possibly, outpatient parenteral therapy, the term *community-acquired* is misleading. These patients are at greater risk for serious infections, including those caused by antibiotic-resistant microorganisms such as MRSA, than are those without prior contact with health care facilities. For example, approximately two of three patients admitted to our hospital with community-acquired bacteremia due to *S. aureus* (methicillin-susceptible *S. aureus* or MRSA) had been hospitalized within the previous year.

The term *nosohusial* has been suggested to describe infections occurring in patients who receive care at home [6]. Whether this new term finds widespread acceptance, the distinction between patients requiring frequent contact with health care settings and those whose infections are truly community-acquired is valid. The increase in the number of patients receiving care at home emphasizes the need to develop methods for studying infections occurring in this setting. Although useful information can be obtained from hospital-based surveillance, an accurate determination of infection rates occurring in non-hospital settings should utilize the population at risk as the denominator. In addition, the use of multiple commercial clinical laboratories for performing culture and sensitivity testing further complicates attempts to determine infection rates in the nonhospital setting. Thus, further home care-based studies are warranted.

These data show a significant increase in the number of intravascular device-associated bacteremias since 1980. Because of the design of this study, we were unable to measure

device utilization or establish infection rates associated with various types of devices. However, the increased use of intravascular devices, particularly central venous catheters, has been implicated in the increased rates of septicemia from 1979 to 1987 that were recorded by the National Hospital Discharge Survey [9]. Fifty-six percent of nosocomial bacteremias in the second study period were device-related, a figure consistent with those of other published series [10]. Fifty-two percent of the nosocomial device-associated infections during the second study period were confirmed by cultures of catheter tips or specimens from device sites that yielded *S. aureus*. A more rigorous definition of device-associated infection, such as requiring cultures of catheter tips or specimens from device sites, could have led to a significant underestimation of the impact of devices. We believe that it is more likely that some of the bacteremias termed as *unknown* were device-related and that our calculation of device-associated infections is conservative.

A surprising finding in our study was the high number of nosocomial and community-acquired port-related bacteremias. Implanted ports have been associated with a low incidence of infection (range, 0.01–0.04 bloodstream infections per 100 device-days [11, 12]). Our data reflect, in part, a large population of patients with diseases such as sickle cell anemia and AIDS and a high utilization of implanted ports in these patients. In addition, suboptimal care of the devices and patient use of ports for injection of illicit drugs were suspected in a few instances. Although we were unable to determine rates of port-related bacteremia, our experience raises concern that the reported incidence of port-related infections may not apply to all settings. Notwithstanding the inferences about rates of port-related infection, these data demonstrate how the use of implanted ports can change the epidemiology of bloodstream infections.

The overall increase in the rates of bacteremia may be confounded by changes in blood culturing methodology and an increase in the number of blood cultures performed. During the 1990–1993 study period, the standard blood specimen for a set of blood cultures was 20 mL, twice that obtained in the 1980–1983 period. A recent controlled trial demonstrated that increasing the volume of blood inoculated into a culture vial from 2.7 mL to 8.7 mL increased the yield of gram-positive pathogens by 19% [13]. The number of blood cultures performed per year doubled from the first to the second study period. This pattern of increased utilization of blood cultures during the 1980s has been reported by other researchers and has been attributed to the presence of a sicker patient population [14].

The annual number of bacteremias also doubled in the second study period, resulting in a stable rate of blood culture positivity. This finding suggests that the methodological changes were not the major cause of the increased rates of bacteremia during 1990–1993. It is also unlikely that these factors accounted for the disproportionate increases in the numbers of *S. aureus* and device-related bacteremias.

In summary, these data show parallel increases in rates of nosocomial and community-acquired *S. aureus* bacteremias since 1980. Device-related infection, the leading source of nosocomial *S. aureus* bacteremia, is emerging as an important source of community-acquired bacteremia in our hospital. Similarly, MRSA bacteremia, once confined to the nosocomial setting, is an increasing problem in the nonhospital setting. The shifting of care, including parenteral therapy, to the outpatient setting likely accounts for some of these observations.

#### Acknowledgments

The authors thank Drs. Monica M. Farley, John E. McGowan, Jr., and Robert W. Pinner for their thoughtful review of the manuscript.

#### References

- Banerjee SN, Emori TG, Culver DH, et al. Secular trends in nosocomial primary bloodstream infections in the United States, 1980–1989. *Am J Med* 1991;91(suppl 3B):86S–9S.
- McGowan JE Jr. Changing etiology of nosocomial bacteremia and fungemia and other hospital-acquired infections. *Rev Infect Dis* 1985;7(suppl 3):S357–70.
- Schaberg DR, Culver DH, Gaynes RP. Major trends in the microbial etiology of nosocomial infection. *Am J Med* 1991;91(suppl 3B):72S–5S.
- Frimodt-Moller N, Espersen F, Skinhoj P, Rosdahl VT. Changing epidemiology of *Staphylococcus aureus* bacteremia during 1960–1990 in Denmark [abstract no 1441]. In: Program and abstracts of the 33rd Interscience Conference on Antimicrobial Agents and Chemotherapy (New Orleans). Washington, DC: American Society for Microbiology, 1993: 134.
- Musher DM, Lamm N, Darouiche RO, Young EJ, Hamill RJ, Landon GC. The current spectrum of *Staphylococcus aureus* infection in a tertiary care hospital. *Medicine (Baltimore)* 1994;73:186–208.
- Graham DR, Keldermans MM, Klemm LW, Semenza NJ, Shafer ML. Infectious complications among patients receiving home intravenous therapy with peripheral, central, or peripherally placed central venous catheters. *Am J Med* 1991;91(suppl 3B):95S–100S.
- Panlilio AL, Culver DH, Gaynes RP, et al. Methicillin-resistant *Staphylococcus aureus* in U.S. hospitals, 1975–1991. *Infect Control Hosp Epidemiol* 1992;13:582–6.
- Garner JS, Jarvis WR, Emori TG, Horan TC, Hughes JM. CDC definitions for nosocomial infections, 1988. *Am J Infect Control* 1988;16:128–40.
- CDC. Increase in national hospital discharge survey rates for septicemia—United States, 1979–1987. *MMWR Morb Mortal Wkly Rep* 1990;39: 31–4.
- Libman H, Arbeit RD. Complications associated with *Staphylococcus aureus* bacteremia. *Arch Intern Med* 1984;144:541–5.
- Maki DG. Infections caused by intravascular devices used for infusion therapy: pathogenesis, prevention, and management. In: Bisno AL, Waldvogel FA, eds. *Infections associated with indwelling medical devices*. 2nd ed. Washington, DC: American Society for Microbiology, 1994: 155–212.
- Groeger JS, Lucas AB, Thaler HT, et al. Infectious morbidity associated with long-term use of venous access devices in patients with cancer. *Ann Intern Med* 1993;119:1168–74.
- Mermel LA, Maki DG. Detection of bacteremia in adults: consequences of culturing an inadequate volume of blood. *Ann Intern Med* 1993;119: 270–2.
- Scheckler WE, Scheibel W, Kresge D. Temporal trends in septicemia in a community hospital. *Am J Med* 1991;91(suppl 3B):90S–4S.