

Three - Dimensional Organization of Chromosome Territories in the Human Interphase Cell Nucleus

Tobias A. Knoch, Christian Münkkel and Jörg Langowski ¹⁾


Joachim Rauch, Harald Bornfleth and Christoph Cremer ²⁾ with Irina Solovei and Thomas Cremer ³⁾

¹⁾ Division Biophysics of Macromolecules, German Cancer Research Center, Im Neuenheimer Feld 280

²⁾ Institute for Applied Physics, Albert Ueberle Str. 3-5, 69120 Heidelberg, FRG

D - 69120 Heidelberg, Federal Republic of Germany
http://www.DKFZ-Heidelberg.de/Macromol/Welcome.html

³⁾ Institute for Anthropology and Human Genetics, Richard Wagner Str. 10, 80333 Munich, FRG




German Human Genome Project

PURPOSE

The eukaryotic cell is a prime example of a functioning nano machinery. The synthesis of proteins, maintenance of structure and duplication of the eukaryotic cell itself are all fine-tuned biochemical processes that depend on the precise structural arrangement of the cellular components. The regulation of genes – their transcription and replication - has been shown to be connected closely to the three-dimensional organization of the genome in the cell nucleus. Despite the successful linear sequencing of the human genome its three-dimensional structure is widely unknown.

With the simulation of chromosomes and cell nuclei in comparison with fluorescence in situ hybridization we show here an approach leading to the detailed determination of the three-dimensional organization of the human genome:

Best agreement between simulations and experiment is reached for a Multi-Loop-Subcompartment model, thus the human genome shows a higher degree of determinism than previously thought.



Typical textbook illustration of the human cell nucleus:
1) human cell nuclei differ from spherical shape,
2) the DNA is not a closed pipe,
3) nucleosomes are not regularly organized into chromatin,
4) chromatin does not float around randomly in the nucleus.
V. Hennings (illustrator) in Molecular and Cellular Biology by Stephen L. Wolfe, 1993.

SIMULATIONS

Random Walk / Giant Loop model (RW/GL)
Sachs et al. (1995)





Fig. 2

Ray traced image of the Random-Walk/Giant-Loop model, loop size 5Mbp, after ~80.000 Monte-Carlo and 1000 relaxing Brownian Dynamics steps. Large loops intermingle freely thus forming no distinct features like in the MLS model.



Multi-Loop-Subcompartment model (MLS)
Münkkel et al. (1997)






Fig. 3

Ray traced image of the Multi-Loop-Subcompartment model, loop size 126kbp, linker size 126 kbp, after ~50.000 Monte-Carlo and 1000 relaxing Brownian- Dynamics steps. Here rosettes form subcompartments as separated organizational and dynamic entities.



With Monte Carlo and Brownian Dynamics methods we simulated various models (see sketch left) of human interphase chromosome 15 assuming a flexible polymer chain. To save computer power we start with ~3,500 300nm=31kbp and later we relax with ~21,000 50nm=5,2 kbp long segments. For simulation of a single chromosome it is placed in a potential well whose height is related to the excluded volume interaction (EVI). The EVI keeps the chain from self crossing. Starting configurations have the approximate form and size of a metaphase chromosome (Fig. 1) from which the following decondensation into interphase resembles the natural process.


Fig. 1



Rosettes in the Multi-Loop-Subcompartment model correspond to the size of chromosomal interphase band domains. For simulation of a whole interphase nucleus 46 metaphase chromosomes are placed randomly in a nucleus confined by an EVI. The simulations are made on two IBM SP2 parallel computers with 80 and 512 nodes.


Fig. 4

Simulation of a human interphase cell nucleus with all 46 chromosomes with 1,200,000 polymer segments after 0.5s Brownian Dynamics simulation with 10s steps. The MLS-model leads to the formation of distinct chromosome territories and subcompartments.




EXPERIMENTS

Measurement of 3D-Distances between Genomic Markers



Distance Distribution

Fig. 5



Fluorescence in situ hybridization (FISH) in connection with confocal light microscopy is used for the specific marking of small chromosomal DNA regions. Despite the low spatial resolution of FISH, it is possible to interpret the results (f. e. the 3D distance between genetic markers as a function of their genomic distance) with our simulations.

Chromosome 15 and the Prader-Labhard-Willi/Angelmann Syndrome region was chosen, because the genomic distance between markers is well known (see sketch left) and because the PLW/A-syndrome is a candidate for structure mutation (in contrast to the common base pair mutation).

Collaboration with B. Horsthemke, Institute for Human Genetics, Essen, FRG.

Methods: Human fibroblast cells grown on coverslips to confluent layers and being assumed to rest now in the same cell cycle phase are fixed in isotonic environment with paraformaldehyde. For Hybridization we use digoxigenin labeled DNA probes. The probes are detected with fluorescent dyes.

Confocal image series were taken with a Leica TCS NT confocal microscope with an axial displacement of $z = 200\text{nm}$. The images are median and background filtered. After manual threshold determination from an extended focus view (Fig. 6) for spot finding we proceed with image reconstruction specially adjusted to the microscope. Finally the 3D-spatial distances are determined between the centers of mass of the spots (Fig. 5). The experimental distance distributions are then compared to the computed ones. We use a workstation cluster of 10 Silicon Graphics Indigo and Indigo II for computation.

The fibroblast nuclei are found to have their in vivo size (~20m * 10m * 6m) so that we conclude that at least on the micrometer length scale we preserved the nuclear structure. With two colour FISH it is possible to detect 3D-distances below the optical resolution.

Fig. 6

Chromosomes form distinct territories in interphase and genomic markers lay clearly separable within the territories. Left: Territory painting by FISH of chromosome 15; by chance the two territories neighbour each other. Right: Genomic markers YAC48 and YAC60, genomic separation 1Mbp.

CONCLUSION

Best agreement between simulations and experiments is reached for a Multi-Loop-Subcompartment model with a loop size of roughly 126kbp and a linker length of 1,200nm. Supposed that defined loop bases exist it might be possible to determine the mean positioning of genes relative to each other.




Fig. 7

For calculating more general properties of chromosomes the fractal dimension of the chromatin fiber was determined from the simulations. The fractal analysis resulted in multifractal behaviour (data not shown here) in good agreement with predictions drawn from porous network research (Avnir, 1989; Mandelbrot, private communications).




Fig. 8

The simulation of a whole human cell nucleus with all 46 human chromosomes in connection with the simulation of single chromosomes resulted in the formation of distinct chromosome territories as predicted. In contrast to the RW/GL-model the MLS-model leads to low overlap between chromosome territories as well as chromosome arms, in agreement with overlap analysis of confocal image series (data not shown here).

Three-Dimensional Organization of Chromosome Territories in the Human Interphase Nucleus

Knoch, T. A., Münkler, C. & Langowski, J.

*11th Heidelberg Cytometry Symposium, Deutsches Krebsforschungszentrum (DKFZ),
Heidelberg, ISSN 0949-5347, 22th - 24th October, 1998.*

Abstract

The synthesis of proteins, maintenance of structure and duplication of the eukaryotic cell itself are all fine-tuned biochemical processes that depend on the precise structural arrangement of the cellular components. The regulation of genes – their transcription and replication - has been shown to be connected closely to the three-dimensional organization of the genome in the cell nucleus. Despite the successful linear sequencing of the human genome its three-dimensional structure is widely unknown.

The nucleus of the cell has for a long time been viewed as a 'spaghetti soup' of DNA bound to various proteins without much internal structure, except during cell division when chromosomes are condensed into separate entities. Only recently has it become apparent that chromosomes occupy distinct 'territories' also in the interphase, i.e. between cell divisions. In an analogy of the Bauhaus principle that "form follows function" we believe that analyzing in which form DNA is organized in these territories will help us to understand genomic function. We use computer models - Monte Carlo and Brownian dynamics simulations - to develop plausible proposals for the structure of the interphase genome and compare them to experimental data. In the work presented here, we simulate interphase chromosomes for different folding morphologies of the chromatin fiber which is organized into loops of 100kbp to 3 Mbp that can be interconnected in various ways. The backbone of the fiber is described by a wormlike-chain polymer whose diameter and stiffness can be estimated from independent measurements. The implementation describes this polymer as a segmented chain with 3000 to 20000 segments for chromosome 15 depending on the phase of the simulation. The modeling is performed on a parallel computer (IBM SP2 with 80 nodes). We also determine genomic marker distributions within the Prader-Willi-Region on chromosome 15q11.2-13.3. For these measurements we use a fluorescence in situ hybridisation method (in collaboration with I. Solovai, J. Craig and T. Cremer, Munich, FRG) conserving the structure of the nucleus. As probes we use 10 kbp long lambda clones (Prof. B. Horsthemke, Essen, FRG) covering genomic marker distances between 8 kbp and 250 kbp. The markers are detected with confocal and standing wavefield light microscopes (in collaboration with J. Rauch, J. Bradl, C. Cremer and E. Stelzer, both Heidelberg, FRG) and using special image reconstruction methods developed solely for this purpose (developed by R. Eils. and W. Jaeger, Heidelberg, FRG).

Best agreement between simulations and experiments is reached for a Multi-Loop-Subcompartment model with a loop size of 126 kbp which are forming rosetts and are linked by a chromatin linker of 126 kbp. We also hypothesize a different folding structure for maternal versus paternal chromosome 15. In simulations of whole cell nuclei this model also leads to distinct chromosome territories and subcompartments. A fractal analysis of the simulations leads to multifractal behavior in good agreement with predictions drawn from porous network research.

The work is part of the Heidelberg 3D Human Genome Study Group, which is part of the German Human Genome Project.

Corresponding author email contact: TA.Knoch@taknoch.org

Keywords:

Genome, genomics, genome organization, genome architecture, structural sequencing, architectural sequencing, systems genomics, coevolution, holistic genetics, genome mechanics, genome function, genetics, gene regulation, replication, transcription, repair, homologous recombination, simultaneous co-transfection, cell division, mitosis, metaphase, interphase, cell nucleus, nuclear structure, nuclear organization, chromatin density distribution, nuclear morphology, chromosome territories, subchromosomal domains, chromatin loop aggregates, chromatin rosettes, chromatin loops, chromatin fibre, chromatin density, persistence length, spatial distance measurement, histones, H1.0, H2A, H2B, H3, H4, mH2A1.2, DNA sequence, complete sequenced genomes, molecular transport, obstructed diffusion, anomalous diffusion, percolation, long-range correlations, fractal analysis, scaling analysis, exact yard-stick dimension, box-counting dimension, lacunarity dimension, local nuclear dimension, nuclear diffuseness, parallel super computing, grid computing, volunteer computing, Brownian Dynamics, Monte Carlo, fluorescence in situ hybridization, confocal laser scanning microscopy, fluorescence correlation spectroscopy, super resolution microscopy, spatial precision distance microscopy, autofluorescent proteins, CFP, GFP, YFP, DsRed, fusion protein, in vivo labelling.

Literature References

- Knoch, T. A.** Dreidimensionale Organisation von Chromosomen-Domänen in Simulation und Experiment. (Three-dimensional organization of chromosome domains in simulation and experiment.) *Diploma Thesis*, Faculty for Physics and Astronomy, Ruperto-Carola University, Heidelberg, Germany, 1998, and TAK Press, Tobias A. Knoch, Mannheim, Germany, ISBN 3-00-010685-5 and ISBN 978-3-00-010685-9 (soft cover, 2nd ed.), ISBN 3-00-035857-9 and ISBN 978-3-00-0358857-0 (hard cover, 2nd ed.), ISBN 3-00-035858-7, and ISBN 978-3-00-035858-6 (DVD, 2nd ed.), 1998.
- Knoch, T. A., Münkler, C. & Langowski, J.** Three-dimensional organization of chromosome territories and the human cell nucleus - about the structure of a self replicating nano fabrication site. *Foresight Institute - Article Archive*, Foresight Institute, Palo Alto, CA, USA, <http://www.foresight.org>, 1- 6, 1998.