

CLINICAL STUDY

Changes in renal tri-iodothyronine and thyroxine handling during fasting

Edgar J Rolleman^{1,2}, Georg Hennemann¹, Hans van Toor¹, Christian H H Schoenmakers³, Eric P Krenning^{1,2} and Marion de Jong²

¹Department of Internal Medicine III, Academic University Hospital Dijkzigt and Erasmus Medical School, Rotterdam, The Netherlands, ²Department of Nuclear Medicine, Academic University Hospital Dijkzigt and Erasmus Medical School, Rotterdam, The Netherlands and ³Department of Clinical Chemistry, Academic University Hospital Dijkzigt and Erasmus Medical School, Rotterdam, The Netherlands

(Correspondence should be addressed to M de Jong, Department of Nuclear Medicine, Room L 208, Academic Hospital Dijkzigt, Dr. Molewaterplein 40, 3015 GD Rotterdam, The Netherlands)

(H H Schoenmakers is now at Elkerliek Ziekenhuis, loc Helmond, Wesselmanlaan 25, 5707 HA Helmond, The Netherlands)

Abstract

Objective: Liver handling of thyroid hormones (TH) has been known to alter significantly during fasting. This study investigates whether renal handling of TH is also changed during fasting.

Methods: We measured urinary excretion rates and clearances of free tri-iodothyronine (T₃) and free thyroxine (T₄) in healthy subjects prior to and on the third day of fasting.

Results: During fasting, both mean T₃ and T₄ urinary excretion decreased significantly to a mean value of 42% of control. Also, total and free (F) serum T₃ concentrations declined significantly, but serum T₄ did not change. Both FT₃ and FT₄ clearance decreased significantly during fasting (62% and 42% of control). The fasting-induced decrease in uric acid clearance correlated well with the decrease in FT₃ clearance ($r = 0.94$; $P < 0.001$). Serum concentrations of non-esterified fatty acids (NEFA) were significantly elevated during fasting.

Conclusions: The findings cannot be fully explained by the fasting-induced decrease in serum T₃, and are in accordance with inhibition of uptake of T₃ and T₄ at the basolateral membrane of the tubular cell. This inhibition may be caused by a decreased energy state of the tubular cell and by other factors such as ketoacidosis and/or increased NEFA concentrations during fasting.

European Journal of Endocrinology 142 125–130

Introduction

In man, eighty percent of plasma tri-iodothyronine (T₃), the bioactive thyroid hormone, is produced outside the thyroid by deiodination of thyroxine (T₄), which is the main secretory product of the thyroid gland (1). Both liver and kidney cells contain high amounts of the enzyme, type I 5'-deiodinase (2, 3), that converts T₄ to T₃. However, before this intracellular conversion can take place, entry over the plasma membrane into liver and kidney cells has to occur. We showed that uptake of T₄, but also of T₃ and reverse (r) T₃ into liver cells is mediated by an active transport process that is dependent on ATP and on the Na⁺-gradient (4). This process appeared to be the rate-limiting step in the metabolism of thyroid hormones (5).

During fasting and non-thyroidal illness (NTI), a decrease in serum T₃ concentration, a rise in rT₃ and mostly unchanged free (F) and total (T) serum T₄ concentrations are seen (6). These alterations in serum thyroid hormone levels are explained by unchanged

plasma T₄ and rT₃ production rates, combined with both decreased T₄ to T₃ conversion and rT₃ degradation in the liver (6). In addition to decreased 5'-deiodinase activity, catalyzing both T₄ to T₃ formation and rT₃ degradation (7), decreased uptakes of T₄ and rT₃ into the liver have been described as causes for these alterations (6). In a study using a three compartment model for thyroid hormone kinetics, we showed that during caloric deprivation, transport of T₄ into the fast equilibrating compartment (composed of liver and kidney) was diminished (8), resulting in a decline of substrate for T₃ production and consequently leading to a lowered plasma T₃ production rate.

Much attention has been given to the changes in liver handling of thyroid hormones seen in NTI and during fasting (6, 9, 10). However, it is not known whether changes in renal handling of thyroid hormones occur during fasting and NTI. In this study, we aim to illuminate some aspects of renal excretion of thyroid hormones during fasting in healthy subjects, thus creating an experimental form of NTI.

Materials, subjects and methods

Subjects

Four (2 female) healthy volunteers aged 20 to 25 years participated in this study. Thyroid function as evaluated by serum concentrations of TT₄, FT₄, TT₃, FT₃, rT₃ and thyrotropin (TSH) was normal in all subjects. Apart from oral contraceptives used by subject 4, no medication was taken. Indices of nutritional status prior to the fast were normal in all subjects studied (normal serum concentrations of T₃, creatinine, uric acid, urea (5.65 ± 1.06 mmol/l) and free fatty acids).

Study design

On day 0 a normal diet was consumed. Urine was collected for 24 h and a blood sample was taken. From day 1 to 3, the subjects fasted, but a maximal caloric intake of 60 kcal per day in the form of low fat bouillon was allowed. On day 3, another 24-h urine sample was collected and a blood sample was taken. All blood samples were taken at the mid-point of a 24-h collection period.

Urine analysis

Samples of the 24-h urine collection were stored at -20°C until analysis. Uric acid, creatinine and urea concentrations were measured on an EPOS analyzer from Eppendorf (Hamburg, Germany). Urinary T₄ and T₃ concentrations were determined by RIA. Using a highly specific antibody, no thyroid hormone conjugates could be measured. In a set of 3 experiments, the recovery of thyroid hormone added to charcoal-treated urine was measured. A regression line: $y = 0.97x - 0.06$, $P = 0.04$ was obtained. Changing the urinary pH by adding 50 μl of either 0.1 mol/l NaOH or 0.1 mol/l HCl to 950 μl urine did not influence the recovery of added thyroid hormone.

Serum analysis

Serum was stored at -20°C until analysis. Serum TT₄, TT₃, and rT₃ were determined by RIA. Serum FT₄ was

determined using an Amerlite kit from Johnson and Johnson Clinical Diagnostics Ltd (Amersham, Bucks, UK). The serum FT₃ fraction was determined by equilibrium dialysis according to Sterling and Brenner (11), with minor modifications (12). Serum TSH was determined using a TSH Amerlite kit (Johnson and Johnson Clinical Diagnostics Ltd). Urea, creatinine and uric acid were determined on a Chem 1 analyzer from Technicon (Tarrytown, NY, USA). Serum non-esterified fatty acids (NEFA) were measured using a NEFA C-test (Wako Chemicals, Tokyo, Japan).

Calculations and statistical analysis

Clearances of creatinine, uric acid, FT₃ and FT₄ were calculated using the formula $U \cdot V / P \cdot 1.44$, where U is urinary concentration in mol/l, V is urinary volume of 24-h collection period in ml/24 h and P is plasma concentration in mol/l (13).

The significance of differences was determined by Student's *t*-test for paired observations. All data are expressed as means \pm S.D. A *P* value < 0.05 was considered statistically significant.

Results

Table 1 shows the parameters measured to evaluate thyroid function before and on the third day of fasting. Data on all subjects follow the same trend. T₃ concentrations are significantly lowered, and those of rT₃ are elevated after fasting. Total and free T₄ concentrations are unchanged. Serum TSH levels decline but all values remain within the reference range.

In Table 2, urinary excretion of T₃ and T₄ on both the control day and the third day of fasting are shown. Urinary excretion of both T₃ and T₄ is significantly diminished during fasting. Since urinary excretion is dependent on the filtration of plasma free hormone, changes in plasma concentrations will lead to changes in urinary excretion. Therefore, renal clearances of FT₃ and FT₄ are given. In the control period, renal clearance of FT₃ is significantly greater and that of FT₄ significantly smaller than the creatinine clearance. Fasting causes a decline in both renal clearances,

Table 1 Serum thyroid function parameters prior to and on the third day of fasting. Data represent the mean \pm S.D. of 4 healthy subjects, except for serum rT₃ data: $n = 3$, before and on the third day of fasting.

	Control	Fasting	Mean % of control
TSH (mU/l)	0.88 ± 0.46	0.37 ± 0.23^a	42.3
TT ₄ (nmol/l)	103.3 ± 24.8	102.8 ± 26.6	99.4
FT ₄ (pmol/l)	19.4 ± 5.4	18.9 ± 3.9	98.5
T ₃ (nmol/l)	1.93 ± 0.53	1.29 ± 0.32^b	67.8
FT ₃ (pmol/l)	4.2 ± 0.6	2.9 ± 0.7^c	68.2
rT ₃ (nmol/l)	0.27 ± 0.06	0.53 ± 0.16^d	194.8

^a $P = 0.0734$; ^b $P = 0.0291$; ^c $P = 0.0184$; ^d $P = 0.05$ versus control period data.

Table 2 Urinary excretion values (in pmol/24 h) of T_4 and T_3 and the clearances (in ml/min) of FT_3 , FT_4 and creatinine prior to and on the third day of fasting. Data represent the mean \pm s.d. of 4 healthy subjects before and on the third day of fasting.

	Control	Fasting	Mean % of control
T_3 excretion	1367 \pm 273	559 \pm 96 ^a	41.7
FT_3 clearance	224.5 \pm 24.6 ^d	138.8 \pm 31.2 ^{a,f}	61.7
T_4 excretion	1826 \pm 702	783 \pm 535 ^b	41.6
FT_4 clearance	70.3 \pm 35.7 ^e	30.8 \pm 22.8 ^c	41.9
Creatinine clearance	121 \pm 6.6	111.7 \pm 23.0	93.6

^a $P = 0.007$; ^b $P = 0.0123$; ^c $P = 0.0231$ versus control period data. ^d $P = 0.0048$; ^e $P = 0.02$; ^f $P = 0.1616$ versus creatinine clearance in the same period.

resulting in a FT_3 renal clearance that is no longer statistically different from the creatinine clearance ($P = 0.1616$).

In Table 3, serum concentrations, 24-h urinary excretion rates and clearance of uric acid are given. Serum uric acid concentrations rise significantly during fasting, but urinary excretion and clearance of uric acid decrease significantly. Figure 1 shows the correlation between renal clearances of FT_3 and uric acid ($r = 0.94$, $P < 0.001$).

Serum NEFA levels, measured in 3 subjects, rise significantly after 3 days of fasting (0.27 ± 0.22 mmol/l in the control period versus 1.03 ± 0.46 mmol/l in the fasted state, $P = 0.0305$).

Discussion

In this study, we investigated renal handling of thyroid hormones during fasting in man. Our control values of T_4 and T_3 excretion and renal clearances of these hormones in healthy euthyroid persons are in good agreement with reported values obtained in other studies (14–20). The FT_3 renal clearance in healthy humans is significantly greater than the creatinine clearance, the latter being a measure of the glomerular filtration rate (GFR), whereas FT_4 clearance is significantly lower, as has been found by others (14–20) and in this study. These data are consistent with active (tubular) excretion of T_3 into the urine and effective reabsorption of T_4 from the glomerular filtrate in the fed man.

We showed a dramatic decline in urinary excretion of both T_3 and T_4 during fasting. Since the urinary fate of a substance is dependent on its serum concentration,

calculation of the renal clearance is needed to correct for differences in serum concentrations. When calculating the renal clearances, we and others (17–20) used the serum free hormone concentrations since only the free fraction of these hormones is available for glomerular filtration and cellular uptake. The decrease in FT_3 and FT_4 clearance during fasting indicates that, in addition to decreased T_3 serum concentrations, other factors are responsible for the lowered excretion of thyroid hormones into the urine.

Figure 2 depicts a schematic proximal tubule cell of the kidney. Urinary T_3 and T_4 may be considered as the sum of glomerularly filtered thyroid hormone (TH) (pathway 1), plus TH excreted to the urine (pathway 2), minus reabsorbed TH (pathway 3). Glomerular filtration of free TH results in a free hormone concentration, at the start of the proximal tubule, equal to that in serum. Reabsorption of both T_4 and T_3 occurs subsequently, but also uptake at the basolateral membrane of the proximal tubule cell takes place. Cavalieri *et al.* (21) showed that, in humans, the unidirectional renal T_3 uptake rate is about 50 nmol per day. This rate is far greater than the filtered load per day, which is around 0.8 nmol per day (serum FT_3 times GFR, this study). Furthermore, Ferguson *et al.* reported unchanged T_4 uptake in the isolated perfused rat kidney after a decrease in the GFR (22). These observations indicate that thyroid hormones may be taken up at the basolateral membrane of the tubule cell and/or in other kidney cells. Thus, in the fed condition, uptake of TH takes place at both the luminal membrane (reabsorption, pathway 3) and the basolateral membrane (pathway 4). Deiodination of T_4 to

Table 3 Serum concentrations, 24-h urinary excretion rates and clearance of uric acid (UA) prior to and on the third day of fasting. Data represent the mean \pm s.d. of 4 healthy subjects before and on the third day of fasting.

	Control	Fasting	Mean % of control
Serum UA (mmol/l)	0.28 \pm 0.06	0.46 \pm 0.07 ^a	168.5
UA excretion (pmol/24 h)	3.37 \pm 0.82	1.27 \pm 0.75 ^b	40.6
UA clearance (ml/min)	8.65 \pm 1.74	2.05 \pm 1.41 ^c	23.4

^a $P = 0.0002$; ^b $P = 0.009$; ^c $P = 0.0049$ versus control period data.




Figure 1 Plot of the correlation between uric acid (UA) clearance and FT₃ clearance measured in 4 healthy subjects prior to and on the third day of fasting. $r=0.94$, $P<0.001$.

T₃ (pathway 5), excretion into the urine (pathway 2) or secretion to the blood (pathway 6) may follow subsequently.

The lowered excretion and clearances of T₃ and T₄ during fasting may either be caused by stimulation of pathway 3, or by a decrease in pathway 2. It is unlikely that in a state of energy depletion, such as fasting, pathway 3 will be stimulated, as during caloric deprivation in humans uptake in the rapid equilibrating compartment (liver and kidney) is decreased (8). Therefore, a decrease in pathway 2 may be a much more likely explanation for the findings in this study. In a recent study, we showed that cellular efflux of thyroid hormones is a passive process, not influenced by changes in cellular

ATP content, and driven by the intracellular free concentration of the hormone (23). Thus, a lowered rate of efflux via pathway 2 is consistent with a lowered intracellular free hormone concentration. This may be explained by a lowered basolateral uptake (pathway 4) and/or by a lowered luminal uptake (pathway 3).

Inhibition of pathway 3 would result in augmented excretion of TH into the urine, which is evidently not the case in our study. Thus, it remains that inhibition of pathway 4 may be the cause for the decrease in pathway 2 via a decrease in intracellular free hormone concentration.

We showed that uptake of T₃ is diminished in perfused livers of fasted rats. Re-perfusion with energy-enriched perfusion medium restored T₃ uptake within half an hour (7), suggesting involvement of ATP in the transport process that is being lowered during fasting. In liver perfusions with fructose in the medium, ATP concentrations were diminished drastically and T₃ uptake decreased in parallel (10). These studies indicate that cellular ATP is a major determinant of cellular thyroid hormone uptake, as was found earlier in rat and human hepatocytes in culture (24, 25). In the present study, cellular kidney tubular ATP may also be lowered due to fasting.

As a result of fasting, serum concentrations of NEFA rise, as reported earlier (26, 27). In a study with obese subjects undergoing caloric restriction, elevated NEFA concentrations in the sera of these subjects were found to be responsible for the inhibited T₄ uptake into rat hepatocytes (27). Thus, elevated NEFA concentrations may, in addition to lowered cellular energy, be responsible for the suggested inhibition of pathway 4 (Fig. 2).

In the proximal tubule, uric acid is both actively reabsorbed and secreted into the urine (28, 29). In the fasted state, a developed ketoacidemia alters renal




Figure 2 Schematic representation of a proximal tubule cell of the kidney. N=nucleus. For explanation of the different pathways (1–6), see Discussion.

clearance of uric acid by an effect on these tubular transport mechanisms (28, 29). These alterations cause a lowered urinary excretion, with subsequent elevated serum concentrations of uric acid (26, 28, 29). The correlation between FT_3 renal clearance and uric acid clearance suggests that tubular handling of T_3 and uric acid may both be affected by similar mechanisms such as ketoacidosis or a decrease in cellular energy stores having similar or different transport pathways.

In this study, we used creatinine clearance (CrCl) as a marker for GFR, since CrCl values have been shown to correspond well with measured inulin clearances in fasting subjects on the third day of fasting in studies by others (30–32). In the studies of Fox *et al.* (28) and Hoffman *et al.* (33), GFR remains stable over three days of fasting, as in our study.

One of our subjects used oral contraceptives. These drugs are known to alter protein binding of thyroid hormones. Therefore, TT_3 and TT_4 were elevated, whereas free concentrations were normal. It can be questioned whether this medication may have induced alterations in renal handling of T_3 and T_4 in one of our fasting subjects. However, Gaitan *et al.* (15) reported unchanged urinary T_3 excretion in subjects using oral contraceptives, so it is unlikely that this drug interfered in our study, especially since excretion and clearance values of this subject were comparable to those of the other subjects.

In conclusion, fasting induced a significant decline in both T_3 and T_4 urinary excretion and renal clearances, which is, in addition to a decrease in serum FT_3 , consistent with inhibited uptake of T_3 and T_4 at the basolateral membrane of the proximal tubular cell. This uptake inhibition may be explained by lowered energy stores of the tubular cells or by other factors such as ketoacidosis and the detected elevation of serum NEFA levels.

References

- Hennemann G. Thyroid hormone deiodination in healthy man. In *Thyroid Hormone Metabolism*, pp 277–295. Ed G Hennemann. New York and Basel: Marcel Dekker, 1986.
- Yamaki M, Murayama N, Yoshida K, Kusano E, Sakurada T & Asano Y. Subcellular distribution of thyroxine 5'-monodeiodinase activity in rabbit kidney. *Hormone and Metabolic Research* 1991 23 7–11.
- O'Mara BA, Dittrich W, Lauterion TJ & St Germain DL. Pre-translational regulation of type I 5'-deiodinase by thyroid hormones and in fasted and diabetic rats. *Endocrinology* 1993 331 1715–1723.
- Docter R & Krenning EP. Role of cellular transport systems in the regulation of thyroid hormone bioactivity. In *The Thyroid Gland*, pp 233–254. Ed MA Greer. New York: Raven Press, 1990.
- Hennemann G, Krenning EP, Polhuys M, Mol JA, Bernard BF, Visser TJ *et al.* Carrier-mediated transport of thyroid hormone into rat hepatocytes is rate-limiting in total cellular uptake and metabolism. *Endocrinology* 1986 119 1870–1872.
- Docter R, Krenning EP, De Jong M & Hennemann G. The sick euthyroid syndrome: changes in thyroid hormone serum parameters and hormone metabolism. *Clinical Endocrinology* 1993 39 499–518.
- De Jong M, Docter R, Van der Hoek HJ, Vos RA, Krenning EP & Hennemann G. Transport of 3,5,3'-triiodothyronine into the perfused rat liver and subsequent metabolism are inhibited by fasting. *Endocrinology* 1992 113 463–470.
- Van der Heyden JTM, Docter R, Van Toor H, Wilson JHP, Hennemann G & Krenning EP. Effects of caloric deprivation on thyroid hormone tissue uptake and generation of low- T_3 syndrome. *American Journal of Physiology* 1986 251 E156–E163.
- Bauer AGC, Wilson JHP, Lamberts SWJ, Docter R, Hennemann G & Visser TJ. Handling of iodothyronines by liver and kidney in patients with chronic liver disease. *Acta Endocrinologica* 1987 116 339–346.
- De Jong M, Docter R, Bernard BF, Van der Heijden JTM, Van Toor H, Krenning EP *et al.* T_4 uptake into the perfused rat liver and liver T_4 uptake in humans are inhibited by fructose. *American Journal of Physiology* 1994 266 E768–E775.
- Sterling K & Brenner MA. Free thyroxine in human serum: simplified measurement with the aid of magnesium precipitation. *Journal of Clinical Investigation* 1966 45 153–163.
- Lim CE, Bernard HF, De Jong M, Docter R, Krenning EP & Hennemann GA. Furan fatty acid and indoxyl sulfate are the putative inhibitors of thyroxine transport in uremia. *Journal of Clinical Endocrinology and Metabolism* 1993 76 318–324.
- Kasike BL & Keane WF. Laboratory assessment of renal disease: clearance, urinalysis and renal biopsy. In *The Kidney*, edn 5, ch 26, pp 1137–1174. Ed BM Brenner. Philadelphia: WB Saunders Company, 1996.
- Hüfner M & Hesch RD. Tri-iodothyronine determinations in urine. *Lancet* 1973 1 101–102.
- Gaitan JE, Wahner HW, Gorman CA & Jiang N. Measurement of tri-iodothyronine in unextracted urine. *Journal of Laboratory and Clinical Medicine* 1975 86 538–546.
- Burke CW & Shakespear RA. Tri-iodothyronine and thyroxine in urine. II. Renal handling, and effect of urinary protein. *Journal of Clinical Endocrinology and Metabolism* 1976 42 504–513.
- Yoshida K, Sakurada T, Kaise N, Yamamoto M, Kaise K, Saito S *et al.* Measurement of tri-iodothyronine in urine. *Tohoku Journal of Experimental Medicine* 1980 132 389–395.
- Rogowski P, Siersbæk-Nielsen & Hansen JM. Urinary excretion of thyroxine and tri-iodothyronine in different thyroid function states in man. *Acta Endocrinologica* 1978 87 525–534.
- Faber J, Siersbæk-Nielsen & Kirkegaard C. Renal handling of thyroxine, 3,5,3'- and 3,3',5'-tri-iodothyronine, 3,3'- and 3',5'-di-iodothyronine in man. *Acta Endocrinologica* 1987 115 144–148.
- Orden I, Pie J, Juste MG, Marsella JA & Blasco C. Thyroxine in unextracted urine. *Acta Endocrinologica* 1987 114 503–508.
- Cavalieri RR, Moser C, Martin P, Shames D & Perez-Mendez V. Kinetics of renal uptake of l-tri-iodothyronine in man. In *Thyroid Hormone Metabolism*, pp 307–318. Eds WA Harland & JS Orr. London: Academic Press, 1975.
- Ferguson DC & Jennings AS. Regulation of conversion of thyroxine to tri-iodothyronine in perfused rat kidney. *American Journal of Physiology* 1983 245 E220–E229.
- Hennemann G, Krenning EP, Bernard B, Huvers F, Mol J, Docter R *et al.* Regulation of influx and efflux of thyroid hormones in rat hepatocytes: possible physiologic significance of the plasma membrane in the regulation of thyroid hormone activity. *Hormone and Metabolic Research* 1984 14 (Suppl) 1–6.
- Krenning EP, Docter R, Bernard HF, Visser TJ & Hennemann G. Decreased transport of thyroxine (T_4), 3,3',5-tri-iodothyronine (T_3) and 3,3',5'-tri-iodothyronine (rT_3) into rat hepatocytes in primary culture due to a decrease of cellular ATP content and various drugs. *FEBS Letters* 1982 140 229–233.
- De Jong M, Visser TJ, Bernard HF, Docter R, Vos RA, Hennemann G *et al.* Transport and metabolism of iodothyronines in cultured human hepatocytes. *Journal of Clinical Endocrinology and Metabolism* 1993 77 139–143.

- 26 Cahill Jr GF. Starvation in man. *New England Journal of Medicine* 1970 **282** 668–675.
- 27 Lim CF, Docter R, Krenning EP, Van Toor H, Bernard B, De Jong M *et al.* Transport of thyroxine into cultured hepatocytes: effects of mild non-thyroidal illness and calorie restriction in obese subjects. *Clinical Endocrinology* 1994 **40** 79–85.
- 28 Fox IH, Halperin ML, Goldstein MB & Marliss EB. Renal excretion of uric acid during prolonged fasting. *Metabolism* 1976 **25** 551–559.
- 29 Wortmann RL. Uric acid and gout. In *The Kidney: Physiology and Pathophysiology*, edn 2, pp 2971–2991. Eds DW Seldin & G Giebisch. New York: Raven Press, 1992.
- 30 Edgren B & Wester PO. Impairment of glomerular filtration in fasting for obesity. *Acta Medica Scandinavica* 1971 **190** 389–393.
- 31 Veverbrants E & Arky RA. Effects of fasting and refeeding. I. Studies on sodium, potassium and water excretion on a constant electrolyte and fluid intake. *Journal of Clinical Endocrinology* 1969 **29** 55–62.
- 32 Gelman A, Sigulem D, Sustovich DR, Ajzen H & Ramos OL. Starvation and renal function. *American Journal of Medical Science* 1972 **263** 465–471.
- 33 Hoffman RS, Martino JA, Wahl G & Arky RA. Effects of fasting and refeeding. II. Tubular sites of sodium reabsorption and effects of oral carbohydrate on potassium, calcium and phosphate excretion. *Journal of Laboratory and Clinical Medicine* 1969 **74** 915–926.

Received 15 April 1999

Accepted 7 October 1999