



Companion Animals as a Source of Viruses for Human Beings and Food Production Animals

L. A. Reperant^{*}, I. H. Brown[†], O. L. Haenen[‡], M. D. de Jong[§],
A. D. M. E. Osterhaus^{*}, A. Papa[¶], E. Rimstad^{||}, J.-F. Valarcher[#]
and T. Kuiken^{*}

** Department of Viroscience, Erasmus Medical Centre, PO Box 2040, 3000 CA Rotterdam, The Netherlands, † Animal and Plant Health Agency Weybridge, New Haw, Addlestone, Surrey, UK, ‡ National Reference Laboratory for Fish, Shellfish and Crustacean Diseases, Central Veterinary Institute of Wageningen UR, PO Box 65, 8200 AB Lelystad, § Department of Medical Microbiology, Academic Medical Centre, Meibergdreef 9, 1105 AZ Amsterdam, The Netherlands, ¶ Department of Microbiology, Medical School Aristotle University of Thessaloniki, Thessaloniki, Greece, || Department of Food Safety and Infection Biology, University of Life Sciences, Oslo, Norway and # Department of Virology, Immunology, and Parasitology, National Veterinary Institute, Uppsala, Sweden*

Summary

Companion animals comprise a wide variety of species, including dogs, cats, horses, ferrets, guinea pigs, reptiles, birds and ornamental fish, as well as food production animal species, such as domestic pigs, kept as companion animals. Despite their prominent place in human society, little is known about the role of companion animals as sources of viruses for people and food production animals. Therefore, we reviewed the literature for accounts of infections of companion animals by zoonotic viruses and viruses of food production animals, and prioritized these viruses in terms of human health and economic importance. In total, 138 virus species reportedly capable of infecting companion animals were of concern for human and food production animal health: 59 of these viruses were infectious for human beings, 135 were infectious for food production mammals and birds, and 22 were infectious for food production fishes. Viruses of highest concern for human health included hantaviruses, Tahyna virus, rabies virus, West Nile virus, tick-borne encephalitis virus, Crimean–Congo haemorrhagic fever virus, Aichi virus, European bat lyssavirus, hepatitis E virus, cowpox virus, G5 rotavirus, influenza A virus and lymphocytic choriomeningitis virus. Viruses of highest concern for food production mammals and birds included bluetongue virus, African swine fever virus, foot-and-mouth disease virus, lumpy skin disease virus, Rift Valley fever virus, porcine circovirus, classical swine fever virus, equine herpesvirus 9, peste des petits ruminants virus and equine infectious anaemia virus. Viruses of highest concern for food production fishes included cyprinid herpesvirus 3 (koi herpesvirus), viral haemorrhagic septicaemia virus and infectious pancreatic necrosis virus. Of particular concern as sources of zoonotic or food production animal viruses were domestic carnivores, rodents and food production animals kept as companion animals. The current list of viruses provides an objective basis for more in-depth analysis of the risk of companion animals as sources of viruses for human and food production animal health.

Crown Copyright © 2016 Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Keywords: companion animal; livestock; virus; zoonosis

Contents

Introduction	S42
Materials and Methods	S42
Literature Search	S42

Correspondence to: T. Kuiken (e-mail: t.kuiken@erasmusmc.nl).

Semiquantitative Prioritization	S43
Results	S46
Discussion	S47
Acknowledgments	S49
Supplementary data	S49
References	S49

Introduction

Little is known about the role of companion animals in the transmission of pathogens, in particular viruses, to man or to livestock. A plethora of viruses have zoonotic potential (Gortazar *et al.*, 2014) or are of economic importance as they infect food production animals (Morgan and Prakash, 2006). Studies aiming at inventorying viruses with zoonotic potential have revealed their expanding diversity with an average of three to four new zoonotic pathogens identified each year (Woolhouse *et al.*, 2012). Wild animal species commonly are the source of these novel or newly identified pathogens. New virus species, and sometimes even new virus genera, have been reported increasingly in wild animal species, such as bats or rodents (Drexler *et al.*, 2012, 2013).

Viruses of economic importance, because of their ability to infect and cause disease in food production animal species (and therefore their ability to undergo transboundary spread), have long been listed by international organizations, such as the Office International des Epizooties (OIE; World Organisation for Animal Health; www.oie.int), with associated regulatory phytosanitary measures. In addition, as for zoonotic pathogens, new pathogens are recognized occasionally to infect food production animals. Wild animal species are likewise often implicated in the transmission of such newly discovered pathogens, both to human beings and to livestock species (Field *et al.*, 2007; Raj *et al.*, 2014).

Companion animals, in particular domestic carnivores, also are hosts of an expanding diversity of viruses. For example, canine parvovirus, now a major pathogen of domestic dogs worldwide, emerged in the 1970s (Parrish, 1999). Influenza A viruses that appear to be maintained in domestic dog populations have only recently emerged (Dubovi, 2010). However, populations of companion animal species are expanding both in size and diversity, in particular in industrialized countries. In this review, companion animals are defined as any domesticated, domestic-bred or wild-caught animals, permanently living in human communities and kept by people for company, amusement, work (e.g. support for blind or deaf people, police or military dogs) or psychological support. These include dogs, cats, horses, rabbits, ferrets,

guinea pigs, reptiles, birds and ornamental fish, but also food production animals, such as domestic pigs, kept as companion animals. These changes in populations of companion animals have so far not been matched with an assessment of the role of these animals as a source of zoonotic viruses and viruses of economic importance in livestock. Here, we review the literature on the occurrence of infections of companion animal species by zoonotic viruses and viruses of food production animals in order to establish a complete list of viruses that could possibly be transmitted from companion animals to man or livestock, based on published records of cross-species transmission. We further prioritize these viruses semiquantitatively, in terms of health and economic importance, by ranking the likelihood that companion animals act as relevant sources for the cross-species transmission of the listed viruses, and associated potential impact, with a special focus on Europe.

Materials and Methods

Literature Search

We retrieved articles published before 1st July 2012 in PubMed to identify and list mammalian, avian and fish viruses that have proven ability to cross the species barriers between companion animals and man or between companion animals and food production animals. To this end, PubMed was searched for combinations of terms belonging to the following general categories: ‘virus’ AND ‘companion animal’ AND ‘food production animal’ OR ‘human’ (Supplementary Tables 1a and 1b). Viruses of food production animals (without report of infection in companion animals) were also included, to account for the risk posed by viruses transmitted from ruminants, pigs and poultry kept as companion animals to people or food production animals. Zoonotic viruses and viruses of food production animals were considered capable of infecting companion animals, based on published positive polymerase chain reaction (PCR) data, viral isolation or serology in the latter. Accordingly, a total of 170 viruses or virus groups at or below genus level were included (Supplementary Table 2).

Semiquantitative Prioritization

Based on a set of priority criteria (Supplementary Table 3a), these viruses were ranked for the likelihood that they could be transmitted from companion animals to man or food production animals, and the associated potential impact of such transmission. Particular attention was given to possible demographic changes that are currently occurring or that may occur in the near future, and which may increase the likelihood of cross-species transmission of viruses from companion animals to man or food production animals. For example, expanding populations of a particular companion animal species, or emergence of a particular virus, may translate into increased likelihood of cross-species transmission.

The prioritization criteria included the trends in the companion animal populations that are putatively the source of the particular virus, its host range, the level of environmental resistance, the clinical impact in man or food production animals and, for viruses transmitted to food production animal species, the population size of the food production animal species, and whether the virus is on the OIE list of notifiable terrestrial and aquatic animal diseases. We further classified the current epidemiological situation for each virus in Europe as absent, endemic or emerging.

As mentioned earlier, expanding populations of a companion animal species may favour cross-species transmission of the pathogens it may host. Populations of companion animals were categorized into either stable, increasing or booming. Pathogens with a wide host range are considered more likely to cross species barriers than more species-specific pathogens. Accordingly, we categorized the listed viruses into viruses infecting less than or more than four different host species. Viruses that may survive in the environment can rely on indirect transmission, which may increase the likelihood of cross-species transmission. We categorized the viruses into labile or resistant, based on the presence or absence of a lipid envelope, respectively. In order to assess the potential impact of the listed viruses, the severity of the resulting clinical disease was categorized into four categories, from no disease to severe disease. Viruses that can infect food production animals were further divided into those listed and those not listed as notifiable by the OIE, and their clinical impact was weighted by the size of the food production animal populations, categorized semiquantitatively into three classes.

The rank was calculated based on the sum of all criteria and weighted by the epidemiological situa-

tion in Europe (Supplementary Table 3b). The ranks were calculated separately for viruses transmitted from companion animals to man, for viruses transmitted from companion animals to mammalian and avian food production animals and for viruses transmitted from ornamental fish to food production fish species. They were standardized on a unit scale for comparison purposes.

The semiquantitative scores of all viruses and of viruses of relevance to Europe were compared, using the non-parametric Mann–Whitney test. Differences were considered significant when $P < 0.05$. The distributions of viruses were represented by Venn diagrams produced in ‘R’ using the Venneuler package.




Fig. 1. Semiquantitative prioritization scores. (A) Zoonotic viruses. (B) Viruses of mammalian and avian food production animals. (C) Viruses of fish. Scores of all listed viruses are represented by light blue bars; scores of viruses that are present or with emerging potential in Europe are represented by dark blue bars.

Table 1
List of viruses with highest prioritization scores that may be transmitted from companion animals to man and/or food production animals

<i>Family</i>	<i>Virus</i>	<i>Zoonotic</i>	<i>Food production animals</i>	<i>Vector-borne</i>	<i>Selected references</i>
Adenoviridae	Avian adenoviruses		X		McFerran and Smyth (2000)
Arteriviridae	Porcine reproductive and respiratory syndrome virus		X		Cho and Dee (2006)
Asfarviridae	African swine fever virus		X	V	Marshall <i>et al.</i> (2007), Sipos <i>et al.</i> (2007), Costard <i>et al.</i> (2009)
Birnaviridae	Infectious bursal disease virus		X		Mahgoub <i>et al.</i> (2012)
	Infectious pancreatic necrosis virus		X		Dobos (1995), Munro and Midtlyng (2011)
Bunyaviridae	Rift Valley fever virus	X	X	V	Gerdes (2004)
	Hantaviruses	X			Desmyter <i>et al.</i> (1983), Vapalahti <i>et al.</i> (2003), Heyman <i>et al.</i> (2009), Olsson <i>et al.</i> (2010), Zhang <i>et al.</i> (2010), Dobby <i>et al.</i> (2012), Papa (2012)
	Dobrava—Belgrade virus	X			Heyman <i>et al.</i> 2009, Papa (2012)
	Crimean—Congo haemorrhagic fever virus	X		V	Shepherd <i>et al.</i> (1987), Capua (1998), Whitehouse (2004), Ergonul (2006)
Caliciviridae	Vesicular exanthema virus		X		Lin and Kitching (2000)
Circoviridae	Chicken infectious anaemia virus		X		Schat (2009)
	Porcine circovirus		X		Li <i>et al.</i> (2010), Grau-Roma <i>et al.</i> (2011), Baekbo <i>et al.</i> (2012)
Coronaviridae	Avian bronchitis virus		X		Sjaak de Wit <i>et al.</i> (2011), Cook <i>et al.</i> (2012)
	Swine transmissible gastroenteritis virus		X		Wood (1979)
Flaviviridae	Bovine viral diarrhoea disease virus		X		Lindberg <i>et al.</i> (2006), Walz <i>et al.</i> (2010)
	Powassan virus	X		V	Charrel <i>et al.</i> (2004)
	Japanese encephalitis virus	X	X	V	Mackenzie <i>et al.</i> (2004)
	Classical swine fever virus		X		Moennig <i>et al.</i> (2003)
	West Nile virus	X	X	V	Hubalek and Halouzka (1999), Campbell <i>et al.</i> (2002), Austgen <i>et al.</i> (2004), Zeller and Schuffenecker (2004), Crespo <i>et al.</i> (2009), Danis <i>et al.</i> (2011)
	Tick-borne encephalitis virus	X		V	Gritsun <i>et al.</i> (2003), Gover <i>et al.</i> (2011), Pfeffer and Dobler (2011), Hubalek and Rudolf (2012)

(Continued)

Table 1 (continued)

Family	Virus	Zoonotic	Food production animals	Vector-borne	Selected references
Hepeviridae	Hepatitis E virus	X			Kuno <i>et al.</i> (2003), Renou <i>et al.</i> (2007), Vasickova <i>et al.</i> (2007), de Deus <i>et al.</i> (2008), Schielke <i>et al.</i> (2009), Lewis <i>et al.</i> (2010), Meng (2010), Pavio <i>et al.</i> (2010), Rutjes <i>et al.</i> (2010), Kamar <i>et al.</i> (2012)
Herpesviridae	Gallid herpesvirus 1 (laryngotracheitis virus)		X		Bagust <i>et al.</i> (2000), Fuchs <i>et al.</i> (2007)
	Ovine herpesvirus 2 (malignant catarrhal fever virus)		X		Russell <i>et al.</i> (2009)
	Suid herpesvirus 1 (pseudorabies/ Aujeszky virus)		X		Tischer and Osterrieder (2010), Muller <i>et al.</i> (2011)
	Equine herpesvirus 9		X		Schrenzel <i>et al.</i> (2008), Osterrieder and Van de Walle (2010)
	Cyprinid herpesvirus 3 (koi herpesvirus)		X		Hedrick <i>et al.</i> (2000), Haenen <i>et al.</i> (2004), Haenen and Hedrick (2006), Perelberg <i>et al.</i> (2008), Way (2009)
Iridoviridae	Infectious spleen and kidney necrosis virus		X		Jeong <i>et al.</i> (2008)
	European catfish virus		X		Whittington <i>et al.</i> (2010)
	Lymphocystis disease virus		X		Paperna <i>et al.</i> (2001), Hossain <i>et al.</i> (2008)
Nodaviridae	Viral nervous necrosis virus		X		Hegde <i>et al.</i> (2003), Furusawa <i>et al.</i> (2007), Mazelet <i>et al.</i> (2011)
Orthomyxoviridae	Infectious salmon anaemia virus		X		Cottet <i>et al.</i> (2011), Rimstad <i>et al.</i> (2011)
	Influenza A virus	X	X		Olsen (2000), Myers <i>et al.</i> (2007), Thiry <i>et al.</i> (2007), Peiris (2009), Harder and Vahlenkamp (2010), Swenson <i>et al.</i> (2010), Reperant <i>et al.</i> (2012)
Paramyxoviridae	Newcastle disease virus (APMV 1)		X		Alexander (2011), Alexander <i>et al.</i> (2012)
	Nipah virus	X	X		Luby and Gurley (2012), Clayton <i>et al.</i> (2013)
Picornaviridae	Hendra virus	X	X		
	Encephalomyocarditis virus		X		Carocci and Bakkali- Kassimi (2012)
	Foot-and-mouth disease virus	X	X		USDA (1994) Anonymous (2006), Marshall <i>et al.</i> (2007), Valarcher <i>et al.</i> (2008)
	Avian encephalomyelitis virus		X		Welchman Dde <i>et al.</i> (2009)
	Aichi virus	X			Le Guyader <i>et al.</i> (2008), Li <i>et al.</i> (2011), Phan <i>et al.</i> (2011), Reuter <i>et al.</i> (2011)

(Continued)

Table 1 (continued)

Family	Virus	Zoonotic	Food production animals	Vector-borne	Selected references
Polyomaviridae	Avian polyomaviruses		X		Johne <i>et al.</i> (2006)
Poxviridae	Monkeypox virus	X			Reed <i>et al.</i> (2004), Tack and Reynolds (2011)
	Lumpy skin disease virus		X		Babiuk <i>et al.</i> (2008), Bowden <i>et al.</i> (2009), Tuppurainen and Oura (2012)
Reoviridae	Salmon gill poxvirus		X		Nylund <i>et al.</i> (2008)
	Bluetongue virus		X	V	Akita <i>et al.</i> (1994), Alexander <i>et al.</i> (1994), Brown <i>et al.</i> (1996), Mellor and Wittmann (2002), Oura and El Harrak (2011)
	Piscine reovirus		X		Palacios <i>et al.</i> (2010)
	G5 rotavirus	X			Esona <i>et al.</i> (2004), Santos and Hoshino (2005), Ahmed <i>et al.</i> (2007), Duan <i>et al.</i> (2007), Midgley <i>et al.</i> (2012), Mladenova <i>et al.</i> (2012)
Rhabdoviridae	Spring viraemia of carp virus		X		Haenen and Davidse (1993), Ahne <i>et al.</i> (2002), Gomez-Casado <i>et al.</i> (2011)
	Viral haemorrhagic septicaemia virus		X		Skall <i>et al.</i> (2005), Smail and Snow (2011)
	Vesicular stomatitis virus		X		Schmitt (2002)
	Rabies virus	X	X		Rupprecht <i>et al.</i> (2001), De Benedictis <i>et al.</i> (2008), Capello <i>et al.</i> (2010)
Togaviridae	Western equine encephalitis virus	X		V	Zacks and Paessler (2010)
	Eastern equine encephalitis virus	X		V	Zacks and Paessler (2010)
	Salmon pancreas disease virus (salmonid alphavirus)		X		McLoughlin and Graham (2007)

Highlighted rows are viruses relevant for Europe. Relevance for zoonotic transmission and transmission to food productions animals is indicated by X; vector-borne viruses are indicated by V.

Results

The list of viruses reportedly capable of infecting companion animal species and man or food production animal species comprises 138 virus species, of which 26 are vector borne, and 32 groups of virus species at the genus level or below (Supplementary Table 2). ‘Vector borne’ refers here to biological rather than mechanical transmission by arthropod vectors. Fifty-nine virus species or groups of species have zoonotic potential, of which 17 are vector borne; 135 infect mammalian or avian food production animal species, of which 24 are vector borne, and 22 infect

food production fish species. Of the fish viruses, only six species have been demonstrated in ornamental fish. The distributions of standardized semiquantitative positive prioritization scores are presented in Fig. 1. The scores are significantly lower for zoonotic viruses and viruses of mammalian and avian food production animals relevant to Europe than those considered in the global list ($Z = 2.21$, $P = 0.02$ and $Z = 6.82$, $P < 0.0001$, respectively). Although a similar trend was observed for fish viruses, this difference was not significant.

Viruses that received scores > 0.5 are highlighted in Supplementary Table 2. Viruses with the highest

prioritization scores, hence those which may be at the highest risk for cross-species transmission from companion animals to man and/or food production animals, are listed in Table 1. Additional virus species with high prioritization scores and of relevance to Europe include European bat lyssavirus (Fooks *et al.*, 2003; van der Poel *et al.*, 2006; Dacheux *et al.*, 2009), cowpox virus (Baxby *et al.*, 1994; Baxby and Bennett, 1997; Pfeffer *et al.*, 2002; Wolfs *et al.*, 2002; Tack, 2011; von Bomhard *et al.*, 2011), lymphocytic choriomeningitis virus (Gregg, 1975; Rousseau *et al.*, 1997; Barton and Mets, 2001; Amman *et al.*, 2007), Puumala virus (Vapalahti *et al.*, 2003; Heyman *et al.*, 2009; Olsson *et al.*, 2010), Tahyna virus (Hubalek *et al.*, 2005; Hubalek, 2008) and peste des petits ruminants virus (Banyard *et al.*, 2010).

Companion animals were categorized subsequently into domestic carnivores, small mammals, horses, birds and food production animals kept as companion animals. These harbored, respectively, 42, 54, 40, 40 and 91 viruses with zoonotic potential or of economic importance. The distribution of viruses with zoonotic

potential, of economic importance for food production animals, and those transmitted by arthropod vectors are shown in Fig. 2.

The companion animal sources of viruses that can infect man and/or food production animals are shown in Fig. 3. Viruses with zoonotic potential have the greatest diversity in companion animal sources, with more limited overlap between host species. So far, there are no reports of fish as a source of zoonotic viruses. Because ruminants, pigs, horses, rabbits and poultry have companion animal counterparts, viruses infecting these species may originate from food production animals kept as companion animals. Small mammals, in particular rodents, are hosts to the second largest number of viruses of economic importance that may infect food production animals.

Discussion

The present study is an initial attempt to delineate the importance of companion animals in the transmission of viruses to man and food production




Fig. 2. Distribution of zoonotic viruses, viruses of mammalian and avian food production animals and vector-borne viruses in companion animal species. (A) All listed viruses. (B) Viruses of domestic carnivores (i.e. dogs, cats and ferrets). (C) Viruses of small mammals (i.e. rodents and rabbits). (D) Viruses of horses. (E) Viruses of birds (i.e. ornamental birds and poultry kept as companion animals). (F) Viruses of mammalian food production animals kept as companion animals (i.e. cattle, sheep, goats and pigs).




Fig. 3. Distribution of viruses of companion animal species in man and in mammalian and avian food production animals. (A) Zoonotic viruses. (B) Viruses of cattle. (C) Viruses of small ruminants (i.e. sheep and goats). (D) Viruses of pigs. (E) Viruses of horses. (F) Viruses of rabbits. (G) Viruses of poultry.

animals. The resulting list of viruses potentially transmitted by companion animals likely is conservative, as reports of actual cross-species transmission events from companion animals to man or food production animals remain rare and non-systematic. Viruses were included based on reports of actual infection in both companion animals and man or food production animals, irrespective of the direction of cross-species transmission. For example, there is no direct evidence to date that companion animals play a role in the transmission of vector-borne viruses to man or food production animals. However, 10 vector-borne virus species cumulated high prioritization scores, suggesting a potential for such cross-species transmission.

Nonetheless, our use of a systematic and objective selection method to assess the potential importance

of companion animal viruses for human health and food production brings a number of advantages compared with the sole reliance on expert opinion. Expert opinion is undeniably valuable, but has the inherent risk of becoming a self-fulfilling prophecy: if pathogen testing is limited to those considered important by experts, the results will be limited to those pathogens. In contrast, the current list includes some pathogens that most experts would not have scored highly (e.g. Aichi virus and G5 rotavirus as zoonotic viruses, and lumpy skin disease virus and equine herpesvirus 9 as viruses of food production animals). Of note, the recent incursions of lumpy skin disease in cattle in Europe (Cyprus, Greece) and neighbouring countries (Turkey and Russia), as reported by Promed, support the need to expect the unexpected.

Due to their population size, in particular in industrialized countries, domestic carnivores and rodents are among the most important companion animal sources of zoonotic viruses. On the other hand, expanding populations of food production animals kept as companion animals (e.g. domestic pigs, Sipos *et al.*, 2007) may also result in the cross-species transmission of zoonotic viruses, which represent about half of the viruses they harbour. Rodents also harbour a large number of viruses of economic importance that can infect food production animals.

The present analysis does not include the potential for direct or indirect contact necessary for cross-species transmission between host species, since data for such an assessment are largely lacking. While the current list of viruses potentially transmitted from companion animals thus has its limitations, it represents a solid and objectively constructed basis for more in-depth risk assessment analyses.

Acknowledgments

We acknowledge N. Tordo as member of the CALLISTO Expert Advisory Group on Virology and for participating in the project meetings. This study was performed as part of the FP7 European project CALLISTO (project number 289316). The present address of L. A. Reperant is: Artemis One Health Research Foundation, Yalelaan 1, Utrecht, The Netherlands. The present address of A. D. M. E. Osterhaus is: Research Center for Emerging Infections and Zoonoses, University of Veterinary Medicine, Bünteweg 17, Hannover, Germany. The present address of J.-F. Valarcher is: Department of Clinical Sciences, Swedish University of Agricultural Sciences, Box 7054, SE-750 07 Uppsala, Sweden.

Supplementary data

Supplementary data related to this article can be found at <http://dx.doi.org/10.1016/j.jcpa.2016.07.006>.

References

- Ahmed K, Anh DD, Nakagomi O (2007) Rotavirus G5P [6] in child with diarrhea, Vietnam. *Emerging Infectious Diseases*, **13**, 1232–1235.
- Ahne W, Bjorklund HV, Essbauer S, Fijan N, Kurath G *et al.* (2002) Spring viremia of carp (SVC). *Diseases of Aquatic Organisms*, **52**, 261–272.
- Akita GY, Ianconescu M, MacLachlan NJ, Osburn BI (1994) Bluetongue disease in dogs associated with contaminated vaccine. *Veterinary Record*, **134**, 283–284.
- Alexander DJ (2011) Newcastle disease in the European Union 2000 to 2009. *Avian Pathology*, **40**, 547–558.
- Alexander DJ, Aldous EW, Fuller CM (2012) The long view: a selective review of 40 years of Newcastle disease research. *Avian Pathology*, **41**, 329–335.
- Alexander KA, MacLachlan NJ, Kat PW, House C, O'Brien SJ *et al.* (1994) Evidence of natural bluetongue virus infection among African carnivores. *American Journal of Tropical Medicine and Hygiene*, **51**, 568–576.
- Amman BR, Pavlin BI, Albarino CG, Comer JA, Erickson BR *et al.* (2007) Pet rodents and fatal lymphocytic choriomeningitis in transplant patients. *Emerging Infectious Diseases*, **13**, 719–725.
- Anonymous (2006) Assessing the risk of foot-and-mouth disease introduction into the EU from developing countries. *European Food Safety Agency Journal*, **313**, 1–34.
- Austgen LE, Bowen RA, Bunning ML, Davis BS, Mitchell CJ *et al.* (2004) Experimental infection of cats and dogs with West Nile virus. *Emerging Infectious Diseases*, **10**, 82–86.
- Babiuk S, Bowden TR, Boyle DB, Wallace DB, Kitching RP (2008) Capripoxviruses: an emerging worldwide threat to sheep, goats and cattle. *Transboundary and Emerging Diseases*, **55**, 263–272.
- Baekbo P, Kristensen CS, Larsen LE (2012) Porcine circovirus diseases: a review of PMWS. *Transboundary and Emerging Diseases*, **59**(Suppl. 1), 60–67.
- Bagust TJ, Jones RC, Guy JS (2000) Avian infectious laryngotracheitis. *Revue Scientifique et Technique de l'OIIE*, **19**, 483–492.
- Banyard AC, Parida S, Batten C, Oura C, Kwiatek O *et al.* (2010) Global distribution of peste des petits ruminants virus and prospects for improved diagnosis and control. *Journal of General Virology*, **91**, 2885–2897.
- Barton LL, Mets MB (2001) Congenital lymphocytic choriomeningitis virus infection: decade of rediscovery. *Clinical Infectious Diseases*, **33**, 370–374.
- Baxby D, Bennett M (1997) Cowpox: a re-evaluation of the risks of human cowpox based on new epidemiological information. *Archives of Virology*, **13**(Suppl.), 1–12.
- Baxby D, Bennett M, Getty B (1994) Human cowpox 1969–93: a review based on 54 cases. *British Journal of Dermatology*, **131**, 598–607.
- Bowden TR, Coupar BE, Babiuk SL, White JR, Boyd V *et al.* (2009) Detection of antibodies specific for sheeppox and goatpox viruses using recombinant capripoxvirus antigens in an indirect enzyme-linked immunosorbent assay. *Journal of Virological Methods*, **161**, 19–29.
- Brown CC, Rhyan JC, Grubman MJ, Wilbur LA (1996) Distribution of bluetongue virus in tissues of experimentally infected pregnant dogs as determined by in-situ hybridization. *Veterinary Pathology*, **33**, 337–340.
- Campbell GL, Marfin AA, Lanciotti RS, Gubler DJ (2002) West Nile virus. *Lancet Infectious Diseases*, **2**, 519–529.
- Capello K, Mulatti P, Comin A, Gagliazzo L, Guberti V *et al.* (2010) Impact of emergency oral rabies vaccination of foxes in northeastern Italy, 28 December 2009–20 January 2010: preliminary evaluation. *Euro Surveillance*, **15**, 19617.

- Capua I (1998) Crimean–Congo haemorrhagic fever in ostriches: a public health risk for countries of the European Union? *Avian Pathology*, **27**, 117–120.
- Carocci M, Bakkali-Kassimi L (2012) The encephalomyocarditis virus. *Virulence*, **3**, 351–367.
- Charrel RN, Attoui H, Butenko AM, Clegg JC, Deubel V *et al.* (2004) Tick-borne virus diseases of human interest in Europe. *Clinical Microbiology and Infection*, **10**, 1040–1055.
- Cho JG, Dee SA (2006) Porcine reproductive and respiratory syndrome virus. *Theriogenology*, **66**, 655–662.
- Clayton BA, Wang LF, Marsh GA (2013) Henipaviruses: an updated review focusing on the pteropid reservoir and features of transmission. *Zoonoses and Public Health*, **60**, 69–83.
- Cook JK, Jackwood M, Jones RC (2012) The long view: 40 years of infectious bronchitis research. *Avian Pathology*, **41**, 239–250.
- Costard S, Wieland B, de Glanville W, Jori F, Rowlands R *et al.* (2009) African swine fever: how can global spread be prevented? *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, **364**, 2683–2696.
- Cottet L, Rivas-Aravena A, Cortez-San Martin M, Sandino AM, Spencer E (2011) Infectious salmon anemia virus – genetics and pathogenesis. *Virus Research*, **155**, 10–19.
- Crespo R, Shivaprasad HL, Franca M, Woolcock PR (2009) Isolation and distribution of West Nile virus in embryonated chicken eggs. *Avian Diseases*, **53**, 608–612.
- Dacheux L, Larrous F, Mailles A, Boisseleau D, Delmas O *et al.* (2009) European bat lyssavirus transmission among cats, Europe. *Emerging Infectious Diseases*, **15**, 280–284.
- Danis K, Papa A, Theocharopoulos G, Dougas G, Athanasiou M *et al.* (2011) Outbreak of West Nile virus infection in Greece, 2010. *Emerging Infectious Diseases*, **17**, 1868–1872.
- De Benedictis P, Gallo T, Iob A, Coassin R, Squecco G *et al.* (2008) Emergence of fox rabies in north-eastern Italy. *Euro Surveillance*, **13**, pii: 19033.
- de Deus N, Peralta B, Pina S, Allepuz A, Mateu E *et al.* (2008) Epidemiological study of hepatitis E virus infection in European wild boars (*Sus scrofa*) in Spain. *Veterinary Microbiology*, **129**, 163–170.
- Desmyter J, LeDuc JW, Johnson KM, Brasseur F, Deckers C *et al.* (1983) Laboratory rat associated outbreak of haemorrhagic fever with renal syndrome due to Hantaan-like virus in Belgium. *Lancet*, **322**, 1445–1448.
- Dobly A, Cochez C, Goossens E, De Bosschere H, Hansen P *et al.* (2012) Sero-epidemiological study of the presence of hantaviruses in domestic dogs and cats from Belgium. *Research in Veterinary Science*, **92**, 221–224.
- Dobos P (1995) The molecular biology of infectious pancreatic necrosis virus. *Annual Review of Fish Diseases*, **5**, 25–54.
- Drexler JF, Corman VM, Muller MA, Lukashev AN, Gmyl A *et al.* (2013) Evidence for novel hepaciviruses in rodents. *PLoS Pathogens*, **9**, e1003438.
- Drexler JF, Corman VM, Muller MA, Maganga GD, Vallo P *et al.* (2012) Bats host major mammalian paramyxoviruses. *Nature Communications*, **3**, 796.
- Duan ZJ, Li DD, Zhang Q, Liu N, Huang CP *et al.* (2007) Novel human rotavirus of genotype G5P[6] identified in a stool specimen from a Chinese girl with diarrhea. *Journal of Clinical Microbiology*, **45**, 1614–1617.
- Dubovi EJ (2010) Canine influenza. *Veterinary Clinics of North America: Small Animal Practice*, **40**, 1063–1071.
- Ergonul O (2006) Crimean–Congo haemorrhagic fever. *Lancet Infectious Diseases*, **6**, 203–214.
- Esona MD, Armah GE, Geyer A, Steele AD (2004) Detection of an unusual human rotavirus strain with G5P[8] specificity in a Cameroonian child with diarrhea. *Journal of Clinical Microbiology*, **42**, 441–444.
- Field HE, Mackenzie JS, Daszak P (2007) Henipaviruses: emerging paramyxoviruses associated with fruit bats. *Current Topics in Microbiology and Immunology*, **315**, 133–159.
- Fooks AR, Brookes SM, Johnson N, McElhinney LM, Hutson AM (2003) European bat lyssaviruses: an emerging zoonosis. *Epidemiology and Infection*, **131**, 1029–1039.
- Fuchs W, Veits J, Helferich D, Granzow H, Teifke JP *et al.* (2007) Molecular biology of avian infectious laryngotracheitis virus. *Veterinary Research*, **38**, 261–279.
- Furusawa R, Okinaka Y, Uematsu K, Nakai T (2007) Screening of freshwater fish species for their susceptibility to a betanodavirus. *Diseases of Aquatic Organisms*, **77**, 119–125.
- Gerdes GH (2004) Rift Valley fever. *Revue Scientifique et Technique de l'OIE*, **23**, 613–623.
- Gomez-Casado E, Estepa A, Coll JM (2011) A comparative review on European-farmed finfish RNA viruses and their vaccines. *Vaccine*, **29**, 2657–2671.
- Gortazar C, Reperant LA, Kuiken T, de la Fuente J, Boadella M *et al.* (2014) Crossing the interspecies barrier: opening the door to zoonotic pathogens. *PLoS Pathogens*, **10**, e1004129.
- Gover L, Kirkbride H, Morgan D (2011) Public health argument to retain current UK national controls for tick and tapeworms under the Pet Travel Scheme. *Zoonoses and Public Health*, **58**, 32–35.
- Grau-Roma L, Fraile L, Segales J (2011) Recent advances in the epidemiology, diagnosis and control of diseases caused by porcine circovirus type 2. *Veterinary Journal*, **187**, 23–32.
- Gregg MB (1975) Recent outbreaks of lymphocytic choriomeningitis in the United States of America. *Bulletin of the World Health Organization*, **52**, 549–553.
- Gritsun TS, Lashkevich VA, Gould EA (2003) Tick-borne encephalitis. *Antiviral Research*, **57**, 129–146.
- Haenen O, Hedrick R (2006) Koi herpesvirus workshop. *Bulletin of the European Association of Fish Pathologists*, **26**, 26–37.
- Haenen OLM, Davidse A (1993) Comparative pathogenicity of two strains of pike fry rhabdovirus and spring viremia of carp virus for young roach, common carp,

- grass carp and rainbow trout. *Diseases of Aquatic Organisms*, **15**, 87–92.
- Haenen OLM, Way K, Bergmann SM, Ariel E (2004) The emergence of koi herpesvirus and its significance to European aquaculture. *Bulletin of the European Association of Fish Pathologists*, **24**, 293–307.
- Harder TC, Vahlenkamp TW (2010) Influenza virus infections in dogs and cats. *Veterinary Immunology and Immunopathology*, **134**, 54–60.
- Hedrick RP, Gilad O, Yun S, Spangenberg JV, Marty GD *et al.* (2000) A herpesvirus associated with mass mortality of juvenile and adult koi, a strain of common carp. *Journal of Aquatic Animal Health*, **12**, 44–57.
- Hegde A, Teh HC, Lam TJ, Sin YM (2003) Nodavirus infection in freshwater ornamental fish, guppy, *Poecilia reticulata* – comparative characterization and pathogenicity studies. *Archives of Virology*, **148**, 575–586.
- Heyman P, Vaheri A, Lundkvist A, Avsic-Zupanc T (2009) Hantavirus infections in Europe: from virus carriers to a major public-health problem. *Expert Review of Anti-Infective Therapy*, **7**, 205–217.
- Hossain M, Song JY, Kitamura SI, Jung SJ, Oh MJ (2008) Phylogenetic analysis of lymphocystis disease virus from tropical ornamental fish species based on a major capsid protein gene. *Journal of Fish Diseases*, **31**, 473–479.
- Hubalek Z (2008) Mosquito-borne viruses in Europe. *Parasitology Research*, **103** (Suppl. 1), S29–S43.
- Hubalek Z, Halouzka J (1999) West Nile fever – a re-emerging mosquito-borne viral disease in Europe. *Emerging Infectious Diseases*, **5**, 643–650.
- Hubalek Z, Rudolf I (2012) Tick-borne viruses in Europe. *Parasitology Research*, **111**, 9–36.
- Hubalek Z, Zeman P, Halouzka J, Juricova Z, Stovickova E *et al.* (2005) Mosquito-borne viruses, Czech Republic, 2002. *Emerging Infectious Diseases*, **11**, 116–118.
- Jeong JB, Kim HY, Jun LJ, Lyu AH, Park NG *et al.* (2008) Outbreaks and risks of infectious spleen and kidney necrosis virus disease in freshwater ornamental fishes. *Diseases of Aquatic Organisms*, **78**, 209–215.
- Johne R, Wittig W, Fernandez-de-Luco D, Hofle U, Muller H (2006) Characterization of two novel polyomaviruses of birds by using multiply primed rolling-circle amplification of their genomes. *Journal of Virology*, **80**, 3523–3531.
- Kamar N, Bendall R, Legrand-Abravanel F, Xia NS, Ijaz S *et al.* (2012) Hepatitis E. *Lancet*, **379**, 2477–2488.
- Kuno A, Ido K, Isoda N, Satoh Y, Ono K *et al.* (2003) Sporadic acute hepatitis E of a 47-year-old man whose pet cat was positive for antibody to hepatitis E virus. *Hepatology Research*, **26**, 237–242.
- Le Guyader FS, Le Saux JC, Ambert-Balay K, Krol J, Serais O *et al.* (2008) Aichi virus, norovirus, astrovirus, enterovirus, and rotavirus involved in clinical cases from a French oyster-related gastroenteritis outbreak. *Journal of Clinical Microbiology*, **46**, 4011–4017.
- Lewis HC, Wichmann O, Duizer E (2010) Transmission routes and risk factors for autochthonous hepatitis E virus infection in Europe: a systematic review. *Epidemiology and Infection*, **138**, 145–166.
- Li L, Kapoor A, Slikas B, Bamidele OS, Wang C *et al.* (2010) Multiple diverse circoviruses infect farm animals and are commonly found in human and chimpanzee feces. *Journal of Virology*, **84**, 1674–1682.
- Li L, Pesavento PA, Shan T, Leutenegger CM, Wang C *et al.* (2011) Viruses in diarrhoeic dogs include novel kobuviruses and sapoviruses. *Journal of General Virology*, **92**, 2534–2541.
- Lin F, Kitching RP (2000) Swine vesicular disease: an overview. *Veterinary Journal*, **160**, 192–201.
- Lindberg A, Brownlie J, Gunn GJ, Houe H, Moennig V *et al.* (2006) The control of bovine viral diarrhoea virus in Europe: today and in the future. *Revue Scientifique et Technique de l'OIE*, **25**, 961–979.
- Luby SP, Gurley ES (2012) Epidemiology of henipavirus disease in humans. *Current Topics in Microbiology and Immunology*, **359**, 25–40.
- Mackenzie JS, Gubler DJ, Petersen LR (2004) Emerging flaviviruses: the spread and resurgence of Japanese encephalitis, West Nile and dengue viruses. *Nature Medicine*, **10**, S98–S109.
- Mahgoub HA, Bailey M, Kaiser P (2012) An overview of infectious bursal disease. *Archives of Virology*, **157**, 2047–2057.
- Marshall ES, Carpenter TE, Thurmond MC (2007) Results of a survey of owners of miniature swine to characterize husbandry practices affecting risks of foreign animal disease. *Journal of the American Veterinary Medical Association*, **230**, 702–707.
- Mazelet L, Dietrich J, Rolland JL (2011) New RT-qPCR assay for viral nervous necrosis virus detection in sea bass, *Dicentrarchus labrax* (L.): application and limits for hatcheries sanitary control. *Fish and Shellfish Immunology*, **30**, 27–32.
- McFerran JB, Smyth JA (2000) Avian adenoviruses. *Revue Scientifique et Technique de l'OIE*, **19**, 589–601.
- McLoughlin MF, Graham DA (2007) Alphavirus infections in salmonids – a review. *Journal of Fish Diseases*, **30**, 511–531.
- Mellor PS, Wittmann EJ (2002) Bluetongue virus in the Mediterranean Basin 1998–2001. *Veterinary Journal*, **164**, 20–37.
- Meng XJ (2010) Hepatitis E virus: animal reservoirs and zoonotic risk. *Veterinary Microbiology*, **140**, 256–265.
- Midgley SE, Banyai K, Buesa J, Halaihel N, Hjulager CK *et al.* (2012) Diversity and zoonotic potential of rotaviruses in swine and cattle across Europe. *Veterinary Microbiology*, **156**, 238–245.
- Mladenova Z, Papp H, Lengyel G, Kisfali P, Steyer A *et al.* (2012) Detection of rare reassortant G5P[6] rotavirus, Bulgaria. *Infection, Genetics and Evolution*, **12**, 1676–1684.
- Moennig V, Floegel-Niesmann G, Greiser-Wilke I (2003) Clinical signs and epidemiology of classical swine fever: a review of new knowledge. *Veterinary Journal*, **165**, 11–20.

- Morgan N, Prakash A (2006) International livestock markets and the impact of animal disease. *Revue Scientifique et Technique de l'OIE*, **25**, 517–528.
- Muller T, Hahn EC, Tottewitz F, Kramer M, Klupp BG *et al.* (2011) Pseudorabies virus in wild swine: a global perspective. *Archives of Virology*, **156**, 1691–1705.
- Munro ES, Midtlyng PJ (2011) Infectious pancreatic necrosis and associated aquatic birnaviruses. In: *Fish Diseases and Disorders*, PTK Woo, DW Bruno, Eds., CAB International, Wallingford, pp. 1–65.
- Myers KP, Olsen CW, Gray GC (2007) Cases of swine influenza in humans: a review of the literature. *Clinical Infectious Diseases*, **44**, 1084–1088.
- Nylund A, Watanabe K, Nylund S, Karlsten M, Saether PA *et al.* (2008) Morphogenesis of salmonid gill poxvirus associated with proliferative gill disease in farmed Atlantic salmon (*Salmo salar*) in Norway. *Archives of Virology*, **153**, 1299–1309.
- Olsen CW (2000) DNA vaccination against influenza viruses: a review with emphasis on equine and swine influenza. *Veterinary Microbiology*, **74**, 149–164.
- Olsson GE, Leirs H, Henttonen H (2010) Hantaviruses and their hosts in Europe: reservoirs here and there, but not everywhere? *Vector Borne and Zoonotic Diseases*, **10**, 549–561.
- Osterrieder N, Van de Walle GR (2010) Pathogenic potential of equine alphaherpesviruses: the importance of the mononuclear cell compartment in disease outcome. *Veterinary Microbiology*, **143**, 21–28.
- Oura CA, El Harrak M (2011) Midge-transmitted blue-tongue in domestic dogs. *Epidemiology and Infection*, **139**, 1396–1400.
- Palacios G, Lovoll M, Tengs T, Hornig M, Hutchison S *et al.* (2010) Heart and skeletal muscle inflammation of farmed salmon is associated with infection with a novel reovirus. *PLoS ONE*, **5**, e11487.
- Papa A (2012) Dobrava-Belgrade virus: phylogeny, epidemiology, disease. *Antiviral Research*, **95**, 104–117.
- Paperna I, Vilenkin M, de Matos AP (2001) Iridovirus infections in farm-reared tropical ornamental fish. *Diseases of Aquatic Organisms*, **48**, 17–25.
- Parrish CR (1999) Host range relationships and the evolution of canine parvovirus. *Veterinary Microbiology*, **69**, 29–40.
- Pavio N, Meng XJ, Renou C (2010) Zoonotic hepatitis E: animal reservoirs and emerging risks. *Veterinary Research*, **41**, 46.
- Peiris JSM (2009) Avian influenza viruses in humans. *Revue Scientifique et Technique de l'OIE*, **28**, 161–173.
- Perelberg A, Ilouze M, Kotler M, Steinitz M (2008) Antibody response and resistance of *Cyprinus carpio* immunized with cyprinid herpes virus 3 (CyHV-3). *Vaccine*, **26**, 3750–3756.
- Pfeffer M, Dobler G (2011) Tick-borne encephalitis virus in dogs – is this an issue? *Parasites and Vectors*, **4**, 59.
- Pfeffer M, Pflieger S, von BD, Kaaden OR, Meyer H (2002) Retrospective investigation of feline cowpox in Germany. *Veterinary Record*, **150**, 50–51.
- Phan TG, Kapusinszky B, Wang C, Rose RK, Lipton HL *et al.* (2011) The fecal viral flora of wild rodents. *PLoS Pathogens*, **7**, e1002218.
- Raj VS, Osterhaus AD, Fouchier RA, Haagmans BL (2014) MERS: emergence of a novel human coronavirus. *Current Opinion in Virology*, **5**, 58–62.
- Reed KD, Melski JW, Graham MB, Regnery RL, Sotir MJ *et al.* (2004) The detection of monkeypox in humans in the Western Hemisphere. *New England Journal of Medicine*, **350**, 342–350.
- Renou C, Cadranet JF, Bourliere M, Halfon P, Ouzan D *et al.* (2007) Possible zoonotic transmission of hepatitis E from pet pig to its owner. *Emerging Infectious Diseases*, **13**, 1094–1096.
- Reperant LA, Kuiken T, Osterhaus AD (2012) Adaptive pathways of zoonotic influenza viruses: from exposure to establishment in humans. *Vaccine*, **30**, 4419–4434.
- Reuter G, Boros A, Pankovics P (2011) Kobuviruses – a comprehensive review. *Reviews in Medical Virology*, **21**, 32–41.
- Rimstad E, Dale OB, Dannevig BH, Falk K (2011) Infectious salmon anaemia. In: *Fish Diseases and Disorders*, Vol. 3, PTK Woo, DW Bruno, Eds., CAB International, Wallingford.
- Rousseau MC, Saron MF, Brouqui P, Bourgeade A (1997) Lymphocytic choriomeningitis virus in southern France: four case reports and a review of the literature. *European Journal of Epidemiology*, **13**, 817–823.
- Rupprecht CE, Stöhr K, Meredith C (2001) Rabies. In: *Infectious Diseases of Wild Mammals*, ES Williams, Ed., Iowa State University Press, Ames, pp. 3–36.
- Russell GC, Stewart JP, Haig DM (2009) Malignant catarrhal fever: a review. *Veterinary Journal*, **179**, 324–335.
- Rutjes SA, Lodder-Verschuur F, Lodder WJ, van der Giessen J, Reesink H *et al.* (2010) Seroprevalence and molecular detection of hepatitis E virus in wild boar and red deer in The Netherlands. *Journal of Virological Methods*, **168**, 197–206.
- Santos N, Hoshino Y (2005) Global distribution of rotavirus serotypes/genotypes and its implication for the development and implementation of an effective rotavirus vaccine. *Reviews in Medical Virology*, **15**, 29–56.
- Schat KA (2009) Chicken anemia virus. *Current Topics in Microbiology and Immunology*, **331**, 151–183.
- Schielke A, Sachs K, Lierz M, Appel B, Jansen A *et al.* (2009) Detection of hepatitis E virus in wild boars of rural and urban regions in Germany and whole genome characterization of an endemic strain. *Virology Journal*, **6**, 58.
- Schmitt B (2002) Vesicular stomatitis. *Veterinary Clinics of North America: Food Animal Practice*, **18**, 453–459.
- Schrenzel MD, Tucker TA, Donovan TA, Busch MD, Wise AG *et al.* (2008) New hosts for equine herpesvirus 1. *Emerging Infectious Diseases*, **14**, 1616–1619.
- Shepherd AJ, Swanepoel R, Shepherd SP, McGillivray GM, Searle LA (1987) Antibody to Crimean–Congo hemorrhagic fever virus in wild mammals

- from southern Africa. *American Journal of Tropical Medicine and Hygiene*, **36**, 133–142.
- Sipos W, Schmoll F, Stumpf I (2007) Minipigs and potbellied pigs as pets in the veterinary practice – a retrospective study. *Journal of Veterinary Medicine. Series A, Physiology, Pathology and Clinical Medicine*, **54**, 504–511.
- Sjaak de Wit JJ, Cook JK, van der Heijden HM (2011) Infectious bronchitis virus variants: a review of the history, current situation and control measures. *Avian Pathology*, **40**, 223–235.
- Skall HF, Olesen NJ, Møllergaard S (2005) Viral haemorrhagic septicaemia virus in marine fish and its implications for fish farming – a review. *Journal of Fish Diseases*, **28**, 509–529.
- Smail DA, Snow M (2011) Viral haemorrhagic septicaemia. In: *Fish Diseases and Disorders*, PTK Woo, DW Bruno, Eds., CAB International, Wallingford, pp. 166–244.
- Swenson SL, Koster LG, Jenkins-Moore M, Killian ML, DeBess EE *et al.* (2010) Natural cases of 2009 pandemic H1N1 influenza A virus in pet ferrets. *Journal of Veterinary Diagnostic Investigation*, **22**, 784–788.
- Tack DM (2011) Zoonotic poxviruses associated with companion animals. *Animals*, **1**, 377–395.
- Tack DM, Reynolds MG (2011) Zoonotic poxviruses associated with companion animals. *Animals (Basel)*, **1**, 377–395.
- Thiry E, Zicola A, Addie D, Egberink H, Hartmann K *et al.* (2007) Highly pathogenic avian influenza H5N1 virus in cats and other carnivores. *Veterinary Microbiology*, **122**, 25–31.
- Tischer BK, Osterrieder N (2010) Herpesviruses – a zoonotic threat? *Veterinary Microbiology*, **140**, 266–270.
- Tuppurainen ES, Oura CA (2012) Review: lumpy skin disease: an emerging threat to Europe, the Middle East and Asia. *Transboundary and Emerging Diseases*, **59**, 40–48.
- USDA (1994) *Foot-and-mouth Disease: Sources of Outbreaks and Hazard Categorization of Modes of Virus Transmission*. US Department of Agriculture, Animal and Plant Health Inspection Service, Veterinary Services, Fort Collins, p. 38.
- Valarcher JF, Leforban Y, Rweyemamu M, Roeder PL, Gerbier G *et al.* (2008) Incursions of foot-and-mouth disease virus into Europe between 1985 and 2006. *Transboundary and Emerging Diseases*, **55**, 14–34.
- van der Poel WH, Lina PH, Kramps JA (2006) Public health awareness of emerging zoonotic viruses of bats: a European perspective. *Vector Borne and Zoonotic Diseases*, **6**, 315–324.
- Vapalahti O, Mustonen J, Lundkvist A, Henttonen H, Plyusnin A *et al.* (2003) Hantavirus infections in Europe. *Lancet Infectious Diseases*, **3**, 653–661.
- Vasickova P, Psikal I, Kralik P, Widen F, Hubalek Z *et al.* (2007) Hepatitis E virus: a review. *Veterinari Medicina*, **52**, 365–384.
- von Bomhard W, Mauldin EA, Breuer W, Pfliegerhaer S, Nitsche A (2011) Localized cowpox infection in a 5-month-old rottweiler. *Veterinary Dermatology*, **22**, 111–114.
- Walz PH, Grooms DL, Passler T, Ridpath JF, Tremblay R *et al.* (2010) Control of bovine viral diarrhoea virus in ruminants. *Journal of Veterinary Internal Medicine*, **24**, 476–486.
- Way K (2009) Viruses in coldwater ornamental fish. *Microbiology Today*, **36**, 200–203.
- Welchman Dde B, Cox WJ, Gough RE, Wood AM, Smyth VJ *et al.* (2009) Avian encephalomyelitis virus in reared pheasants: a case study. *Avian Pathology*, **38**, 251–256.
- Whitehouse CA (2004) Crimean–Congo hemorrhagic fever. *Antiviral Research*, **64**, 145–160.
- Whittington RJ, Becker JA, Dennis MM (2010) Iridovirus infections in finfish – critical review with emphasis on ranaviruses. *Journal of Fish Diseases*, **33**, 95–122.
- Wolfs TF, Wagenaar JA, Niesters HG, Osterhaus AD (2002) Rat-to-human transmission of cowpox infection. *Emerging Infectious Diseases*, **8**, 1495–1496.
- Wood EN (1979) Transmissible gastroenteritis and epidemic diarrhoea of pigs. *British Veterinary Journal*, **135**, 305–314.
- Woolhouse M, Scott F, Hudson Z, Howey R, Chase-Topping M (2012) Human viruses: discovery and emergence. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, **367**, 2864–2871.
- Zacks MA, Paessler S (2010) Encephalitic alphaviruses. *Veterinary Microbiology*, **140**, 281–286.
- Zeller HG, Schuffenecker I (2004) West Nile virus: an overview of its spread in Europe and the Mediterranean basin in contrast to its spread in the Americas. *European Journal of Clinical Microbiology and Infectious Diseases*, **23**, 147–156.
- Zhang Y, Zhang H, Dong X, Yuan J, Yang X *et al.* (2010) Hantavirus outbreak associated with laboratory rats in Yunnan, China. *Infection, Genetics and Evolution*, **10**, 638–644.

[Received, January 19th, 2016]
 [Accepted, July 7th, 2016]