

- attenuated vaccine for chimpanzees. *J Infect Dis.* 1996;173:592–7. <http://dx.doi.org/10.1093/infdis/173.3.592>
7. Drexler JF, Corman VM, Lukashev AN, Brand JMA, van den Gmyl AP, Brünink S, et al. Evolutionary origins of hepatitis A virus in small mammals. *Proc Natl Acad Sci U S A.* 2015;112:15190–5. <http://dx.doi.org/10.1073/pnas.1516992112>
 8. Drewe JA, O’Riain MJ, Beamish E, Currie H, Parsons S. Survey of infections transmissible between baboons and humans, Cape Town, South Africa. *Emerg Infect Dis.* 2012;18:298–301. <http://dx.doi.org/10.3201/eid1802.111309>
 9. Goldberg TL, Gillespie TR, Rwego IB, Estoff EL, Chapman CA. Forest fragmentation as cause of bacterial transmission among nonhuman primates, humans, and livestock, Uganda. *Emerg Infect Dis.* 2008;14:1375–82. <http://dx.doi.org/10.3201/eid1409.071196>
 10. Naughton-Treves L. Predicting patterns of crop damage by wildlife around Kibale National Park, Uganda. *Conserv Biol.* 1998;12:156–68. <http://dx.doi.org/10.1046/j.1523-1739.1998.96346.x>

Address for correspondence: Tony L. Goldberg, Department of Pathobiological Sciences, University of Wisconsin–Madison, 1656 Linden Dr, Madison, WI 53706, USA; email: tgoldberg@vetmed.wisc.edu

Porcine Bocavirus Infection Associated with Encephalomyelitis in a Pig, Germany¹

Vanessa M. Pfankuche, Rogier Bodewes, Kerstin Hahn, Christina Puff, Andreas Beineke, André Habierski, Albert D.M.E. Osterhaus, Wolfgang Baumgärtner

Author affiliations: University of Veterinary Medicine, Hannover, Germany (V.M. Pfankuche, K. Hahn, C. Puff, A. Beineke, A. Habierski, A.D.M.E. Osterhaus, W. Baumgärtner); Center for Systems Neuroscience, Hannover (V.M. Pfankuche, K. Hahn, A. Beineke, W. Baumgärtner); The Erasmus University Medical Center, Rotterdam, the Netherlands (R. Bodewes); Utrecht University, Utrecht, the Netherlands (R. Bodewes)

DOI: <http://dx.doi.org/10.3201/eid2207.152049>

To the Editor: In 2013, a 6-week-old female piglet kept in a flatdeck cage had coughing, growth retardation, and diarrhea and was taken to a local veterinarian in Hannover, Germany; the piglet was euthanized. After necropsy at the University of Veterinary Medicine in Hannover, histologic investigation found interstitial pneumonia; a mild, multifocal, lymphohistiocytic panencephalitis that affected the cerebrum and cerebellum, including brain stem and

medulla oblongata; and a mild, multifocal, lymphohistiocytic panmyelitis. Results from screening for typical neurotropic viruses (classical swine fever virus, suid herpesvirus 1, rabies virus, teschovirus, porcine enterovirus 8, 9, and 10) were negative; *Mycoplasma hyorhinis* was detected by multiplex PCR (Institute of Virology, University of Veterinary Medicine Hannover) within the lung and pulmonary lymph nodes. Cerebral tissue from the pig was processed for viral metagenomics by random RNA and DNA virus screening and next-generation sequencing (NGS) with the 454 sequencing platform (GS Junior; Roche, Basel, Switzerland), as described (1), and 21,359 reads were obtained. Analysis by using blastn and blastx (2) showed 10 reads had $\geq 97\%$ nt identity with porcine bocavirus (PBoV) KU14. No other viral sequences were detected.



By using primers based on sequence data of the PBoV, partially overlapping PCR amplicons were obtained to confirm and extend the NGS data of the isolate, which was named PBoV S1142/13 (1; GenBank accession no. KU311698). A total of 2,176 nt of PBoV S1142/13 were obtained, consisting of the partial nucleoprotein (NP) 1 and the nearly complete viral protein (VP) 1 gene. By using MAFFT version 7 (<http://mafft.cbrc.jp/alignment/server/>), we aligned the nearly complete VP1 gene of PBoV S1142/13 with various closely related members of the genus *Bocaparvovirus* and built a maximum-likelihood tree by using the general time reversible plus invariable sites plus gamma distribution method, as determined by jModelTest 2.0 (3) and default parameters in MEGA6.06 (4). Results confirmed that PBoV S1142/13 was most closely related to PBoV KU14 (Figure, panel A). The partial genome of PBoV S1142/13 differed at 8 nt positions from PBoV KU14, resulting in 99.6% nt identity. Of these nucleotide differences, 4 resulted in an amino acid difference, including position 2733 (T→C on the basis of PBoV KU14 as a reference genome), which is part of the NP1 stopcodon of PBoV KU14. These results indicate that the stopcodon was located 39 nt farther downstream than for PBoV KU14. The other 3 aa differences were present in the VP1 protein; each of these differences was within the same group of amino acids as those detected in PBoV KU14.

For further substantiation of a potential cause-effect relationship of histologic (Figure, panel B) and NGS results, we performed fluorescent in situ hybridization (FISH) on formalin-fixed, paraffin-embedded central nervous system (CNS) sections of the diseased animal and of a control pig with no CNS lesions. We used an RNA probe specific for the obtained NP1 and VP1 sequences covering 1,153 nt (Affymetrix, Santa Clara, CA, USA) according to the manufacturer’s protocol, with minor variations (ViewRNA ISH Tissue 1-Plex Assay Kit and ViewRNA Chromogenic Signal Amplification Kit, Affymetrix). A probe specific for porcine ubiquitin (*Sus scrofa* ubiquitin

¹Preliminary results from this study were presented at the 3rd International One Health Congress, March 15–18, 2015, Amsterdam, the Netherlands; and at the Conference of the German Veterinary Medical Association, March 8–10, 2015, Fulda, Germany.

Figure. Phylogenetic analysis and staining of porcine bocavirus (PBoV) from the spinal cord of a diseased pig, Hannover, Germany.

A) Phylogenetic relationship of PBoV isolate S1142/13 (bold) with other bocaviruses. The nucleotide sequence of the nearly complete viral protein 1 of PBoV S1142/13 was aligned with other members of the genus *Bocaparvovirus*, and a maximum-likelihood phylogenetic tree was prepared by using the general time reversible plus invariable sites plus gamma distribution model and 500 bootstrap replicates. Only bootstrap values >70 are shown. Scale bar indicates nucleotide substitutions per site.



B) Spinal cord of the diseased pig showing perivascular accentuated mild, focal, nonsuppurative inflammation (arrow). Hematoxylin and eosin stain. Scale bar indicates 100 μ m. C) Intracytoplasmic and intranuclear PBoV-specific positive signals in neurons of the spinal cord of the diseased pig (arrowheads), determined by using fluorescent in situ hybridization (Fast Red; ViewRNA Chromogenic Signal Amplification Kit; Affymetrix, Santa Clara, CA, USA). Phase contrast and fluorescent microscopy. Scale bar indicates 100 μ m.

C; GenBank accession no. XM_005657305; nt 2–890) served as a positive control.

The spinal cord of the diseased pig showed diffuse intracytoplasmic and intranuclear PBoV-specific signals within scattered neurons adjacent to the histologically detected inflammatory lesions (Figure, panel C). The negative control and the nonprobe incubation lacked PBoV-specific signals. The porcine ubiquitin probe provided a strong intracellular and extracellular staining within the CNS of both pigs.

PBoV (genus *Bocaparvovirus*, family *Parvoviridae*) was first described in 2009 as porcine boca-like virus in pigs in Sweden with postweaning multisystemic wasting syndrome (5). PBoV is usually involved in respiratory and intestinal diseases in pigs (5) but has not been detected in the CNS. In the pig in our study, the lack of detection of other viral sequences by using NGS indicates the potential role of PBoV as a pathogen that triggers encephalomyelitis. FISH substantiated the NGS results and revealed neuronal intracytoplasmic and intranuclear PBoV-specific signals adjacent to the lesion, indicating intraneuronal transcription and replication (6). Nevertheless, a potential synergistic effect of *M. hyorhinis* on the PBoV pathogenesis cannot be ruled out. Similarly, co-infection of *M. hyorhinis* and porcine circovirus type 2 has been associated with enhanced inflammatory lesions in the lungs of pigs (7).

The CNS tropism of PBoV S1142/13 could result from various factors, including specific amino acid changes that

enable the virus to pass the blood–brain barrier and infect neurons. Additional studies are necessary to elucidate a possible role of the amino acid differences between PBoV S1142/13 and PBoV KU14 in the tropism of these viruses.

Human bocavirus has recently been found in the cerebrospinal fluid of patients having encephalitis (8), and related human parvovirus 4 (9) and human parvovirus B19 (10) have been reported in human encephalitis. The correlation of PBoV-specific signals by using FISH for histologic detection of encephalomyelitis assigns PBoV a potential role in provoking CNS lesions. PBoV should be considered as a cause of encephalomyelitis but needs further investigation.

Acknowledgments

The authors thank the Lower Saxony State Office for Consumer Protection and Food Safety, the Institute of Virology at the University of Veterinary Medicine Hannover, and the Friedrich-Loeffler-Institute in Jena for performing the routine diagnostic analyses of common porcine pathogens. The authors also thank Danuta Waschke, Kerstin Rohn, Bettina Buck, Caroline Schütz, and Kerstin Schöne for excellent technical assistance.

This study was in part supported by the Niedersachsen-Research Network on Neuroinfectiology of the Ministry of Science and Culture of Lower Saxony and by the COMPARE project and received funding from the European Union's Horizon 2020 research and innovation program COMPARE (grant agreement no. 643476). V.M.P. received a scholarship from the Akademie für Tiergesundheit e.V. in Bonn, Germany.

References

1. van Leeuwen M, Williams MM, Koraka P, Simon JH, Smits SL, Osterhaus AD. Human picobirnaviruses identified by molecular screening of diarrhea samples. *J Clin Microbiol*. 2010;48:1787–94. <http://dx.doi.org/10.1128/JCM.02452-09>
2. Schürch AC, Schipper D, Bijl MA, Dau J, Beckmen KB, Schapendonk CM, et al. Metagenomic survey for viruses in Western Arctic caribou, Alaska, through iterative assembly of taxonomic units. *PLoS One*. 2014;9:e105227. <http://dx.doi.org/10.1371/journal.pone.0105227>
3. Darriba D, Taboada GL, Doallo R, Posada D. jModelTest 2: more models, new heuristics and parallel computing. *Nat Methods*. 2012;9:772. <http://dx.doi.org/10.1038/nmeth.2109>
4. Tamura K, Stecher G, Peterson D, Filipinski A, Kumar S. MEGA6: Molecular evolutionary genetics analysis version 6.0. *Mol Biol Evol*. 2013;30:2725–9. <http://dx.doi.org/10.1093/molbev/mst197>
5. Zhou F, Sun H, Wang Y. Porcine bocavirus: achievements in the past five years. *Viruses*. 2014;6:4946–60. <http://dx.doi.org/10.3390/v6124946>
6. Bodewes R, Lapp S, Hahn K, Habierski A, Förster C, König M, et al. Novel canine bocavirus strain associated with severe enteritis in a dog litter. *Vet Microbiol*. 2014;174:1–8. <http://dx.doi.org/10.1016/j.vetmic.2014.08.025>
7. Chen D, Wei Y, Huang L, Wang Y, Sun J, Du W, et al. Synergistic pathogenicity in sequential coinfection with *Mycoplasma hyorhinis* and porcine circovirus type 2. *Vet Microbiol*. 2016;182:123–30. <http://dx.doi.org/10.1016/j.vetmic.2015.11.003>
8. Mori D, Ranawaka U, Yamada K, Rajindrajith S, Miya K, Perera HK, et al. Human bocavirus in patients with encephalitis, Sri Lanka, 2009–2010. *Emerg Infect Dis*. 2013;19:1859–62. <http://dx.doi.org/10.3201/eid1911.121548>
9. Benjamin LA, Lewthwaite P, Vasanthapuram R, Zhao G, Sharp C, Simmonds P, et al. Human parvovirus 4 as potential cause of encephalitis in children, India. *Emerg Infect Dis*. 2011;17:1484–7.
10. Barah F, Whiteside S, Batista S, Morris J. Neurological aspects of human parvovirus B19 infection: a systematic review. *Rev Med Virol*. 2014;24:154–68. <http://dx.doi.org/10.1002/rmv.1782>

Address for correspondence: Wolfgang Baumgärtner, Department of Pathology, University of Veterinary Medicine, Bünteweg 17 D-30559 Hannover, Germany; email: wolfgang.baumgaertner@tiho-hannover.de

Pegivirus Infection in Domestic Pigs, Germany

Christine Baechlein,¹ Adam Grundhoff,¹
Nicole Fischer, Malik Alawi, Doris Hoeltig,
Karl-Heinz Waldmann, Paul Becher

Author affiliations: University of Veterinary Medicine Hannover, Hannover, Germany (C. Baechlein, D. Hoeltig, K.-H. Waldmann, P. Becher); German Center for Infection Research Partner Site Hannover–Braunschweig, Hannover (C. Baechlein, P. Becher); German Center for Infection Research Partner Site Hamburg–Lübeck–Borstel, Hamburg (N. Fischer, A. Grundhoff); Heinrich Pette Institute, Hamburg, Germany (A. Grundhoff, M. Alawi); University Medical Center Hamburg–Eppendorf, Hamburg (N. Fischer, M. Alawi)

DOI: <http://dx.doi.org/10.3201/eid2207.160024>

¹These authors contributed equally to this article.

To the Editor: The family *Flaviviridae* includes many human and animal virus pathogens. Recently, in addition to the genera *Flavivirus*, *Hepacivirus*, and *Pestivirus*, a fourth genus, *Pegivirus*, has been identified (1). In addition to human pegiviruses, a range of phylogenetic, highly divergent pegiviral sequences have been identified in various animal species, including primates, bats, rodents, and horses (2). We report the detection of a porcine pegivirus (PPgV) in serum samples from pigs.

Initially, we investigated pooled serum samples by using high-throughput sequencing methods and isolated RNA from individual porcine serum samples by using the QI-Amp Viral RNA Mini Kit (QIAGEN, Hilden, Germany). We prepared libraries compatible with Illumina (San Diego, CA, USA) sequencing from pooled samples and individual serum samples by using the ScriptSeq version 2 RNA-Seq Library Preparation Kit (Epicenter, Madison, WI, USA) and sequenced them by using a HiSeq 2500 (2 × 150 cycles paired-end; Illumina) for pooled samples and MiSeq (2 × 250 cycles paired-end; Illumina) for individual samples (3).

We conducted quantitative reverse transcription PCR (RT-PCR) by using a Quantitect-SYBR Green Assay (QIAGEN) and primers PPgV_fwd: 5'-CTGTCTATGCTGGTCACTGGA-3' and PPgV_rev: 5'-GCCATAGAACGGGAAGTCGC-3'. By using high-throughput sequencing of the pooled serum sample library (23,167,090 reads), we identified 1 contig (4,582 bp) that had distant nucleotide sequence similarity to bat pegivirus (69% and 4% sequence coverage) and 2 contigs (2,683 bp and 665 bp) that had 73% sequence coverage, thereby covering 8% and 37% of the identified sequence. RT-PCR with primers designed on basis of recovered sequences identified the sample containing pegivirus sequences. Subsequent MiSeq analysis (7,085,595 reads) of an RNA library prepared from a sample from 1 animal identified 1 contig (9,145 nt) with sequence similarity to pegivirus sequences.

We performed 3' end completion of the viral genome by rapid amplification of cDNA ends and identified the entire open reading frame of PPgV_903 encoding 2,972 aa (GenBank accession no. KU351669). Analysis of the pegivirus 5' untranslated region identified a highly structured internal ribosome entry site motif (online Technical Appendix, <http://wwwnc.cdc.gov/EID/article/22/7/16-0024-Techapp1.pdf>), which was similar in structure to previously described 5' untranslated region structures of other pegiviruses (4,5).

Pegiviruses do not encode a protein homologous to the capsid protein of other viruses of the family *Flaviviridae*, another common feature of pegiviruses (6). The presence of cleavage sites for cellular signal peptidases and viral proteases indicates that, similar to polyproteins of other pegiviruses and members of the genus *Hepacivirus*, the pegivirus polyprotein NH₂-E1-E2-Px-NS2-NS3-NS4A-NS4B-NS5A-NS5B-COOH (E [envelope], NS [nonstructural],