



The Detailed 3D Multi-Loop Aggregate/Rosette Chromatin Architecture and Functional Dynamic Organization of the Human and Mouse Genomes







Tobias A. Knoch

Biophysical Genomics & Erasmus Computing Grid

Erasmus Medical Center

TA.Knoch@taknoch.org








The Detailed 3D Multi-Loop Aggregate/ Chromatin Architecture and Functional Dynamics Human Genomes


**From Sequence to Morphology:
Towards a Holistic Understanding of Genomes!**

Dr. Tobias A. Knoch
Genomics & Erasmus Computing Grid
Erasmus Medical Center
TA.Knoch@taknoch.org



Dynamic and Hierarchical Genome Organization

The different organization levels of genomes bridge several orders of magnitude concerning space and time. How all of these organization levels connect to processes like gene regulation, replication, embryogeneses, or cancer development is still unclear?




WE KNOW THAT WE HARDLY KNOW ANYTHING

From the sequence to the morphology we just have begun to elucidate the organization of genomes and we also just have begun to understand that it is not only the detailed knowledge about one organizational level but beyond the holistic entity of the whole cell nucleus or genome which makes genomes function!



Approaching the Three-Dimensional Organization of the Human Genome

Structural-, Scaling- and Dynamic-Properties
in the Simulation of
Interphase Chromosomes and Cell Nuclei
Long-Range Correlations in Complete Genomes
In Vivo Quantification of the Chromatin Distribution
Construct Conversions in Simultaneous Co-Transfections



Tobias A. Knoch



2002
ISBN 3-00-009959-X

Nucleus:

- > Nuclear organization: chromosome arrangement, morphology?
- > Nuclear Code: information content, regulation, variability?

Chromosome:

- > Chromosome organization: loops, loop aggregates, extension?
- > Chromosome code: information content, regulation, variability?

Chromatin:


- > Chromatin fiber organization: prevalence, variation, dynamics?
- > Chromatin code: coding, regulation, modification?

Nucleosome:

- > Nucleosome organization: tail position, mobility, modification?
- > Histon code: coding, regulation, modification?

DNA Sequence:

- > DNA local structure: bending, melting, stability, modification?
- > General sequence organization: coding, regulating, and the rest?




Chromatin Conformation and Higher-Order Topologies


It becomes increasingly clearer, that the chromatin conformation is a random organization of nucleosomes, which depending on external or modification conditions has different condensation degrees, with a prevalence for the 30nm fiber with ~6nucleosomes per 11nm. This seems to make loops which further cluster to form aggregates more or less rosette-like which then constitute the chromosome.




A-C: Voet & Voigt; D: Reznik *et al.*



Courtesy P. Fransz, Amsterdam



A: B. Aramova *et al.*; C: Salganik *et al.*; D-G: Frznik *et al.*; G-K: Tsvetkov & Pavlov





Integral Models of Cell Nuclear Organization I

Already Rabl and Boveri were aware of the obvious fact that the organization of genomes has to be consistent from the sequence level to the morphology of the whole cell nucleus. Although they might be different in detail their common seem is recursive folding and clustering thereof with variation/ modification and dynamics accounting for different nuclear states and function.





Rabl & Boveri





A: Bloom & Fawcett
B: Alberts *et al.*
C: Paulson & Laemmli

A, B: Pienta & Coffey; C: Alberts *et al.*; D: Belmont & Bruce





Integral Models of Cell Nuclear Organization II

The biggest advantage of integral models is the again obvious and simple fact, that they allow the validation from the consistency of different levels of organization from the other levels. Thus, e.g. the so called Interchromosomal Domain Model can be ruled out by simple voluminous thought...






Random-Walk/Giant-Loop Multi-Loop-Subcompartment Model





A: courtesy K. Richter; B: courtesy K. Greulich-Bode

Selective Chromosome Interaction Capture (T2C)

T2C is a novel selective high-resolution high-throughput chromosome interaction capture, in which the relation between, region size, resolution, interaction frequency range, and sequencing depth can be designed towards the goal of the experiment. T2C reaches the limit of the “genomic” uncertainty principle and statistical mechanics.








We Reached the Level of the Genomic Uncertainty Principle!
The Conformation of Chromatin and the 3D Architecture
can be derived with Tremendous Resolution !

Stable Consensus Architecture of Genomes


Due to the high signal-to-noise ratio of T2C reaching 5-6 orders of magnitude interaction maps reveal definitely an extremely high degree of similarity between different species, cell types, or functional states, thus functional differences are variation of a stable theme persisting through the cell cycle






HB2



TEV



HRV




Stable Consensus Architecture of Genomes


Due to the high signal-to-noise ratio of T2C reaching 5-6 orders of magnitude interaction maps reveal definitely an extremely high degree of similarity between different species, cell types, or functional states, thus functional differences are variation of a stable theme persisting through the cell cycle



fetal brain



fetal liver





Simulation of Single Chromosomes

The 30 nm chromatin fiber is modeled as a polymer chain with stretching, bending, and excluded volume interactions. Monte Carlo and Brownian Dynamic methods lead to thermodynamical equilibrium configurations.


All models form chromosome territories with big voids and different chromatin morphologies. Experimental territory and subcompartment diameters agree best with an MLS model with 80 to 120 kbp loops and linkers.




Metaphase starting configuration with ideogram bands in red/green, linker in grey.




RW/GL model, loop size 5 Mbp, after ~80.000 MC and 1000 relaxing BD steps. Large loops intermingle freely and reach out of the chromosome territory, thus forming no distinct features like in MLS model.



MLS model, loop size 126kbp, linker size 126 kbp, after ~50.000 MC and 1000 relaxing BD steps. Here rosettes form subcompartments as separated organizational and dynamic entities.






RW/GL model, loop size 126 kbp, after ~80.000 MC and 1000 relaxing BD steps. Large loops intermingle freely thus forming no distinct features like in MLS model.



Simulation of Single Chromosomes

The 30 nm chromatin fiber is modeled as a polymer chain with stretching, bending, and excluded volume interactions. Monte Carlo and Brownian Dynamic methods lead to thermodynamical equilibrium configurations.


All models form chromosome territories with big voids and different chromatin morphologies. Experimental territory and subcompartment diameters agree best with an MLS model with 80 to 120 kbp loops and linkers.



RW/GL model, loop size ~80.000 MC and 1000 relaxing BD steps. Large loops intermingle freely and reach out of the chromosome territory, thus forming no distinct features like in MLS model.

120 kbp, after ~30.000 MC and 1000 relaxing BD steps. Here rosettes form subcompartments as separated organizational and dynamic entities.


model, loop size 126 kbp, after ~80.000 MC and 1000 relaxing BD steps. Large loops intermingle freely thus forming no distinct features like in MLS model.



Simulation of Whole Nuclei with all 46 Chromosomes

Starting with some metaphase arrangement of cylindrical chromosomes, interphase nuclei with a 30 nm fiber resolution and at thermodynamical equilibrium are created in 4 steps using simulated annealing and Brownian Dynamics methods with stretching, bending, excluded volume and a spherical boundary interactions.


The chromosome territory position depends on their metaphase position and is reasonably stable.



Simulation of Whole Nuclei with all 46 Chromosomes

Starting with some metaphase arrangement of cylindrical chromosomes, interphase nuclei with a 30 nm fiber resolution and at thermodynamical equilibrium are created in 4 steps using simulated annealing and Brownian Dynamics methods with stretching, bending, excluded volume and a spherical boundary interactions.



The chromosome territory position depends on their metaphase position and is reasonably stable.



Simulation of Whole Nuclei with all 46 Chromosomes

Starting with some metaphase arrangement of cylindrical chromosomes, interphase nuclei with a 30 nm fiber resolution and at thermodynamical equilibrium are created in 4 steps using simulated annealing and Brownian Dynamics methods with stretching, bending, excluded volume and a spherical boundary interactions.




The chromosome territory position depends on their metaphase position and is reasonably stable.



Fine Structure of Loop Aggregates/Rosettes




Depending on the resolution, the loops within a domain and their arrangement in loop aggregates/rosettes can be shown as well as the details of how the loops are organized at their base as well as their aggregated rosette core: parallel loop fibres exist at the loop base with ~6kbp and these form the core.

A collage of logos and diagrams. At the top left is the logo for 'Bundesministerium für Bildung und Forschung'. To its right is 'N7/O' with the text 'Nucleosome Organization and Chromatin Dynamics'. Further right is 'BBRC'. Below these are the 'European Commission' logo and 'Erasmus MC' logo. In the center is a circular diagram with 'ExGenSys' and 'Chromatin' written around it, and 'Structure - Function' at the bottom. To the right of this diagram is 'Erasmus University' and 'Utrecht University'. At the bottom are logos for 'University of Oxford', 'dkfz', 'Fachhochschule', and 'U'. There is also a logo for 'Erasmus MC' with the text 'Erasmus Research Center'.



Fine Structure of Loop Aggregates/Rosettes




Depending on the resolution, the loops within a domain and their arrangement in loop aggregates/rosettes can be shown as well as the details of how the loops are organized at their base as well as their aggregated rosette core: parallel loop fibres exist at the loop base with ~6kbp and these form the core.



Fine Structure of Loop Aggregates/Rosettes

Depending on the resolution, the loops within a domain and their arrangement in loop aggregates/rosettes can be shown as well as the details of how the loops are organized at their base as well as their aggregated rosette core: parallel loop fibres exist at the loop base with ~6kbp and these form the core.


A collage of logos from various funding agencies and research institutions, including BBSRC, NWO, European Commission, and several research centers like the Centre for Systems Biology and the University of Oxford.



Fine Structure of Loop Aggregates/Rosettes



Depending on the resolution, the loops within a domain and their arrangement in loop aggregates/rosettes are shown as well as the details of how the loops are organized at their base as well as their aggregate core: parallel loop fibres exist at the loop base with ~6kbp and these form the core

The Rosalind Franklin Images!
The 3D Architecture of Chromosomes can be Directly Derived Visually !!!








Simulated Interaction Maps

Simulated spatial distance maps as well as simulated interaction maps result in the representation of every parameter variation, and also exhibit the fine-structure describing the loop base as well as rosette core. Thus from the quasi-fibre to the entire chromosome the architecture can be understood in detail.






Simulated Interaction Maps

Simulated spatial distance maps as well as simulated interaction maps result in the representation of every parameter variation, and also exhibit the fine-structure describing the loop base as well as rosette core. Thus from the quasi-fibre to the entire chromosome the architecture can be understood in detail.







Variation of a Consensus Architecture Scheme

The difference between different cell types, functional states or even species is minor despite depending on the region. From this, the chromatin fibre conformation, loop position, and their association into loop aggregates/rosettes can be derived, simulated by polymer models and finally visualized.




Variation of a Consensus Architecture Scheme

The difference between different cell types, functional states or even species is minor despite depending on the region. From this, the chromatin fibre conformation, loop position, and their association into loop aggregates/rosettes can be derived, simulated by polymer models and finally visualized.



DNA Sequence Organization


Determination of the concentration fluctuation function $C(l)$ and its local slope the correlation coefficient $\delta(l)$ are an indication for the i) degree of long-rang scaling behaviour, ii) general multi-scaling, and iii) fine-structure features, which all are connected to all levels of genome organization and especially also the three-dimensional genome architecture.



$$C(l) = \sqrt{\langle (c_l - \bar{c}_L)^2 \rangle_s}$$






$$C(l) = \sqrt{\frac{1}{L-l+1} \sum_{s=1}^{L-l} \left(\frac{1}{l} \sum_{k=1}^l n - \frac{1}{L} \sum_{k=1}^L N \right)^2}$$

$$C(l) = \frac{1}{Ll} \sqrt{\frac{1}{L-l} \sum_{s=1}^{L-l} \left[\left(\sum_{k=1}^l Ln \right) - \left(\sum_{k=1}^L lN \right) \right]^2}$$






DNA Sequence Organization

Determination of the concentration fluctuation function $C(l)$ and its local slope the correlation coefficient $\delta(l)$ are an indication for the i) degree of long-rang scaling behaviour, ii) general multi-scaling, and iii) fine-structure features, which all are connected to all levels of genome organization and especially also the three-dimensional genome architecture.





DNA Sequence Organization

Determination of the concentration fluctuation function $C(l)$ and its local slope the correlation coefficient $\delta(l)$ are an indication for the i) degree of long-rang scaling behaviour, ii) general multi-scaling, and iii) fine-structure features, which all are connected to all levels of genome organization and especially also the three-dimensional genome architecture.






Scaling Analysis

Scaling analysis show again the entire bandwidth of architectural effects in an aggregated manner. Beyond, they show the scale bridging of the structures and the evolutionary holistic entanglement between the 3D architecture and the DNA sequence organization itself.






Scaling Analysis

Scaling analysis show again the entire bandwidth of architectural effects in an aggregated manner. Beyond, they show the scale bridging of the structures and the evolutionary holistic entanglement between the 3D architecture and the DNA sequence organization itself.






Scaling Analysis

Scaling analysis show again the entire bandwidth of architectural effects in an aggregated manner. Beyond, they show the scale bridging of the structures and the evolutionary holistic entanglement between the 3D architecture and the DNA sequence organization itself.




Scaling Analysis

Scaling analysis show again the entire bandwidth of architectural effects in an aggregated manner. Beyond, they show the scale bridging of the structures and the evolutionary holistic entanglement between the 3D architecture and the DNA sequence organization itself.




Scaling Analysis

Scaling analysis show again the entire bandwidth of architectural effects in an aggregated manner. Beyond they show the scale bridging of the structures and the evolutionary holistic entanglement between the 3D architecture and the DNA sequence organization itself.





The 3D Architecture and the DNA Sequence Organization are Evolutionary Entangled!



3D Architecture of the Prader-Willi Region

Fluorescence *in situ* hybridization with various protocols of small probes within the Prader-Willi region combined with spectral precision distance confocal laser scanning microscopy and comparison with large-scale computer simulations shows a Multi-Loop Subcompartment organization of the Prader-Willi region.





3D Architecture of the Prader-Willi Region



Fluorescence *in situ* hybridization with various protocols of small probes within the Prader-Willi region combined with spectral precision distance confocal laser scanning microscopy and comparison with large-scale computer simulations shows a Multi-Loop Subcompartment organization of the Prader-Willi region.



650 kbp
6205 nm
520



2160 nm
480 nm





3D Architecture of the Prader-Willi Region

Fluorescence *in situ* hybridization with various protocols of small probes within the Prader-Willi region combined with spectral precision distance confocal laser scanning microscopy and comparison with large-scale computer simulations shows a Multi-Loop Subcompartment organization of the Prader-Willi region.





650 kbp

6205 nm







2160 nm

480 nm




3D Architecture & Function of the IgH Locus

Fluorescence *in situ* hybridization of the IgH locus combined with spectral precision distance epifluorescence microscopy, analytical trilateration and comparison with computer simulations shows again a Multi-Loop Subcompartment organization of the IgH locus with functional relevant distances.






3D Architecture & Function of the IgH Locus

Fluorescence *in situ* hybridization of the IgH locus combined with spectral precision distance epifluorescence microscopy, analytical trilateration and comparison with computer simulations shows again a Multi-Loop Subcompartment organization of the IgH locus with functional relevant distances.




3D Architecture & Function of the IgH Locus

Fluorescence *in situ* hybridization of the IgH locus combined with spectral precision distance epifluorescence microscopy, analytical trilateration and comparison with computer simulations shows again a Multi-Loop Subcompartment organization of the IgH locus with functional relevant distances.



3D Architecture & Function of the IgH Locus

Fluorescence *in situ* hybridization of the IgH locus combined with spectral precision distance epifluorescence microscopy, analytical trilateration and comparison with computer simulations shows again a Multi-Loop Subcompartment organization of the IgH locus with functional relevant distances.





“Synoptic” 3D Architecture of Various Loci

A history “synoptic” comparison of the spatial distance mapping from their original background and aim, FISH methodological protocols, via microscopic imaging and restoration analysis procedures, to their interpretation, reveals that with time Multi-Loop Subcompartment models are favoured.



Study	Location	Preparation of Cells			FISH			Microscopy		Fit to model
		Cell cycle	KCl [nM]	Fixative	Melting	Label	Colours	# of nuclei	Image acquisition	
Fig. 3B, Trask '89	DHFR	UA41 G1-cf	75 dropped	MAA 3:1	FM 50 %	Biotin	1	20-37	photo, wall	RWGL 0.08-J RWGL 1.0
Fig. 3B, Lawrence '90	Dystrophin	WI38F G1	75 dropped	MAA 3:1	FM 50 %	Biotin	1	20-60	photo, wall	RWGL 0.5-1
Fig. 3A, Trask '91	Xq28	F G1-cf	75 dropped	MAA 3:1	FM 50 %	Biotin	1	30-60	photo, wall	RWGL 0.7 J RWGL 2.0- >5.0
Fig. 3B, Trask '91	Xq28	F G1-cf	75 dropped	MAA 3:1	FM 50 %	Biotin Dig	2	30-60	photo, wall	RWGL 1.0-3.0
Fig. 3, v.d. Engh '92 or Fig. 5A, Trask '93	4p16.3	F G1-cf	75 dropped	MAA 3:1	FM 50 %	Biotin Dig	2	?	photo, d-board	$L_S \leq 0.1$ for $GS < 0.5 <$ $RWGL > 5.0$
Fig. 5B, Trask '93	6p21	F G1-cf	75	MAA 3:1	FM 50-70 %	Biotin Dig	2	?	photo, d-board	$L_S \leq 0.1$ for $GS < 1.0 <$ $RWGL 1.0-5.0$
Fig. 5, Senger '93	MHC 6p21.31	HFF G1-cf	?	?	FM 50 %	Biotin	1	> 30	photo, wall	$MLS L_S \& LI_S =$ $0.12-0.25$ $RWGL 0.1-0.5$
Fig. 5, Senger '93	MHC 6p21.31	HFF G1-cf	?	?	FM 50 %	Biotin Dig	2	> 30	photo, wall	$MLS L_S = 0.1$ $LI_S = 0.18$ $RWGL 0.1-0.5$
Fig. 1, Warrington '94	4p16.3	F G1-cf	75	MAA 3:1	FM 50 %	Biotin Dig	2	?	?	$RWGL > 5.0$
Tab. 1, Warrington '94	5q31-33	L	?	?	?	?	?	?	CLSM BioRad	$RWGL > 5.0$
Fig. 2B, Yokota '95	4p16.3	F G1-cf	40 dropped	MAA 3:1	FM 70 %	Biotin Dig	2	40-360	photo, d-board	$RWGL 2.0-4.0$
Fig. 3B, Yokota '95	4p16.3	F G1-cf	-	PFA 4 %	FM 70 %	Biotin Dig	2	40-350	photo, d-board	$MLS L_S \& LI_S =$ $0.1-0.125$

Study	Location	Preparation of Cells			FISH			Microscopy		Fit to model
		Cell cycle	KCl [nM]	Fixative	Melting	Label	Colours	# of nuclei	Image acquisition	
Fig. 2A, Yokota '97	4p16.3 R-band	F G1-cf	40 dropped	MAA 3:1	FM 70 %	Biotin Dig	2	37-178	photo, d-board	$RWGL 2.0-3.0$
Fig. 2B, Yokota '97	6p21.3 R-band	F G1-o	40 dropped	MAA 3:1	FM 70 %	Biotin Dig	2	37-178	photo, d-board	$RWGL 4.0-5.0$
Fig. 2C, Yokota '97	21q22.2 G-band	F G1-o	40 dropped	MAA 3:1	FM 70 %	Biotin Dig	2	37-178	photo, d-board	$RWGL 1.0-2.0$
Fig. 2D, Yokota '97	Xp21.3 G-band	F G1-o	40 dropped	MAA 3:1	FM 70 %	Biotin Dig	2	37-178	photo, d-board	$RWGL 0.5-0.9$
Fig. 2D, Yokota '97	Xq28 R-band	F G1-of	40 dropped	MAA 3:1	FM 70 %	Biotin Dig	2	37-178	photo, d-board	$RWGL 1.0-5.0j$
Fig. 2A, Yokota '97	Xp21.3 G-band	F	-	PFA 4 %	FM 70 %	Biotin Dig	2	37-178	photo, d-board	$RWGL 0.25$ $MLS L_S = 0.126$ $LI_S = 0.200$
Fig. 2A, Yokota '97	Xq28 R-band	F	-	PFA 4 %	FM 70 %	Biotin Dig	2	37-178	photo, d-board	$RWGL 1.0$
Fig. 4B, Yokota '97	Xp21.3 G-band	HeLa	40 dropped	MAA 3:1	FM 70 %	Biotin Dig	2	37-178	photo, d-board	$RWGL 0.25$ $MLS L_S = 0.126$ $LI_S = 0.200$
Fig. 4B, Yokota '97	Xq28 R-band	HeLa	40 dropped	MAA 3:1	FM 70 %	Biotin Dig	2	37-178	photo, d-board	$RWGL 1.0$
Monier '97	11q13	F	-	PFA 4 %	FM 70 %	Biotin Dig	1	22-69	CLSM	$MLS L_S = 0.126$ $LI_S = 180$
Monier '97	11q13	L	-	PFA 4 %	FM 70 %	Biotin Dig	1	22-69	CLSM	$MLS L_S = 0.1$ $LI_S = 0.18-0.24$
Knoch '98/ Rauch '99	15q11-21	F	-	PFA 4 %	FM 70 %	Biotin Dig	1 & 2	60-120	CLSM	$MLS L_S = 0.1$ $LI_S = 0.06-0.125$







“Synoptic” 3D Architecture of Various Loci


A history “synoptic” comparison of the spatial distance mapping from their original background and aim, FISH methodological protocols, via microscopic imaging and restoration analysis procedures, to their interpretation, reveals that with time Multi-Loop Subcompartment models are favoured.



Study
Fig. 3B, Trask '89
Fig. 3B, Lawrence '91
Fig. 3A, Trask '91
Fig. 3B, Trask '91
Fig. 3, v.d. Trask '93
Fig. 5B, Trask '93
Fig. 5, Senger '93
Fig. 5, Senger '93
Fig. 1, Warrington
Tab. 1, Warrington
Fig. 2B, Ye '95
Fig. 3B, Ye '95





to model
L 2.0-3.0
L 4.0-5.0
L 1.0-2.0
L 0.5-0.9
L 1.0-5.0j
L 0.25 L _S =-0.126 0.200
L 1.0
L 0.25 L _S =-0.126 0.200
L 1.0
L _S =-0.126 180
L _S =-0.1 0.18-0.24
L _S =-0.1 0.06-0.125



From Fiber Topology to Nuclear Morphology

Chromosome territories form in the RW/GL and the MLS model. However, only the MLS model leads distinct subcompartments and low chromosome and subcompartment overlap. Best agreement is reached for an MLS model with 80 to 120 kbp loops and linkers in nuclei with 8 to 10 μm diameter.

The simulated nuclear morphology reflects the chromosome fiber topology of different models in detail.





A: MLS in 6 μm nucleus
 I: 63 kbp loops, 63 kbp linkers
 II: 63 kbp loops, 252 kbp linkers
 III: 126 kbp loops, 252 kbp linkers

B: MLS in 8 μm nucleus
 I: 126 kbp loops, 126 kbp linkers
 II: 84 kbp loops, 126 kbp linkers

C: MLS in 10 μm nucleus
 126 kbp loops, 126 kbp linker,
 not totally relaxed

D: RW/GL in 12 μm nucleus
 5 Mbp loops
 not totally relaxed



confocal microscopy
 100x objective, theoretic resolution

confocal microscopy
 63x objective, real resolution

confocal microscopy
 territory painting

rendering

electron microscopy

electron microscopy
 territory painting



In vivo Morphology & Chromatin Distribution

The stable expression of fusions between histones and autofluorescent proteins and the integration into nucleosomes allows the minimal invasive investigation of the structure and dynamics of chromatin.

The clustered morphology in detail favour an MLS like chromatin topology.





HeLa, H2A-YFP





Cos7, H1.0-GFP

LCLC 103H, H2A-CFP





ID13, H2A-YFP

HeLa, mH2A1.2-YFP





HeLa, H2A-YFP,
natural promoter




Fine Morphology of Nuclei

High resolution rendering and simulated electron microscopy including territory painting reveal not only again the model details but also that any location in the nucleus is accessible to biological molecules <15 nm in diameter and that even the Extended Interchromosomal Domain hypothesis is oversimplified.







MLS models with 126 kbp loops and linkers in a 10 μm nucleus.



Scaling of the Chromatin Fiber Topology


The spatial-distance and exact yard-stick dimension distinguish between the simulated models in detail. The MLS model shows a globular and fine-structured multi scaling behaviour due to the loops forming rosettes. This agrees with DNA fragmentation by Carbon ion irradiation and the appearance of fine-structured multi-scaling long-range correlations found in the sequential organization of genomes.




Scaling of the Chromatin Morphology & Distribution

The local (inverse-) mass dimension distinguishes between the models in detail and show also a multi-scaling behaviour with globular feature for the MLS model like the scaling of the fiber topology. With the mass dimension as function of intensity separates very well between different nuclei *in vivo*.




Consequently, the chromatin morphology is causally and quantitatively connected to the fiber topology.



(inverse-) mass dimension distribution







mass dimension as function of image intensity








Quantified TSA induced Morphology Changes

Trichostatin A induced histone acetylation can be quantified by *in vivo* H2A-GFP confocal images and image correlation spectroscopy (iFCS), which is a scaling analysis, and reveals the opening of chromatin, and thus reorganization changes on scales from 0.2 to $\sim 1\mu\text{m}$, consistent with MLS models.






Quantified TSA induced Morphology Changes

Trichostatin A induced histone acetylation can be quantified by *in vivo* H2A-GFP confocal images and image correlation spectroscopy (iFCS), which is a scaling analysis, and reveals the opening of chromatin, and thus reorganization changes on scales from 0.2 to $\sim 1\mu\text{m}$, consistent with MLS models.










Counting Nucleosomes *In Vivo*

Counting nucleosomes in living cells with a combination of fluorescence correlation spectroscopy and confocal laser scanning imaging reveals not only the free unbound histone component but also the concentration in absolute numbers of bound histones. Thus, the absolute concentration distribution of histones can be determined and reveals again the typical expected distribution of aggregated chromatin loops.



Counting Nucleosomes *In Vivo*

Counting nucleosomes in living cells with a combination of fluorescence correlation spectroscopy and confocal laser scanning imaging reveals not only the free unbound histone component but also the concentration in absolute numbers of bound histones. Thus, the absolute concentration distribution of histones can be determined and reveals again the typical expected distribution of aggregated chromatin loops.



Also The Dynamics Can Only Be Based On Quasi-Fibre Forming Rosettes !

Diffusion of Particles in the Nucleus





Due to the volume and spatial relationships in the nucleus typical particles reach almost any location in the nucleus by moderately obstructed diffusion: a 10 nm particle moves 1 to 2 μm within 10 ms.

The structural influence on the obstruction degree is random for Alexa 568 as function of the chromatin distribution visualized by H2A CFP in vivo and measured by fluorescence correlation spectroscopy (FCS)






$$\langle r^2 \rangle \propto t^{2/D_w}$$

Nuclear diameter [μm]	Nuclear Volume [μm^3]	Mean Nucleosome Concentration [μM]	Chromatin Volume Fraction [%]	Mean Isotropic Mesh Spacing [nm]
6	115	251	20.1	41
8	268	107	8.6	64
10	523	55	4.4	90
12	904	32	2.6	117





DNA Fragment Distribution after Ione-Irradiation

The length distribution of DNA fragments after irradiation with e. g. C or Ca with an inhomogeneous spatial double strand breakage probability depends on the detailed folding topology of the chromatin fiber and the RW/GL and MLS models differ largely.




Systems Biological Result Integration via the GLOBE 3D Genome Platform

All results will be integrated using our GLOBE 3D Genome Platform, established for analysis, manipulation and understanding of multi-dimensional complex genome wide data. Thus in reiterative cycles between experiments and simulations a systems biological/medical genome model will be achieved.






Systems Biological Result Integration via the GLOBE 3D Genome Platform

All results will be integrated using our GLOBE 3D Genome Platform, established for analysis, manipulation and understanding of multi-dimensional complex genome wide data. Thus in reiterative cycles between experiments and simulations a systems biological/medical genome model will be achieved.



Systems Biological Result Integration via the GLOBE 3D Genome Platform

All results will be integrated using our GLOBE 3D Genome Platform, established for analysis, manipulation and understanding of multi-dimensional complex genome wide data. Thus in reiterative cycles between experiments and simulations a systems biological/medical genome model will be achieved.





Conclusion

The compacted chromatin quasi-fibre, folds into loop-aggregates connected by a linker !

Every structural level of nuclear organization including its dynamics is connected and represented in all the other levels in a holistic systems genomics manner.



- The 3D genomes architecture consists of chromatin quasi-fibres (5 ± 1 nuc. / 11 nm, L_p of 80-120 nm), forming stable loop aggregates/rosettes (~ 40 -100 kbp loops, ~ 60 kbp linkers).
- The dynamics of genomes follows the 3D genome architecture in detail and determines in an inseparable entanglement with the architecture genome function.
- From the single base pair to the entire cell nucleus, all genomic levels represent all other levels and by modification a code is present and used to store genetic information.
- Genomes have a consensus organization with only small variation from the basic theme on each compaction level of the genome and these small variations determine genome function.
- Genome organization and function cannot be determined or understood from a single organizational level but only in a holistic systems genomics manner integrating all parts of the system.
- The genome behaves on the basis of a genomic statistical mechanics with a genomic uncertainty principle attached !



Acknowledgements

Thanks go to all the lab local lab members, those people who supported this work in the last decades, the institutions providing their infrastructure, and the national and international computing infrastructures.

Special thanks go to the reviewers, the EraSysBio Plus initiative and the national and EU funding bodies.



Nick Kepper
Michael Lesnussa
Anis Abuseiris

Petros Kolovos
Jessica Zuin

Harmen J. G. van de Werken
A.M. Ali Imam



Christel E. M. Kockx
Rutger W. W. Brouwer
Wilfred F. J. van Ijken

Kerstin S. Wendt
Frank G. Grosveld


Cell Biology & Biophysics

EMBL

Malte Wachsmuth



Acknowledgements




	Biophysical Genomics, Cell Biology, Erasmus MC	Biological Sciences, UCSD	The Cremer Labs	
		Suchit Jhunjhunwala	Joachim Rauch	
Biophysics, BTZ Dresden	Petros Kolovos	Menno van Zelm	Irina Solovei	Biophysics of Macromolecules DKFZ
Thomas Weidemann	Anis Abuseiris	Cornelis Murre	Michael Hausmann	Gabriele Müller
	Michael Lesnussa		Christoph Cremer	Waldemar Waldeck
Cell Biology & Biophysics	Rob de Graaf	Clinical Genetics Erasmus MC	Thomas Cremer	Jörg Langowski
EMBL	Nick Kepper	Bert Eussen		
Malte Wachsmuth	Frank Grossveld	Annelies de Klein	Molecular Genetics DKFZ	Supercomputing Center Karlsruhe
Molecular Biophysics KIP	LMU Munich	University Tübingen	Karsten Richter	Rudolph Lohner
Katalin Fejes-Toth	Peter Quicken	Markus Göker	Peter Lichter	
Karsten Rippe	Anna Friedl			

Erasmus Medical Center and BioQuant & German Cancer Research Center

High-Performance Computing Center Stuttgart, University of Stuttgart; Supercomputing Center Karlsruhe, University of Karlsruhe; Computing Center, Deutsches Krebsforschungszentrum Heidelberg (DKFZ)

Erasmus Medical Center, Hogeschool Rotterdam, The Fraunhofer Society, The German MediGRID and Services@MediGRID, The German D-Grid Initiatives, The German Ministry for Science and Technology, The Dutch Science Organization (NOW), The European EGEE Initiative, The European EDGES Consortium, The German Society for Human Ecology, The International Society for Human Ecology, The European Commission



Acknowledgements

Biophysical Genomics, Biological Sciences, UCSD The Cremer I
Cell Biology, Erasmus MC Suchit Jhunjhunwala Joachim

Biophysics, BTZ Dresden Petros Kolovos Menno van Zelm
Thomas Weidemann Anis Abuseiris Cornelis Murre

Cell Biology & Biophysics Michael Lesnussa
EMBL Rob de Graaf
Malte Wachsmuth Nick Kepper
Frank Grossveld

Molecular Biophysics LMU M Petrus
KIP Katalin Fejes-Toth
Karsten Rippe

Center and BioQuant & German Cancer Research Center
High Performance Computing Center, University of Stuttgart; Supercomputing Center Karlsruhe, University of
Computing Center, Deutsches Krebsforschungszentrum Heidelberg (DKFZ)


University of Amsterdam, The Netherlands; Eindhoven University of Technology, The Netherlands; Radboud University Nijmegen, The Netherlands; The Fraunhofer Society, The German MediGRID and Services@MediGRID,
Research Initiatives, The German Ministry for Science and Technology, The Dutch Science Organization (NOW),
the EGEE Initiative, The European EDGES Consortium, The German Society for Human Ecology, The
International Society for Human Ecology,
The European Commission

Macromolecules
DKFZ
Gabriele Müller
Waldemar Waldeck
Jörg Langowski


Molecular Genetics
DKFZ
Karsten Richter
Peter Lichter
Rudolph Lohner

Supercomputing Center
Karlsruhe

The Real Fun
Is Yet To Come!




15
 Decipher
 Affy 100KXba
 Refseqb36



43750000

15

1:500000



The
Detailed 3D Multi-Loop Aggregate/Rosette Chromatin Architecture
and
Functional Dynamic Organization
of the
Human and Mouse Genome

Knoch, T. A.,

Wachsmuth, M., Kepper, N., Lesnussa, M., Abuseiris, A., Kolovos, P., Zuin, J., van den Werken, H. J. G., Wendt, K. S. & Grosveld, F. G.

Chromatin and Epigenetics: From Omics to Single Cells 2014, Institut de Genetique et de Biologie Moleculaire et Cellulaire (IGBMC), Strasbourg, France, 14th - 15th October, 2014.

Abstract

The dynamic three-dimensional chromatin architecture of genomes and the obvious co-evolutionary connection to its function – the storage and expression of genetic information – is still, after ~170 years of concentrated research, one of the central issues of our time. With a systems genomics combination of T2C and FCS together with analytical and simulated polymer models as well as scaling analysis of the architecture and the DNA sequence itself, we determined the architecture and dynamics with molecular resolution from the single base to the mega base pair level! T2C is a novel superior selective high-throughput high-resolution chromosomal interaction capture of all physical genomic interactions (T2C) which has great advantages since it allows to reach an optimal combination of resolution, interaction frequency range, and multi-plexability, at unprecedented signal-to-noise ratio of multiple orders of magnitude. Actually, with T2C the ultimate limits at the level of the “genomic uncertainty principle” and “genomic statistical mechanics”, is reached. In parallel, with a novel FCS approach it is possible to determine the relaxation of local concentration fluctuations of the chromatin fibre and derive based on an analytical polymer model the relation between average persistence length, mass density and chromatin architecture. Whereas T2C measures architectural parameters, the FCS approach allows to measure *in vivo* the dynamics of the architecture. As common ground polymer simulations using Monte Carlo and Brownian Dynamics approaches as well as scaling analysis of the architecture as well as the DNA sequence organization itself round-off the experimental techniques in detail. In combination all four approaches reach agreement concerning the 3D architecture and dynamics of genomes in detail. With this we investigated several loci in detail of different species, cell type, and functional states from *some to the mega-base pair level and thus spanning 6 orders of magnitude (!)*, and finally determined the three-dimensional organization and dynamics for the first time in a consistent system genomics manner from the four mentioned angles. Beyond, the results are also in agreement with the heuristics of the research of the last 170 years. Consequently, we have not only finally determined the architecture and dynamics of the mouse and human genome with base pair/molecular resolution, but this also opens now the detailed architectural and dynamic sequencing of genomes in a systems genomic manner which is of fundamental importance from fundamental genome understanding to diagnosis and treatment.

Corresponding author email contact: TA.Knoch@taknoch.org

Keywords:

Genome, genomics, genome organization, genome architecture, structural sequencing, architectural sequencing, systems genomics, coevolution, holistic genetics, genome mechanics, genome statistical mechanics, genomic uncertainty principle, multilism genotype-phenotype, genome function, genetics, gene regulation, replication, transcription, repair, homologous recombination, simultaneous co-transfection, cell division, mitosis, metaphase, interphase, cell nucleus, nuclear structure, nuclear organization, chromatin density distribution, nuclear morphology, chromosome territories, subchromosomal domains, chromatin loop aggregates, chromatin rosettes, chromatin loops, chromatin quasi fibre, chromatin density, persistence length, spatial distance measurement, histones, H1.0, H2A, H2B, H3, H4, mH2A1.2, DNA sequence, complete sequenced genomes, molecular transport, obstructed diffusion, anomalous diffusion, percolation, long-range correlations, fractal analysis, scaling analysis, exact yard-stick dimension, box-counting dimension, lacunarity dimension, local nuclear dimension, nuclear diffuseness, parallel super computing, grid computing, volunteer computing, polymer model, analytic mathematical model, Brownian Dynamics, Monte Carlo, fluorescence *in situ* hybridization (FISH), targeted chromatin capture (T2C) confocal laser scanning microscopy, fluorescence correlation spectroscopy, spatial precision distance microscopy, super-resolution microscopy, two dimensional fluorescence correlations spectroscopy (2D-FCS) auto-fluorescent proteins, CFP, GFP, YFP, DsRed, fusion protein, *in vivo* labelling, information browser, visual data base access, holistic viewing system, integrative data management, extreme visualization, three-dimensional virtual environment, virtual paper tool, human ecology, e-human grid ecology, society, social systems, e-social challenge, inverse tragedy of the commons, grid phenomenon, micro-sociality, macro-sociality, autopoietic tragedy of social sub-systems, micro subsystems, macro subsystems, micro operationality, macro operationality, grid psychology micro riskmanagement, macro riskmanagement.

Literature References

- Knoch, T. A.** Dreidimensionale Organisation von Chromosomen-Domänen in Simulation und Experiment. (Three-dimensional organization of chromosome domains in simulation and experiment.) *Diploma Thesis*, Faculty for Physics and Astronomy, Ruperto-Carola University, Heidelberg, Germany, 1998, and TAK Press, Tobias A. Knoch, Mannheim, Germany, ISBN 3-00-010685-5 and ISBN 978-3-00-010685-9 (soft cover, 2rd ed.), ISBN 3-00-035857-9 and ISBN 978-3-00-035885-0 (hard cover, 2rd ed.), ISBN 3-00-035858-7, and ISBN 978-3-00-035858-6 (DVD, 2rd ed.), 1998.
- Knoch, T. A., Münkkel, C. & Langowski, J.** Three-dimensional organization of chromosome territories and the human cell nucleus - about the structure of a self replicating nano fabrication site. *Foresight Institute - Article Archive*, Foresight Institute, Palo Alto, CA, USA, <http://www.foresight.org>, 1- 6, 1998.
- Knoch, T. A., Münkkel, C. & Langowski, J.** Three-Dimensional Organization of Chromosome Territories and the Human Interphase Nucleus. *High Performance Scientific Supercomputing*, editor Wilfried Juling, Scientific Supercomputing Center (SSC) Karlsruhe, University of Karlsruhe (TH), 27- 29, 1999.
- Knoch, T. A., Münkkel, C. & Langowski, J.** Three-dimensional organization of chromosome territories in the human interphase nucleus. *High Performance Computing in Science and Engineering 1999*, editors Krause, E. & Jäger, W., High-Performance Computing Center (HLRS) Stuttgart, University of Stuttgart, Springer Berlin-Heidelberg-New York, ISBN 3-540-66504-8, 229-238, 2000.
- Bestvater, F., **Knoch, T. A.**, Langowski, J. & Spiess, E. GFP-Walking: Artificial construct conversions caused by simultaneous cotransfection. *BioTechniques* 32(4), 844-854, 2002.
- Gil-Parado, S., Fernández-Montalván, A., Assfalg-Machleidt, I., Popp, O., Bestvater, F., Holloschi, A., **Knoch, T. A.**, Auerswald, E. A., Welsh, K., Reed, J. C., Fritz, H., Fuentes-Prior, P., Spiess, E., Salvesen, G. & Machleidt, W. Ionomycin-activated calpain triggers apoptosis: A probable role for Bcl-2 family members. *J. Biol. Chem.* 277(30), 27217-27226, 2002.
- Knoch, T. A. (editor)**, Backes, M., Baumgärtner, V., Eysel, G., Fehrenbach, H., Göker, M., Hampl, J., Hampl, U., Hartmann, D., Hitzelberger, H., Nambena, J., Rehberg, U., Schmidt, S., Weber, A., & Weidemann, T. Humanökologische Perspektiven Wechsel - Festschrift zu Ehren des 70. Geburtstags von Prof. Dr. Kurt Egger. Human Ecology Working Group, Ruperto-Carola University of Heidelberg, Heidelberg, Germany, 2002.

- Knoch, T. A.** Approaching the three-dimensional organization of the human genome: structural-, scaling- and dynamic properties in the simulation of interphase chromosomes and cell nuclei, long-range correlations in complete genomes, *in vivo* quantification of the chromatin distribution, construct conversions in simultaneous co-transfections. *Dissertation*, Ruperto-Carola University, Heidelberg, Germany, and TAK†Press, Tobias A. Knoch, Mannheim, Germany, ISBN 3-00-009959-X and ISBN 978-3-00-009959-5 (soft cover, 3rd ed.), ISBN 3-00-009960-3 and ISBN 978-3-00-009960-1 (hard cover, 3rd ed.), ISBN 3-00-035856-9 and ISBN 978-3-00-010685-9 (DVD, 3rd ed.) 2002.
- Westphal, G., van den Berg-Stein, S., Braun, K., **Knoch, T. A.**, Dümmerling, M., Langowski, J., Debus, J. & Friedrich, E. Detection of the NGF receptors TrkaA and p75NTR and effect of NGF on the growth characteristics of human tumor cell lines. *J. Exp. Clin. Canc. Res.* 21(2), 255-267, 2002.
- Westphal, G., Niederberger, E., Blum, C., Wollman, Y., **Knoch, T. A.**, Dümmerling, M., Rebel, W., Debus, J. & Friedrich, E. Erythropoietin Receptor in Human Tumor Cells: Expression and Aspects Regarding Functionality. *Tumori* 88(2), 150-159, 2002.
- Gil-Parado, S., Popp, O., **Knoch, T. A.**, Zahler, S., Bestvater, F., Felgenträger, M., Holoshi, A., Fernández-Montalván, A., Auerswald, E. A., Fritz, H., Fuentes-Prior, P., Machleidt, W. & Spiess, E. Subcellular localization and subunit interactions of over-expressed human μ -calpain. *J. Biol. Chem.* 278(18), 16336-15346, 2003.
- Knoch, T. A.** Towards a holistic understanding of the human genome by determination and integration of its sequential and three-dimensional organization. *High Performance Computing in Science and Engineering 2003*, editors Krause, E., Jäger, W. & Resch, M., High-Performance Computing Center (HLRS) Stuttgart, University of Stuttgart, Springer Berlin-Heidelberg-New York, ISBN 3- 540-40850-9, 421-440, 2003.
- Wachsmuth, M., Weidemann, T., Müller, G., Urs W. Hoffmann-Rohrer, **Knoch, T. A.**, Waldeck, W. & Langowski, J. Analyzing intracellular binding and diffusion with continuous fluorescence photobleaching. *Biophys. J.* 84(5), 3353-3363, 2003.
- Weidemann, T., Wachsmuth, M., **Knoch, T. A.**, Müller, G., Waldeck, W. & Langowski, J. Counting nucleosomes in living cells with a combination of fluorescence correlation spectroscopy and confocal imaging. *J. Mol. Biol.* 334(2), 229-240, 2003.
- Fejes Tóth, K., **Knoch, T. A.**, Wachsmuth, M., Frank-Stöhr, M., Stöhr, M., Bacher, C. P., Müller, G. & Rippe, K. Trichostatin A induced histone acetylation causes decondensation of interphase chromatin. *J. Cell Science* 117, 4277-4287, 2004.
- Ermler, S., Kronic, D., **Knoch, T. A.**, Moshir, S., Mai, S., Greulich-Bode, K. M. & Boukamp, P. Cell cycle-dependent 3D distribution of telomeres and telomere repeat-binding factor 2 (TRF2) in HaCaT and HaCaT-myc cells. *Europ. J. Cell Biol.* 83(11-12), 681-690, 2004.
- Kost, C., Gama de Oliveira, E., **Knoch, T. A.** & Wirth, R. Spatio-temporal permanence and plasticity of foraging trails in young and mature leaf-cutting ant colonies (*Atta spp.*). *J. Trop. Ecol.* 21(6), 677- 688, 2005.
- Winnefeld, M., Grewenig, A., Schnölzer, M., Spring, H., **Knoch, T. A.**, Gan, E. C., Rommelaere, J. & Cziepluch, C. Human SGT interacts with BAG-6/Bat-3/Scythe and cells with reduced levels of either protein display persistence of few misaligned chromosomes and mitotic arrest. *Exp. Cell Res.* 312, 2500-2514, 2006.
- Sax, U., Weisbecker, A., Falkner, J., Viezens, F., Yassene, M., Hartung, M., Bart, J., Krefting, D., **Knoch, T. A.** & Semler, S. Grid-basierte Services für die elektronische Patientenakte der Zukunft. *E- HEALTH-COM - Magazin für Gesundheitstelematik und Telemedizin*, 4(2), 61-63, 2007.
- de Zeeuw, L. V., **Knoch, T. A.**, van den Berg, J. & Grosveld, F. G. Erasmus Computing Grid - Het bouwen van een 20 TeraFLOP virtuele supercomputer. *NIOC proceedings 2007 - het perspectief of lange termijn*. editor Frederik, H. NIOC, Amsterdam, The Netherlands, 52-59, 2007.
- Rauch, J., **Knoch, T. A.**, Solovei, I., Teller, K. Stein, S., Buiting, K., Horsthemke, B., Langowski, J., Cremer, T., Hausmann, M. & Cremer, C. Lightoptical precision measurements of the Prader- Willi/Angelman Syndrome imprinting locus in human cell nuclei indicate maximum condensation changes in the few hundred nanometer range. *Differentiation* 76(1), 66-82, 2008.
- Sax, U., Weisbecker, A., Falkner, J., Viezens, F., Mohammed, Y., Hartung, M., Bart, J., Krefting, D., **Knoch, T. A.** & Semler, S. C. Auf dem Weg zur individualisierten Medizin - Grid-basierte Services für die EPA der Zukunft. *Telemedizinführer Deutschland 2008*, editor Jäckel, A. Deutsches Medizinforum, Minerva KG, Darmstadt, ISBN 3-937948-06-6, ISBN-13 9783937948065, 47-51, 2008.

- Drägestein, K. A., van Capellen, W. A., van Haren, J. Tsibidis, G. D., Akhmanova, A., **Knoch, T. A.**, Grosveld, F. G. & Galjart, N. Dynamic behavior of GFP-CLIP-170 reveals fast protein turnover on microtubule plus ends. *J. Cell Biol.* 180(4), 729-737, 2008.
- Jhunjhunwala, S., van Zelm, M. C., Peak, M. M., Cutchin, S., Riblet, R., van Dongen, J. J. M., Grosveld, F. G., **Knoch, T. A.**⁺ & Murre, C.⁺ The 3D-structure of the Immunoglobulin Heavy Chain Locus: implications for long-range genomic interactions. *Cell* 133(2), 265-279, 2008.
- Krefting, D., Bart, J., Beronov, K., Dzhimova, O., Falkner, J., Hartung, M., Hoheisel, A., **Knoch, T. A.**, Lingner, T., Mohammed, Y., Peter, K., Rahm, E., Sax, U., Sommerfeld, D., Steinke, T., Tolxdorff, T., Vossberg, M., Viezens, F. & Weisbecker, A. MediGRID - Towards a user friendly secured grid infrastructure. *Future Generation Computer Systems* 25(3), 326-336, 2008.
- Knoch, T. A.**, Lesnussa, M., Kepper, F. N., Eussen, H. B., & Grosveld, F. G. The GLOBE 3D Genome Platform - Towards a novel system-biological paper tool to integrate the huge complexity of genome organization and function. *Stud. Health. Technol. Inform.* 147, 105-116, 2009.
- Knoch, T. A.**, Baumgärtner, V., de Zeeuw, L. V., Grosveld, F. G., & Egger, K. e-Human Grid Ecology: Understanding and approaching the Inverse Tragedy of the Commons in the e-Grid Society. *Stud. Health. Technol. Inform.* 147, 269-276, 2009.
- Dickmann, F., Kaspar, M., Löhnardt, B., **Knoch, T. A.**, & Sax, U. Perspectives of MediGRID. *Stud. Health. Technol. Inform.* 147, 173-182, 2009.
- Knoch, T. A.**, Göcker, M., Lohner, R., Abuseiris, A. & Grosveld, F. G. Fine-structured multi-scaling long-range correlations in completely sequenced genomes - features, origin and classification. *Eur. Biophys. J.* 38(6), 757-779, 2009.
- Dickmann, F., Kaspar, M., Löhnardt, B., Kepper, N., Viezens, F., Hertel, F., Lesnussa, M., Mohammed, Y., Thiel, A., Steinke, T., Bernarding, J., Krefting, D., **Knoch, T. A.** & Sax, U. Visualization in health-grid environments: a novel service and business approach. *LNCS 5745*, 150-159, 2009.
- Dickmann, F., Kaspar, M., Löhnardt, B., Kepper, N., Viezens, F., Hertel, F., Lesnussa, M., Mohammed, Y., Thiel, A., Steinke, T., Bernarding, J., Krefting, D., **Knoch, T. A.** & Sax, U. Visualization in health-grid environments: a novel service and business approach. *Grid economics and business models - GECON 2009 Proceedings, 6th international workshop, Delft, The Netherlands.* editors Altmann, J., Buyya, R. & Rana, O. F., GECON 2009, LNCS 5745, Springer-Verlag Berlin Heidelberg, ISBN 978-3-642-03863-1, 150-159, 2009.
- Estrada, K.* , Abuseiris, A.* , Grosveld, F. G., Uitterlinden, A. G., **Knoch, T. A.**⁺ & Rivadeneira, F.⁺ GRIMP: A web- and grid-based tool for high-speed analysis of large-scale genome-wide association using imputed data. *Bioinformatics* 25(20), 2750-2752, 2009.
- de Wit, T., Dekker, S., Maas, A., Breedveld, G., **Knoch, T. A.**, Langeveld, A., Szumska, D., Craig, R., Bhattacharya, S., Grosveld, F. G.⁺ & Drabek, D. Tagged mutagenesis of efficient minos based germ line transposition. *Mol. Cell Biol* 30(1), 66-77, 2010.
- Kepper, N., Schmitt, E., Lesnussa, M., Weiland, Y., Eussen, H. B., Grosveld, F. G., Hausmann, M. & **Knoch T. A.**, Visualization, Analysis, and Design of COMBO-FISH Probes in the Grid-Based GLOBE 3D Genome Platform. *Stud. Health Technol. Inform.* 159, 171-180, 2010.
- Kepper, N., Ettig, R., Dickmann, F., Stehr, R., Grosveld, F. G., Wedemann, G. & **Knoch, T. A.** Parallel high-performance grid computing: capabilities and opportunities of a novel demanding service and business class allowing highest resource efficiency. *Stud. Health Technol. Inform.* 159, 264-271, 2010.
- Skrownny, D., Dickmann, F., Löhnardt, B., **Knoch, T. A.** & Sax, U. Development of an information platform for new grid users in the biomedical field. *Stud. Health Technol. Inform.* 159, 277-282, 2010.
- Knoch, T. A.**, Baumgärtner, V., Grosveld, F. G. & Egger, K. Approaching the internalization challenge of grid technologies into e-Society by e-Human "Grid" Ecology. *Economics of Grids, Clouds, Systems, and Services – GECON 2010 Proceedings, 7th International Workshop, Ischia, Italy,* editors Altman, J., & Rana, O. F., Lecture Notes in Computer Science (LNCS) 6296, Springer Berlin Heidelberg New York, ISSN 0302-9743, ISBN-10 3-642-15680-0, ISBN-13 978-3-642-15680-9, 116-128, 2010.
- Dickmann, F., Brodhun, M., Falkner, J., **Knoch, T. A.** & Sax, U. Technology transfer of dynamic IT outsourcing requires security measures in SLAs. *Economics of Grids, Clouds, Systems, and Services – GECON 2010 Proceedings, 7th International Workshop, Ischia, Italy,* editors Altman, J., & Rana, O. F., Lecture Notes in Computer Science (LNCS) 6296, Springer Berlin Heidelberg New York, ISSN 0302-9743, ISBN-10 3-642-15680-0, ISBN-13 978-3-642-15680-9, 1-115, 2010.

- Knoch, T. A.** Sustained Renewability: approached by systems theory and human ecology. *Renewable Energy 2*, editors M. Nayeripour & M. Keshti, Intech, ISBN 978-953-307-573-0, 21-48, 2011.
- Kolovos, P., **Knoch, T. A.**, F. G. Grosveld, P. R. Cook, & Papantonis, A. Enhancers and silencers: an integrated and simple model for their function. *Epigenetics and Chromatin 5(1)*, 1-8, 2012.
- Dickmann, F., Falkner, J., Gunia, W., Hampe, J., Hausmann, M., Herrmann, A., Kepper, N., **Knoch, T. A.**, Lauterbach, S., Lippert, J., Peter, K., Schmitt, E., Schwardmann, U., Solodenko, J., Sommerfeld, D., Steinke, T., Weisbecker, A. & Sax, U. Solutions for Biomedical Grid Computing - Case Studies from the D-Grid Project Services@MediGRID. *JOCS 3(5)*, 280-297, 2012.
- Estrada, K., Abuseiris, A., Grosveld, F. G., Uitterlinden, A. G., **Knoch, T. A.** & Rivadeneira, F. GRIMP: A web- and grid-based tool for high-speed analysis of large-scale genome-wide association using imputed data. *Dissection of the complex genetic architecture of human stature and osteoporosis*. cumulative dissertation, editor Estrada K., Erasmus Medical Center, Erasmus University Rotterdam, Rotterdam, The Netherlands, ISBN 978-94-6169-246-7, 25-30, 1st June 2012.
- van de Corput, M. P. C., de Boer, E., **Knoch, T. A.**, van Cappellen, W. A., Quintanilla, A., Ferrand, L., & Grosveld, F. G. Super-resolution imaging reveals 3D folding dynamics of the β -globin locus upon gene activation. *J. Cell Sci. 125 (Pt 19)*, 4630-4639, 2012.
- da Silva, P. S. D., Delgado Bieber, A. G., Leal, I. R., **Knoch, T. A.**, Tabarelli, M., Leal, I. R., & Wirth, R. Foraging in highly dynamic environments: leaf-cutting ants adjust foraging trail networks to pioneer plant availability. *Entomologia Experimentalis et Applicata 147*, 110-119. 2013.
- Zuin, J., Dixon, J. R., van der Reijden, M. I. J. A., Ye, Z., Kolovos, P., Brouwer, R. W. W., van de Corput, M. P. C., van de Werken, H. J. G., **Knoch, T. A.**, van IJcken, W. F. J., Grosveld, F. G., Ren, B. & Wendt, K. S. Cohesin and CTCF differentially affect chromatin architecture and gene expression in human cells. *PNAS 111(3)*, 9906-1001, 2014.
- Kolovos, P., Kepper, N., van den Werken, H. J. G., Lesnussa, M., Zuin, J., Brouwer, R. W. W., Kockx, C. E. M., van IJcken, W. F. J., Grosveld, F. G. & **Knoch, T. A.** Targeted Chromatin Capture (T2C): A novel high resolution high throughput method to detect genomic interactions and regulatory elements. *Epigenetics & Chromatin 7:10*, 1-17, 2014.
- Diermeier, S., Kolovos, P., Heizinger, L., Schwartz, U., Georgomanolis, T., Zirkel, A., Wedemann, G., Grosveld, F. G., **Knoch, T. A.**, Merkl, R., Cook, P. R., Längst, G. & Papantonis, A. TNF α signalling primes chromatin for NF- κ B binding and induces rapid and widespread nucleosome repositioning. *Genome Biology 15(12)*, 536-548, 2014.
- Knoch, T. A.**, Wachsmuth, M., Kepper, N., Lesnussa, M., Abuseiris, A., A. M. Ali Imam, Kolovos, P., Zuin, J., Kockx, C. E. M., Brouwer, R. W. W., van de Werken, H. J. G., van IJcken, W. F. J., Wendt, K. S. & Grosveld, F. G. The detailed 3D multi-loop aggregate/rosette chromatin architecture and functional dynamic organization of the human and mouse genomes. *bioRxiv preprint*, 16.07.2016.
- Kolovos, P., Georgomanolis, T., Koeflerle, A., Larkin, J. D., Brant, J., Nikolić, M., Gusmao, E. G., Zirkel, A., **Knoch, T. A.**, van IJcken, W. F. J., Cook, P. R., Costa, I. G., Grosveld, F. G. & Papantonis, A. Binding of nuclear kappa-B to non-canonical consensus sites reveals its multimodal role during the early inflammatory response. *Genome Research 26(11)*, 1478-1489, 2016.
- Wachsmuth, M., **Knoch, T. A.** & Rippe, K. Dynamic properties of independent chromatin domains measured by correlation spectroscopy in living cells. *Epigenetics & Chromatin 9:57*, 1-20, 2016.
- Knoch, T. A.**, Wachsmuth, M., Kepper, N., Lesnussa, M., Abuseiris, A., A. M. Ali Imam, Kolovos, P., Zuin, J., Kockx, C. E. M., Brouwer, R. W. W., van de Werken, H. J. G., van IJcken, W. F. J., Wendt, K. S. & Grosveld, F. G. The detailed 3D multi-loop aggregate/rosette chromatin architecture and functional dynamic organization of the human and mouse genomes. *Epigenetics & Chromatin 9:58*, 1-22, 2016.